



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.2653867>Available online at: <http://www.iajps.com>

Research Article

**ANALYSIS OF ABDOMINAL ULTRASONOGRAPHY FOR
PREDICTION OF SCAR DEHISCENCE IN FEMALES WITH
PREVIOUS CESAREAN SECTION**Dr M. Faizan Arshad¹, Dr Fawad Ashraf², Dr Urooj Iqbal³¹Tehsil Headquarter Hospital Shorkot, Jhang²Ittefaq Hospital, Lahore³District Head Quarter Hospital, Hafizabad

Article Received: February 2019

Accepted: March 2019

Published: April 2019

Abstract:

Introduction: Cesarean section (CS) rates are increasing worldwide. As a result, women presenting with pregnancy with previous CS are also rising. **Objectives of the study:** The basic objective of the study is to analyze the abdominal ultrasonography for prediction of scar dehiscence in females with previous cesarean section. **Methodology of the study:** This cross-sectional study was conducted in THQ hospital Shorkot, Jhang from March 2018 to December 2018. The data was collected from 150 patients. Females of age: 20 - 40 years with parity <5, presenting at gestational age >37 weeks with history of previous one delivery through cesarean section and planned to undergo delivery through elective cesarean section under spinal anesthesia were included in this study. The data was collected through a questionnaire. Demographic information (name, age, gestational age, BMI, parity) was also obtained. Then females undergo transabdominal ultrasonography by a single senior radiologist. Females were labeled as positive or negative for scar dehiscence. At time of delivery, uterine scar was assessed and scar dehiscence was confirmed. **Results:** The data were collected from 160 patients. The mean age of the patients was 30.22±6.33 years with minimum and maximum ages of 20 & 40 years respectively. The mean value of gestational age of the patients was 38.81±0.740 weeks with minimum and maximum gestational ages of 38 & 40 weeks respectively. The mean value of BMI of the patients was 24.22±3.321 kg/m² with minimum and maximum BMI values of 18.60 & 29.83 kg/m² respectively. The patients with age ≤ 30 years, the sensitivity, specificity and diagnostic accuracy of scar dehiscence on USG was 93.75%, 97.78% & 96.10% respectively taking cesarean section as gold standard. **Conclusion:** It is concluded that trans abdominal ultrasonography is a useful and reliable method having high value of diagnostic accuracy for prediction of scar dehiscence in females with history of cesarean.

Corresponding author:

Dr. M. Faizan Arshad,

Tehsil Headquarter Hospital Shorkot, Jhang

QR code



Please cite this article in press M. Faizan Arshad et al., *Analysis Of Abdominal Ultrasonography For Prediction Of Scar Dehiscence In Females With Previous Cesarean Section.*, Indo Am. J. P. Sci, 2019; 06(04).

INTRODUCTION:

Cesarean section (CS) rates are increasing worldwide. As a result, women presenting with pregnancy with previous CS are also rising. Previous CS is becoming the most common indication for CS, confirming the age-old dictum proposed by Edward Craigin in 1914 "Once a cesarean always a cesarean." Although the absolute risk of uterine dehiscence/rupture in lower segment CS is very low (0.2–1.5 %), the unpredictable nature of this complication and its grave consequences for both mother and baby has resulted in decreased rates of trial of labor after CS (TOLAC) in many countries [1].

The main cause of uterine rupture in a scarred uterus is lack of appropriate counseling and inadequate or absence of antenatal care with increasing number of women undergoing trial of labor after a previous caesarean section, in an anticipation of vaginal delivery [2], separation of previous caesarean scar has become a common cause of rupture especially in unskilled hands [3].

One study showed that frequency of scar dehiscence present in 69% cases after previous cesarean section. The frequency of ruptured uteri was calculated to be 0.67%, giving a ratio of 1:148 deliveries [4]. Ultrasound estimation of LUS provides a fairly simple and non-invasive method for prediction of scar dehiscence/rupture. Evaluation of thickness of LUS has been found to be a potential factor for predicting scar dehiscence. The risk of scar dehiscence/rupture has been directly related to the thinning of LUS. However, there is controversy over the thickness of LUS. One study showed that the sensitivity and specificity of Trans abdominal Sonography or trans abdominal ultrasonography were 91% and 93% respectively for prediction of scar dehiscence [5]. But another study showed that trans abdominal ultrasonography showed the sensitivity 25%, specificity 100% for prediction of scar dehiscence [6].

Objectives of the study

The basic objective of the study is to analyze the abdominal ultrasonography for prediction of scar dehiscence in females with previous cesarean section.

METHODOLOGY OF THE STUDY:

This cross-sectional study was conducted in THQ hospital Shorkot, Jhang from March 2018 to December 2018. The data was collected from 150 patients. Females of age: 20 - 40 years with parity <5, presenting at gestational age >37 weeks with history of previous one delivery through cesarean section and planned to undergo delivery through elective cesarean section under spinal anesthesia were included in this study. The data was collected through a questionnaire. Demographic information (name, age, gestational age, BMI, parity) was also obtained. Then females undergo transabdominal ultrasonography by a single senior radiologist. Females were labeled as positive or negative for scar dehiscence. At time of delivery, uterine scar was assessed and scar dehiscence was confirmed.

Statistical analysis

All the data were analyzed using SPSS version 18.0. All the values were expressed in mean and standard deviation.

RESULTS:

The data were collected from 160 patients. The mean age of the patients was 30.22 ± 6.33 years with minimum and maximum ages of 20 & 40 years respectively. In this study, 21(13.13%) patients had parity 01, 62(38.75%) patients had parity 02, 53(33.13%) patients had parity 03 and 24(15%) patients had parity 04. The mean value of gestational age of the patients was 38.81 ± 0.740 weeks with minimum and maximum gestational ages of 38 & 40 weeks respectively. The mean value of BMI of the patients was 24.22 ± 3.321 kg/m² with minimum and maximum BMI values of 18.60 & 29.83 kg/m² respectively. According to this study in patients with age ≤ 30 years, the sensitivity, specificity and diagnostic accuracy of scar dehiscence on USG was 93.75%, 97.78% & 96.10% respectively taking cesarean section as gold standard.

Table 01: Frequency distribution of scar dehiscence diagnosed on USG

		Frequency	Percent
Scar dehiscence USG	Positive	68	42.5
	Negative	92	57.5
	Total	160	100.0

Table 01: Comparison of diagnosis of scar dehiscence on USG with cesarean stratified by age

Age (years)	Scar dehiscence on USG	Scar dehiscence on cesarean		Total
		Positive	Negative	
≤ 30	Positive	30(93.8%)	1(2.2%)	31(40.3%)
	Negative	2(6.3%)	44(97.8%)	46(59.7%)
> 30	Positive	33(94.3%)	4(8.3%)	37(44.6%)
	Negative	2(5.7%)	44(91.7%)	46(55.4%)

DISCUSSION:

Cesarean section rates are increasing worldwide. As a result, women presenting with pregnancy with previous Caesarean Section are also rising [7]. Previous Caesarean Section is becoming the most common indication for Caesarean Section, confirming the age old dictum proposed by Edward Craigin in 1914 "Once a cesarean always a cesarean". In this study the sensitivity, specificity and diagnostic accuracy transabdominal ultrasonography for prediction of scar dehiscence in females with history of cesarean section was 94.03%, 94.62% & 94.38% respectively taking cesarean section as gold standard. Some of the studies are discussed below showing their results as [8].

A study by Chanderdeep Sharma et al concluded that sonographic evaluation of LUS scar and myometrial thickness (both with trans abdominal and vaginal ultrasonography) is a safe, reliable, and non-invasive method for predicting the risk of scar dehiscence/rupture [9]. Specific guidelines for TOLAC, after sono graphic assessment of women with previous Caesarean Section, are need of the hour. They showed the sensitivity and specificity of trans abdominal ultrasonography were 91% and 93% respectively for prediction of scar dehiscence [10].

CONCLUSION:

It is concluded that trans abdominal ultrasonography is a useful and reliable method having high value of diagnostic accuracy for prediction of scar dehiscence in females with history of cesarean.

REFERENCES:

1. Aziz N, Yousfani S. Analysis of uterine rupture at university teaching hospital Pakistan. *Pakistan journal of medical sciences*. 2015;31(4):920.
2. Sharma C, Surya M, Soni A, Soni PK, Verma A, Verma S. Sonographic prediction of scar dehiscence in women with previous cesarean

section. *The Journal of Obstetrics and Gynecology of India*. 2015;65(2):97-103.

3. Abosrie M, Farag MAEM. Prediction of cesarean section scar dehiscence before delivery using three-dimensional transabdominal ultrasonography. *Benha Medical Journal*. 2015;32(2):101.
4. Mohammed ABF, Al-Moghazi DA, Hamdy MT, Mohammed EM. Ultrasonographic evaluation of lower uterine segment thickness in pregnant women with previous cesarean section. *Middle East Fertility Society Journal*. 2010;15(3):188-93.
5. Caughey AB, Cahill AG, Guise J-M, Rouse DJ. Safe prevention of the primary cesarean delivery. *American Journal of Obstetrics & Gynecology*. 2014;210(3):179-93.
6. Judith L, Debbie S, Sandra B. *Counseling the nursing mother: a lactation consultant's guide*. Canada Jones & Barlett publishers. 2010.
7. Yeniel A, Petri E. Pregnancy, childbirth, and sexual function: perceptions and facts. *International urogynecology journal*. 2014;25(1):5-14.
8. Geiger W. *The Dīpavaṃsa and Mahāvāṃsa and their historical development in Ceylon*: Government Printer, South Africa; 1908.
9. Lurie S. The changing motives of cesarean section: from the ancient world to the twenty-first century. *Archives of gynecology and obstetrics*. 2005;271(4):281-5.
10. Torpin R, Vafaie I. The birth of Rustam: An early account of cesarean section in Iran. *American journal of obstetrics and gynecology*. 1961;81(1):185-9.
11. Souza JP, Gülmezoglu A, Lumbiganon P, Laopaiboon M, Carroli G, Fawole B, et al. Caesarean section without medical indications is associated with an increased risk of adverse short-term maternal outcomes: the 2004-2008 WHO Global Survey on Maternal and Perinatal Health. *BMC medicine*. 2010;8(1):71.