



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3358938>Available online at: <http://www.iajps.com>

Research Article

**FOETO-MATERNAL OUTCOME IN SCARRED AND
UNSCARRED RUPTURE UTERUS OF SOUTHERN PUNJAB
POPULATION**Dr Ayesha hanif¹, Dr Sidra Naeem², Dr Affaf Shaukat³¹Allama Iqbal Memorial Teaching Hospital, Sialkot, ²Sharif Medical and Dental College Lahore, ³Mayo Hospital Lahore.

Article Received: June 2019

Accepted: July 2019

Published: August 2019

Abstract:**Background:** Uterine rupture is a rare devastating obstetrical complication that endanger the women life.**Objective:** To explore and compare risk factors and complications of foetal and maternal outcome in cases of uterine rupture either unscarred or scarred uterus.**Material and Methods:** Retrospective study was carried out in Department of Obstetrics and Gynaecology at tertiary teaching hospital. Records of all pregnant women who were admitted for delivery during a follow up period of January 2017 to June 2018, data were collected from hospitals records and analysed. 96 patients were included in the study who were identified with rupture uterus in total and only 92 records were analysed the remaining 4 women died before investigation. The remaining subjects were categorized into two groups, group one consisted of unscarred rupture uterus and group two were comprised of scarred uterus with dehiscence.**Results:** Rupture uterus was found in 0.74% pregnant women in our study. Rupture uterus was found in unscarred uterus 52.2% and 47.8% in scarred uterus whereas 70.8% rupture was found at term pregnancy in unscarred uterus and 63.6% in scarred uterus. Complete rupture was found incurring commonly in unscarred uterus 95.8% and scarred uterus (81.8%). Incomplete rupture was found more in scarred uterus (18.1%) in comparison to unscarred uterus (4.1%) ($p < 0.044$). Uterus repair with bilateral tubal ligation was performed more in scarred 25% (11) as compared to unscarred uterus 4 (8.3%) ($p < 0.047$). Total maternal deaths were 9.37% (9) but 4 died without investigation. Perinatal deaths were 91.30% (84).**Conclusion:** Morbidity remained high in unscarred rupture in comparison with rupture of scarred uterus, more hysterectomies performed in unscarred rupture due to involvement of both segment of uterus.**Corresponding author:****Dr. Ayesha hanif,**

Allama Iqbal Memorial Teaching Hospital, Sialkot.

QR code



Please cite this article in press Ayesha hanif et al., *Foeto-Maternal Outcome In Scarred And Unscarred Rupture Uterus Of Southern Punjab Population.*, Indo Am. J. P. Sci, 2019; 06[08].

INTRODUCTION:

Uterine rupture is defined as a full-thickness separation of all the layers of uterine wall including over-lying serosa. The tear could be present on anterior, posterior, or lateral wall of uterus or combination of these. Uterine rupture is typically classified as either complete or incomplete.

Complete uterine rupture, when all layers of the uterine wall are separated and incomplete rupture or scar dehiscence means uterine wall is separated but the visceral peritoneum is intact.

Risk factors for uterine rupture included grand multiparity, mal-presentation, prolonged labour, obstructed labour, uterine anomaly, instrumental deliveries, delivery by unskilled personnel, use of over doses of oxytocin and prostaglandins for induction of labour, Previous uterine surgery (Myomectomy and caesarean section), foetal macrosomia.¹ Nowadays caesarean section is an important risk factor responsible for rupture uterus all over the world.^{2,3} Uterine rupture is a clinical diagnosis and there must be a high index of suspicion by clinician because of variable presentation. Most Women presented with maternal tachycardia, hypovolemic shock, history of constant abdominal pain followed by cessation of uterine contractions, loss of foetal station, uterine tenderness and change in uterine contour and slight vaginal bleeding. The most consistent early indicator of uterine rupture is the on-set of a prolonged, persistent foetal bradycardia but there is no foetal heart rate pattern pathognomonic of rupture.⁴⁻⁶

Aim of this study was to compare the risk factors, complications, foetal and maternal outcome in cases of uterine rupture either unscarred or scarred uterus.

MATERIAL AND METHODS:

This retrospective study was carried out in the department of Obstetrics and Gynaecology, at tertiary care hospital. After ethical clearance re-cords of all pregnant women who were admitted for delivery during a period of January 2017 to June 2018, data were collected from hospitals records and analysed. A total of 96 patient's records had been identified with rupture uterus and we had analysed in detail only 92 women records because 4 women died before any interventions.

As per records these patients who had rupture of uterus were divided into two groups on the basis of previous

surgeries over the uterus: Group A included unscarred 'uterus without any previous history of uterine surgery while Group B consist of scarred uterus with history of one or more previous caesarean section, uterine curettage, previous myomectomy or uterine rupture repair. Both groups were compared for parameters like maternal age, parity, risk factor, site of rupture, operative procedures, foetal outcome, maternal morbidity and mortality.

Every women with rupture uterus received emergency obstetric care first then planned for surgery. Surgical management was performed as repair of uterus with bilateral tubal ligation, repair of uterus without tubal ligation and hysterectomy either total or subtotal.

STATISTICAL ANALYSIS:

The data was collected on predetermined questionnaire from hospital records and analysis was done by using descriptive statistics and frequency with percentages. For group comparison student t-test and chi-square test was used. *P-value* <0.5 considered significant.

RESULTS:

Rupture uterus were identified in 96 cases out of 12,960 admission of pregnant women for delivery during a follow up, thus the rupture of uterus was reported 0.74% in this study.

In present study out of 92 cases of rupture uterus, 52.2% (48) unscarred uterus underwent rupture and 47.8% (44) had rupture of scarred uterus. Here maximum rupture were reported within age group of 26 to 30 year, unscarred uterus rupture found in 54.2% (26) and scarred uterus rupture in 52.2% (23) at this age group. Overall maximum rupture uterus was observed at term gestation age. Uterine rupture at >37 weeks of gestation, present in 70.8% (34) cases of unscarred uterus and 63.6% (28) in scarred uterus. Both groups did not differ significantly in the terms of age, antenatal booking status, history of curettage. There were significant difference in terms of parity and it was also observed that rupture of uterus mainly happened in unbooked and unsupervised deliveries. Rupture of unscarred uterus more common in grand multiparous women (parity > 5) whereas scarred uterine rupture commonly found with par-ity 1 to 2 ($p < 0.001$). (Table 1).

Table 1: Demographic Profile of Women with Rupture Uterus.

Sl.No	Clinical feature	Unscarred uterus (48)	Scarred uterus (44)	p Values
1-	AGE			
	20-25 yr	6(12.5%)	10(22.7%)	0.196
	26-30 yr	26(54.2%)	23(52.2%)	0.856
	31-40 yr	16(33.3%)	11(25%)	0.381
2-	Parity			
	1-2	4(8.3%)	20(45.4%)	<0.001**
	3-4	24(50%)	19(43.1%)	0.513
	>5	20(41.66%)	5(11.4%)	0.001**
3-	H/o of curettage	6(12.5%)	10(22.72%)	0.196
4-	Period of gestation			
	20-28 week	6(12.5%)	0 (0%)	0.027*
	29-37 week	8(16.6%)	16(36.4%)	0.032*
	>37week	34(70.8%)	28(63.6%)	0.462
5-	Booked	0 (0%)	2(4.5%)	0.226
	Unbooked	48(100%)	42(95.4%)	0.226

In present study common causes of uterine rupture in unscarred uterus were obstructed labour 31.5% (29) and multi-parity 16.3% (16) whereas previous caesarean section was the most common risk factor 45.6% (42) for dehiscence of scarred uterus. (Figure 1)

Complete uterine rupture was more common in unscarred uterus 46 (95.8%) as compared to scarred uterus 36 (81.8%) while incomplete uterine rupture was detected more in scarred uterus 8 (18.1%) as compared to unscarred uterus 2 (4.1%) and results were statistically significant ($p<0.044$). Lower segment rupture was found more in scarred uterus 22 (50%) as compared to unscarred uterus 15 (31.2%). After laparotomy records, evaluation both upper and lower segment involvement was observed in 58.3% of unscarred uterus and in 31.8% of scarred uterus ($p<0.011$) Vagina and broad ligament involvement was found more in unscarred uterus rupture as compared to involvement of bladder this might be because of high intrauterine pressure. While cervix and bladder involvement was more commonly found in scarred uterus might be because of passive dehiscence of previous scar. (Table 2)

Total and subtotal abdominal hysterectomies were performed more in unscarred uterus as compared to scarred uterus, this might be because of complete rupture often involve cervix and vagina or extension in broad ligament, making difficult to repair. Uterus repair with ligation was performed more in scarred uterus 11(25%) as compared to unscarred uterus 4 (8.3%) and this difference was statistically significant ($p<0.047$). Requirement of bladder repair was more

with scarred uterus 4 (9%) along with hysterectomy while with unscarred rupture uterus repair in less cases, only 4.1% (2) required bladder repair. Out of 96 women with rupture uterus total 9 (9.37%) maternal deaths were reported in both the groups. Out of 9 deceased, 4 women expired without investigation and 5 (5.43%) women expired after surgery. These 5 (5.4%) deaths were mainly due to haemorrhagic shock and septicaemia. Blood was transfused in each and every women either during intra-operative or postoperative period. Maximum women required three to four-unit packed red cells in both the group, group 1 and 2, in 70.8% and 59.1% respectively. Requirement of Fresh frozen plasma transfusion was 54.2% and 22 (50%) in women with rupture of unscarred and scarred uterus respectively.

Maximum women were discharged in satisfactory condition after this catastrophic event within eight to ten days, 58.3% (28) and 86.4% (38) in unscarred and scarred rupture group respectively. Requirement of Prolonged hospital stay more in unscarred uterus rupture 20 (41.7%) as compared to scarred uterus rupture 6(13.4%) and this difference was statistically significant ($p<0.003$). (Table 3) Perinatal mortality was 91.30% in present study. Total Perinatal deaths 84 out of total 92 cases of rupture uterus. Almost half of the foetus died within the uterus, only 8 live babies were born from scarred uterus group whom incomplete scar dehiscence was detected during emergency caesarean section. These 8 women were admitted in emergency with acute foetal distress and scar tenderness thus suspicion of impending rupture arose. Live birth in scarred uterus rupture was high as

compared to unscarred rupture ($p < 0.002$). Out of 8 live born babies three neonates required neonatal intensive care and out of these two neonates died. (Table 4)

Anaemia was the common complication in women of both groups but it was more in unscarred rupture uterus (79.1%) versus 70.5% in scarred uterine

rupture, followed by paralytic ileum 3 (6.3%) in unscarred versus 4 (9.1%) in scarred rupture uterus. Surgical site infection found in 4.1% and 3 (6.8%) cases of unscarred versus scarred rupture uterus. No statistically significant difference was observed between both two groups in the terms of complication. (Table 5)

Table 2: Distribution According to Site of Rupture Uterus.

Sl.	Site	Unscarred	Scarred	<i>p</i> value
No		uterus	uterus	
		(48)	(44)	
1-		Types of rupture		
	Complete rupture	46(95.8%)	36(81.8%)	
	Incomplete rupture	2(4.1%)	8(18.1%)	0.044*
2-		Site of rupture		
	upper segment rupture	5(10.4%)	8(18.1%)	0.286
	lower segment rupture	15(31.2%)	22(50%)	0.067
	Both US+LS rupture	28(58.3%)	14(31.8%)	0.011*
3-		Involvement of nearest organ		
	No injury	32(66.8 %)	29(65.9 %)	0.002**
	Cervix	2(4.1%)	3(6.8%)	0.667
	Vagina	5(10.4%)	4(9%)	1.000
	Broad ligament	5(10.4%)	3(6.8%)	0.716
	Bladder involvement	4(8.3%)	5(11.4%)	0.732

Table 3: Management and Intra-Operative complications.


Clinical Feature	Unscarred	Scarred	<i>p</i> value
	uterus (48)	uterus (44)	
Uterus Repair	16(33.3%)	13(29.5%)	0.696
Uterus repair with ligation	4(8.3%)	11(25%)	0.047*
Hysterectomy			
Total abdominal hysterectomy	23(47.9%)	17(38.6%)	0.370
Subtotal hysterectomy			
	5(10.4%)	3(6.8%)	0.716
Other structure involvement			
Bladder repair	2(4.1%)	4(9%)	0.421
Iontropic support	12 (25%)	10(22.7%)	0.799
Ventillatory support	18(37.5%)	12(27.3%)	0.296
Mortality	3(6.8%)	2(4.7%)	1.000
Duration in ventilatory unit.	14(29.2%)	10(22.7%)	0.482
(days)			
Total hospital stay	28(58.3%)	38(86.4%)	
8-10	20(41.7%)	6(13.4%)	0.003**
>14 days			
Blood transfusion	48	44	
One-unit PRBC	3(6.3%)	4(9.1%)	0.706
Two-unit PRBC	11(22.9%)	14(31.8%)	0.338
Three to four-unit PRBC	34(70.8%)	26(59.1%)	0.238
Fresh frozen plasma	26(54.2%)	22(50%)	0.689

Table 4: Foetal Outcome.

Sl.No	Clinical feature	Unscarred uterus (48)	Scarred uterus (44)	p value
1-	Live	0 (0%)	8(18.1%)	0.002**
2-	Still birth	48(100%)	36(81.8%)	0.310
3-	Neonatal intensive unit	0 (0%)	3(6.8%)	0.106
4-	Neonatal death	0 (0%)	2(4.5%)	0.226
5-	Birth weight of baby			
	<2.5 kg	6(12.5%)	8(18.1%)	0.449*
	2.5 to 3.5kg	25(52%)	18(40.9%)	0.283
	>3.5	17(35.4%)	22(50%)	0.157

Table 5: Post-Operative Morbidity.

	Maternal morbidity/ Morbidity	Unscarred uterus (48)	Scarred uterus (44)	p value
	Anaemia	38(79.1%)	31(70.5%)	0.335
	Paralytic ileus	3(6.3%)	4(9.1%)	0.706
	Surgical site infection	2(4.1%)	3(6.8%)	0.667
	Fever	5(10.41%)	6(13.6%)	0.635

Risk factors T-92**Figure 1:** Risk Factors Responsible for uterus rupture.

DISCUSSION:

Uterine rupture is an uncommon devastating obstetrical emergency associated with significant maternal and perinatal morbidity and mortality.⁷ Usually it is more frequently associated with scarred uterus as compared to unscarred uterus. Maternal mortality is the worst complication of uterine rupture, it occurs in approximately 1 in 500 uterine ruptures.⁸ According to WHO the incidence of rupture uterus in general population is 5.3/10,000 birth.⁹ In developing countries incidence of uterine rupture is far higher (1 in 106)¹⁰ as compared to the developed countries where it is approximately 74 in 10,000.¹¹ Rupture of an unscarred uterus is a rare event involving 1:17,000–20,000 deliveries.¹² In present study incidence of uterine rupture was 0.74%, while study done by other authors reported incidence of rupture, 0.038% and 0.057%.^{13,14} This high incidence might be due to increased caesarean section. In our study all women were unbooked (100%) and had not received any antenatal care during pregnancy similarly other studies reported 80% and 78.68% women were unbooked in their study.^{15,7}

Here rupture commonly found in 26 to 30 years of age group and next common age group was 31 to 40 year similarly another study reported increasing maternal age has detrimental effect on uterine rupture.¹⁶ In present study among unscarred group 50% rupture were found within parity of 3 to 4 followed by 41.66% with parity >5, whereas maximum cases 43.1% rupture in scarred uterus group found within the parity of 3 to 4, similarly another studies also reported multiparity is a very important predisposing factor which was found in 97.9% of cases.¹⁷ One study reported 32% unscarred uterine rupture had a parity of 4 and more.¹⁸

In present study common causes of unscarred uterine rupture were obstructed labour in 31.5% and multiparity in 16.3%, cephalopelvic disproportion 3.20% whereas previous caesarean section was the most common risk factor in 45.6% for scarred uterine rupture. In contrast study done by other author common causes of rupture uterus were Pitocin induced labour in 51.6%, great multiparity in 42.2%, Previous uterine scar in 18.8% and obstructed labour in 12.5%.¹⁹ Another studies reported common causes of rupture uterus were great multiparity in 41.5%, Pitocin induced labour in 58.5%, malpresentation and malposition in 12.1% and CPD 9.8%.^{20,21}

One study reported 19.2 % association of previous caesarean section with rupture uterus.²² In present

study 52.2% in unscarred group and 47.8% in scarred group had uterine rupture similarly one study reported higher incidence 85.3% unscarred rupture while rupture in scarred uterus 14.7% were found with previous scar.²³ Hysterectomy performed in present study, in 47.9% cases of unscarred rupture while in 38.6% scarred rupture uterus and study done by another authors reported hysterectomies in 34.86% unscarred rupture and in 35% scarred rupture uterus.^{24,25} In present study lower uterine segment rupture commonly observed in both scarred and unscarred uterus in contrast one author reported lower segment rupture in 80% of cases.²⁶ Total maternal mortality was 9.3% in our study while another study reported 11.43%.²⁷ In our study perinatal mortality was 91.30% while other study reported 78.66%.¹⁵ Subtotal hysterectomy done in 10.4% and 6.8% cases of unscarred and scarred rupture respectively while another study reported subtotal hysterectomy in 32.1% cases.¹³

CONCLUSION:

In our study morbidity is high with unscarred rupture as compared to rupture of scarred uterus, more hysterectomies performed in unscarred rupture due to involvement of both segment of uterus and extension of tear to the vaginal fornices. Thus, suggested primary focus is to increase institutional deliveries, proper antenatal care and to avoid unnecessary caesarean section. Early surgical intervention is the main key to achieve good results. In present study perinatal mortality 91.30% and maternal deaths were 9.37%. Thus, reduction of maternal morbidity, mortality and perinatal mortality possible by preventing rupture of uterus.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

REFERENCES:

1. Smith JG, Mertz HL, Merrill DC. Identifying risk factors for uterine rupture. *Clin Perinatol.* 2008;35(1):85-99.
2. Sahin H, Kulusari A, Yildizhan R, Kurdoglu M, Adali E, Kamaci M. Uterine rupture: A twelve-year clinical analysis. *J Maternal Fetal Neonatal Med.* 2008;21(7):503-6.
3. Vanden AT, Beatrice M, James I, Roosmalen JV. Using audits to reduce the incidence of uterine rupture in a Malawian district hospital. *Int J Obstetric Gynecol.* 2009;10(3):289-94.
4. Menihan CA. Uterine rupture in women attempting a vaginal birth following prior cesarean birth. *J Perinatol.* 1998;18(6 Pt 1):440-3.

5. Hawe JA, Olah KS. Posterior uterine rupture in a patient with a lower segment caesarean section scar complicating prostaglandin induction of labor. *BJOG Int J Obstet Gynaecol.* 1998;104:857-8.
6. Ozdemir I, Yucel N, Yucel O. Rupture of the pregnant uterus: a 9-year review. *Arch Gynecol Obstet.* 2005;272(3):229-31.
7. Aziz N, Yousfani S. Analysis of uterine rupture at university teaching hospital Pakistan. *Pak J Med Sci.* 2015;31(4):920-4.
8. Wen SW, Huang L, Liston R, *et al.* Severe maternal morbidity in Canada, 1991–2001. *Canadian Medical Association Journal.* 2005;173(7):759-63.
9. Hofmeyr GJ, Say L, Gulmezoglu AM. WHO Systemic review of maternal mortality and morbidity: The prevalence of Uterine Rupture. *BJOG.* 2005;112(9):1221-8.
10. Ezegwui HU, Nwogu-Ikojo EE. Trends in uterine rupture in Enugu, Nigeria. *J Obstet Gynaecol.* 2005;25(3):260-2.
11. Landon MB, Hauth JC, Leveno KJ, *et al.* Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *N Engl J Med.* 2004;351(25):2581-9.
12. Ofir , Sheiner E, Levy A, Katz M, Mazor M. Uterine rupture: differences between a scarred and an unscarred uterus. *American Journal of Obstetrics and Gynecology.* 2004;191(2):425-9.
13. Ahmadi S, Nouria M, Bibi M. Uterine rupture of the unscarred uterus. About 28 cases. *Gynecol Obstet Fertil.* 2003;31(9):713-7.
14. Aggarwal P, Terhase N. Unscarred uterine rupture: a retrospective study. *Int J Reprod Contracept Obstet Gynecol.* 2015;4(6):1997-9.
15. Rashmi, Radhakrishnan G, Vaid NB, Agarwal N. Rupture uterus- changing Indian scenario. *J Indian Med Assoc.* 2001;99(11):634-7.
16. Shipp TO, Zelop C, Repke JT, *et al.* The association of maternal age and symptomatic uterine rupture during a trial of labor after prior cesarean delivery. *Ob-stet Gynecol.* 2002;99(4):585-8.
17. Duhan N, Sangwan N, Dahiya K, Sirohiwal D, Paul A. Rupture of the gravid uterus: A review of 92 cases. *J Gynecologic Surg.* 2011;27(1):17-9.
18. Schrinky DC, Benson RC. Rupture of the pregnant uterus: a review. *Obstet Gynecol Surv.* 1978;33(4):217-32.
19. Qazi Q, Akhtar Z, Khan K, Khan AH. Woman Health, Uterus Rupture, Its Complications and Management in Teaching Hospital Bannu, Pakistan. *Maedica.* 2012;7(1):49-53.
20. Kidantou HL, Mwampagatwa I, Roosmalen JV. Uterine rupture. Retrospective analysis of causes, complications and management outcomes at Muhimbili Na-tional Hospital in Dar es Salaam, Tanzania. *Tanzan J Health Res.* 2012;14(3):220-5.
21. Fofie C, Baffoe P. A two-year review of uterine rupture in a regional hospital. *Ghana Med J.* 2010;44(3):98-102.
22. Mishra SK, Maris N, Uprty OK. Uterus repture: Preventable obstetric tragdies?. *Aust NZJ Obstet Gynecol.* 2006;46(6):541-5.
23. Khan S, Parveen Z, Begum S, Alam I. Uterus rupture a review of 34 cases at Ayub Teaching Hospital. *Abbothabad.* 2003;15(4):50-2.
24. Khan B, Sultana R, Bashir R, Deeba F. A Ten yearn review of emergency peri-partum hysterectomy in a tertiary care hospital. *J Ayub Med Coll Abbottabad.* 2012;24(1):14-7.
25. Shah N, Khan N. Emergency Obstetric Hysterectomy. A review of 69 cases. *R Med J* 2009;34(1):75-8.
26. Rizwan N, Abbasi RM. Uterine rupture, frequency of cases and fetomaternal outcome. *JPMA.* 2011;61(4):322.
27. Dwivedi S, Kumar A. Uterine rupture a retrospective analysis of referral cases at a tertiary care centre in Kanpur City. *Internat J Reproductive Contracept, Obstetrics and Gynaecol.* 2017;4(4):1148-52.