



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3362490>Available online at: <http://www.iajps.com>

Research Article

**CENTRAL NERVOUS SYSTEM TUMORS: A HOSPITAL
BASED ANALYSIS**¹Madiha Sarwar, ²Fatima Ashfaq, ³Alina Fatima¹Rawalpindi Medical University, ²WMO, Gen. Surgery, Aziz Bhatti Shaheed Teaching Hospital, Gujrat, ³KEMU, Mayo.**Article Received:** June 2019**Accepted:** July 2019**Published:** August 2019**Abstract:**

Objective: To evaluate the range of CNS tumours and give benchmark information to future studies surveying information in the continuum.

Patients and Methods: This descriptive research was carried out at Jinnah Hospital, Lahore from January 2018 to February 2019. One hundred cases satisfying the consideration criteria were incorporated between the age of 1-85 years selected from both genders.

Results: Larger parts of the cases were found in the year 2008 with the most usually experienced injury being the frosty tumours pursued by the meningotheial neoplasms. Our discoveries were like past comparative examinations in our setup with little change in patterns.

Conclusion: Glial tumours seem, by all accounts, to be increasingly normal in our setup while the famously trusted Meningotheial tumours however regular came straightaway. Our investigation can frame the benchmark information whereupon future examinations can be led.

Keywords: Tumors, Meningotheial, Frosty, Neoplasms, Glial, Mortality and Morbidity.

Corresponding author:**Madiha Sarwar,**

Rawalpindi Medical University.

QR code



Please cite this article in press Madiha Sarwar et al., *Central Nervous System Tumors: A Hospital Based Analysis.*, Indo Am. J. P. Sci, 2019; 06(08).

INTRODUCTION:

Focal sensory system (CNS) tumours have dependably been a reason for worry among histopathologists because of the wide assortment in their appearances. Diagnosing a CNS tumour represents a noteworthy test, numerous multiple times requiring exceptional demonstrative systems, for example, and immunohistochemical staining [1]. Early determination and the executives of mind tumours are imperative to diminish bleakness and mortality. Frequency and Mortality Rate Trends in the United States in 2007 appeared more than 50,000 new instances of CNS tumours with 7.4% cases from adolescence group [2]. While in the United Kingdom the evaluated lifetime danger of creating CNS malignant growth was observed to be 1 of every 133 for men and 1 of every 185 for women [3]. Most focuses in Pakistan have cross-sectional information in regards to CNS tumours because of the way that this feature of pathology is generally uncommon when contrasted with tumours of different locales with CNS tumours happening at under 2% of all essential tumours [4, 5]. Anyway, they seem, by all accounts, to be regular in children [6]. Glial neoplasms have been viewed as basic CNS tumors in numerous studies [7, 12]. Spinal rope tumors structure about 15% of focal sensory system tumors. Most regular sorts incorporate neurofibromas and meningiomas [13]. In spite of the fact that information about the range of CNS tumors in our populace is accessible from focuses over our nation [4, 5, 8 10 – 12], everyone has its own restrictions as far as sick populace, along these lines a solitary arrangement of results can't be connected consistently in all cases. In our middle past investigations concentrated on malignancies all in all and not especially upon CNS tumors hence an appraisal of this region is without a doubt justified. Since close checking of the movements in patterns are indispensable to evaluate the malady load as well as

to control essentially required enhancements in the clinical consideration rehearses. The motivation behind this examination was to evaluate the range of CNS tumors and give benchmark information to future investigations surveying information in the continuum.

PATIENTS AND METHODS:

This descriptive research was carried out at Jinnah Hospital, Lahore from January 2018 to February 2019. Sequential non-likelihood examining method was utilized. Tests marked as CNS tumours were incorporated; both kindhearted and harmful, from patients having a place with the two sexes between the ages of 1-85 years. All inappropriately transported and/or autolyzed tests were prohibited. An aggregate of one hundred instances of CNS tumours satisfied the incorporation criteria. The histopathological reports of every single included case were looked into in detail while Hematoxylin and Eosin (H and E) slides of irregular cases were likewise explored from the slides record. Tiny judgments, patient's age, sexual orientation were noted down in Microsoft Excel database document. The information was broken down on the Statistical Package for Social Sciences (SPSS). Engaging insights were utilized to depict the information.

RESULTS:

Of the one hundred included cases, 27% were from the year 2008 pursued constantly 2004 which contributed 19% of cases. Patient's ages went from 2 years to 85 years with a mean period of 39.7 years. Grown-up populace represented 91% cases while 9% of patients had a place with the pediatric age gathering. Most patients had a place with the fifth decade pursued constantly most elevated gathering in the fourth decade of life though patients in their third decade of life came straightaway. (Fig.1).




Figure – I: Distribution of CNS tumours according to various age groups in our study

There was a male power in our examination with male patient's framing 73 % of the included cases. Our outcomes likewise demonstrated a prevalence of Glial tumours at 58 % pursued by 18% Meningothelial neoplasm out of which five were atypical meningiomas. The point by point range of

histological injuries experienced in our examination cases is given in Table 1.

Photomicrograph of an Astrocytoma WHO Grade-II from our study group is shown in (Fig.2).

Tumor Type(s)	Histological Diagnosis	Number of Cases	Percentage
Glial Tumors 58 (58%)	Astrocytoma WHO Grade-IV (Obablastoma Multiform)	23	23%
	Astrocytoma WHO Grade-III	10	10%
	Astrocytoma WHO Grade-II	11	11%
	Oligodendroglioma WHO Grade-II	8	8%
	Astrocytoma WHO Grade-I (Pilocytic Astrocytoma)	4	4%
	Mixed Oligo-Astro Neoplasm	2	2%
Meningothelial Tumors 18 (18%)	Meningioma WHO Grade-I	13	13%
	Meningioma WHO Grade-II (A typical Meningioma)	5	5%
Metastatic 7 (7%)	Metastatic Carcinoma	7	7%
Others 17 (17%)	Medalloblastoma	4	4%
	Hemangioblastoma	4	4%
	Ependymoma WHO Grade-II	6	6%
	Schwannoma	1	1%
	Non-Hodgkin's Lymphoma-Diffuse Large Cell	1	1%
	Completely Necrotic Tumor	1	1%




Figure – II: Astrocytoma WHO Grade-II H&E

DISCUSSION:

Our outcomes are in concurrence with different investigations over the country [4, 5, 8, 10 – 12], which demonstrate the prevalence of Glial tumours particularly in patients exhibiting in their fourth to sixth decade of life. Astrocytomas regularly show high rates of neighbourhood intrusion that lead to a nearby repeat of the ailment which is a noteworthy reason for horribleness and mortality. It has been demonstrated that multiplication and relocation are fundamentally unrelated practices. An ensemble of these empowers these tumours to attack and repeat. As of now referenced the sickness weight of these tumours is additionally expanded by the perplexing administration frequently required to help these patients which incorporates careful resection, radiotherapy and chemotherapy [14 – 16]. Visualization is by all accounts useful for second rate totally resected tumours [17]. It is contended that the frequency and death rates for CNS tumours have changed almost no throughout the decade, with men having higher occurrence and death rates than ladies. Four and a half thousand new instances of CNS malignant growths were analyzed in Europe in the year 2006 alone [3, 18]. In studies led at our middle just as at Armed Forces Institute of Pathology (AFIP) Rawalpindi which concentrated on ten commonest threatening tumours demonstrated low recurrence of the CNS tumours when contrasted with other sites [4, 5].

Gliomas and meningiomas have been regularly viewed as the most well-known CNS tumours pursued by craniopharyngiomas, pineal locale tumours, acoustic neuromas and choroid plexus papilloma though in youth age bunch additionally, the example is similar [6, 7, 8, 10]. Different focuses crosswise over Pakistan likewise share the comparative results [4, 5, 8, 10 – 12]. CNS lymphomas involve 4.6% of all CNS neoplasms and contain about 2.2% of the non-Hodgkin's lymphomas [19]. Different rarities with respect to CNS incorporate intense myeloid leukaemia (AML) penetrate. Meningothelial tumours shaped the second biggest gathering of tumours in our examination involving 18% of the cases which converts into the colossal issue of tumour repeat looked by the patients just as the treating doctors of meningothelial tumours. The assessed repeat rate is in the scope of 10%–15% at 5 years and 25%–37% at an interim of 10-years following a corrective surgery [20]. The significance of H and E morphology can't be over underlined as the tumour morphology has been to a great extent acknowledged as a solid and dependable prognostic factor in anticipating odds of infection repeat. This contention is upheld by the way that a

solitary couple of World Health Organization (WHO) Grade I meningiomas repeat while higher evaluation meningothelial tumours, for example, atypical (WHO Grade-II) and anaplastic meningiomas (WHO Grade-III) do undoubtedly repeat in the principal years following total careful evacuation of the essential lesion [21]. This apparently honest sore, when contrasted with the glial tumours, causes critical bleakness if not mortality attributable to the weight side effects, particularly in spinal areas. When contrasted with different investigations metastatic tumours were more for example 7% in number in our investigation, maybe because of the reason of referrals of patients from fringe medical clinics and country setups where odds of early recognition of malignancies are troubling. Medulloblastoma is believed to be the commonest threatening CNS tumour of youth, of which 4 cases were accounted for in this investigation, comparatively four instances of Hemangioblastoma was additionally detailed. Schwannoma and Non-Hodgkin's Lymphoma – Diffuse Large Cell Type were moderately uncommon elements at 1 % each. Tumour morphology couldn't be found out in one situation where the whole tumour tissue was totally necrotic.

CONCLUSION:

Glial tumours seem, by all accounts, to be increasingly basic in our setup while the prevalently trusted meningothelial tumours however normal shaped the second biggest gathering of neoplasms. Moreover, assessment of CNS tumours' morphology and recurrence can help in checking of infection inclines and evolving designs. Our investigation frames the benchmark information whereupon future examinations can be directed to shape information in continuum for successfully regulating any enhancements which become visible amid the symptomatic procedures.

REFERENCES:

1. Ahmed Z, Azad NS, Bhurgari Y, Ahmed R, Kayani N, Pervez S et al. Essentialness of Immunohistochemistry in the precise portrayal of threatening tumours. *J Ayyub Med Coll* 2006; 18(2): 38-43.
2. Central Brain Tumor Registry of the United States. Hotspot for rate and mortality information: Surveillance, Epidemiology, and End Results (SEER) Program and the National Center for Health Statistics. US: National community for wellbeing insights. Accessible from: <http://www.cbtrus.org/factsheet/factsheet.html>
3. Office for National Statistics, Cancer Statistics enrollments: Registrations of malignant growth

- analyzed in 2006, England. Arrangement MB1 no.37. 2009, National Statistics: London.
4. Zahir N, Mubarak An, Abdullah P, Mehmood A, Rehman Z. Example of Malignant Tumors in Karachi is it unique? *J Coll Physicians Surg Pak* 2000; 10(9): 338-41.
 5. Jamal S, Mughal S, Mamoon N, Mushtaq S, Luqman M, Anwar M. The example of threatening tumours: Tumor library information investigation, AFIP, Rawalpindi, Pakistan (1992-2001). *J Pak Med Assoc* 2006; 56(8): 359-62.
 6. Tanwani AK, Ahmad I, Qureshi AM. Histological example of youth tumours at Children Hospital, P.I.M.S., Islamabad. *J Pak Inst Med Sci* 2002; 13(2): 694-7.
 7. Shah S, Soomro I, Hussainy A, Hassan S. Clinico-morphological example of intracranial tumours in youngsters. *J Pak Med Assoc* 1999; 49(3): 63-5.
 8. Bhatti SN, Ali J, Ahmed An, Ahmed Z, Aziz T. Sorts of mind tumours in Hazara Division and Northern territories of Pakistan. *J Ayyub Med Coll* 2004; 16(1): 44-6.
 9. Salahuddin T, Hussain MA. Mind stem tumours: Surgical administration. *Proficient Med J* 2002; 9(3): 222-5.
 10. Ahmed Z, Azad NS, Muzaffer S, Nasir I, Hasan S. CNS tumours at AKU: an update in addition to a concise dialogue on Intraventricular Tumors with exceptional accentuation on Central Neurocytoma *J Ayyub Med Coll* 2004; 16(4): 12-5.11.
 11. Jamal S, Mamoon N, Mushtaq S, Luqman M. Example of focal sensory system (CNS) tumours: an investigation of 430 cases. *Pak J Pathol* 2005; 16(4): 106-9.
 12. Yaqoob M, Masood MK, Waseem R, Zaheer an, Izhar TS, Qureshi AW et al. Clinical introduction, morphological sorts and anatomical appropriation of essential intracranial tumours in kids. *Pak Paed J* 2005; 29(2): 69-75.
 13. Salahuddin T, Shams S, Hussain N. Spinal string tumours; the job of microsurgery. *Proficient Med J* 2001; 8(4): 530-3.
 14. RL Souhami, I Tannock, P Hohenberger, J-C Horiot, eds. *Course reading of Oncology*. Second ed. Oxford University Press; 2002.
 15. Souhami, Tobias, eds. *malignant growth and Its Management*. Fourth ed. Oxford Blackwell Scientific Publications; 2003.
 16. Levin, eds. *Disease in the Nervous System*. second ed. Oxford University Press; 2002.
 17. Buckner JC, Brown PD, O'Neill BP, Meyer FB, Wetmore CJ, Uhm JH. Focal Nervous System Tumors. *Mayo Clinic Proc* 2007; (10): 1271-86
 17. Northern Ireland Cancer Registry, *Cancer Incidence and Mortality*. 2009
 18. Muzaffar S, Siddiqui M, Siddiqui I, Soomro I, Pervez S, Kayani N, Hasan S. Focal sensory system lymphomas: a histologic and immunophenotypic investigation. *J Pak Med Assoc* 2000; 50(5): 141-4.
 19. Ayerbe J, Lobato RD, de la Cruz J, Alday R, Rivas JJ, Gómez PA et al. Hazard factors anticipating repeat in patients worked on for intracranial meningioma. A multivariate examination. *Acta Neurochir (Wien)* 1999; 141: 921-32.
 20. Hilbig A, Barbosa-Coutinho LM. Meningiomas: Histopathological viewpoints and repeat. *Arq Neuropsiquiatr*. 1997; 55:431-7.