



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3370299>Available online at: <http://www.iajps.com>

Research Article

**PERIOPERATIVE MANAGEMENT OF AN ADOLESCENT
WITH TRANSPOSITION OF THE GREAT ARTERIES, STATUS
POST SENNING PROCEDURE, FOR POSTERIOR SPINAL
FUSION**¹Dr Raja Hassan Talal, ²Dr Ateeqa Azhar, ³Dr Fozia Saeed¹Medical Officer Bashir Hospital, Sialkot. ²Woman Medical Officer, Anesthesia Department, Allama Iqbal Memorial Teaching Hospital, Sialkot. ³Woman Medical Officer, Federal Government Polyclinic, Islamabad.**Article Received:** June 2019**Accepted:** July 2019**Published:** August 2019**Abstract:**

With advances in careful procedures and preoperative consideration, the survival rate of patients with intrinsic coronary disease keeps on expanding. All things considered, patients with vindicated or amended inborn coronary illness may exhibit for major surgeries. Albeit elective strategies are currently accessible for the anatomic fix of patients with transposition of the extraordinary corridors (TGA), an atrial dimension baffle (Mustard or Senning Method) was recently performed. As these patients age, long haul unfriendly impacts may happen including myocardial brokenness, cadence unsettling influences requiring pacemaker situation, pneumonic hypertension, and baffle impediment. We present a case report of a multiyear old young lady with TGA, who experienced a Senning system amid the earliest stages and now introduced for back spinal combination in the treatment of kyphoscoliosis. The preoperative consideration of such patients is talked about.

Keywords: Congenital Heart Disease; Transposition of the Great Arteries; Senning Procedure; Mustard Procedure; Myocardial Dysfunction; Pulmonary Hypertension; Baffle Obstruction.

Corresponding author:

Dr. Raja Hassan Talal,
Medical Officer Bashir Hospital, Sialkot.

QR code



Please cite this article in press Raja Hassan Talal et al., *Perioperative Management of an Adolescent with Transposition of the Great Arteries, Status Post Senning Procedure, For Posterior Spinal Fusion.*, Indo Am. J. P. Sci, 2019; 06(08).

INTRODUCTION:

Inborn coronary illness has a detailed occurrence of roughly 0.8 per 100 births [1,2]. As such, patients with mitigated or rectified intrinsic coronary illness (CHD) may show for major surgeries. In spite of the fact that the objective of current careful methodologies is to accomplish an anatomic fix, preceding the present advances in systems, surgeries oftentimes come about just in whitewashing. In such patients, suitable tender loving care amid the preoperative workup, just as the intraoperative and postoperative consideration can encourage a fruitful preoperative course. We present a patient with transposition of the incredible veins (TGA) who had experienced a Senning technique amid the earliest stages who currently shows for back spinal combination (PSF) in the treatment of kyphoscoliosis. The perioperative consideration of such patients is talked about.

CASE REPORT:

A14-year-old young lady with TGA, who experienced a Senning methodology amid early stage had now exhibited for back spinal combination in the treatment of kyphoscoliosis. Audit of her restorative record and introduction of the material in this arrangement were endorsed by Jinnah Hospital, Lahore (October 2018). Her previous history was significant for D-transposition of the extraordinary vessels analyzed not long after birth. Amid earliest stages, she had a Senning methodology. At 3 years old, a VVI pacemaker was set because of discontinuous third-degree heart square. In spite of the fact that she had no other real therapeutic issues, her mom expressed that she was taking successive evening rests because of weariness and did not play aggressive b-ball this year as she couldn't stay aware of her companions. Current prescriptions included sotalol 40 mg two times per day and enalapril 5 mg two times every day. The preoperative assessment uncovered an all around grew, very much fed pre-adult in no intense trouble. Imperative signs uncovered a sporadic mood with a pulse of 84 beats/minute, circulatory strain (BP) of 104/68 mmHg, respiratory rate of 16 breaths/minute, and a room air oxygen immersion of 97%. Physical examination was unremarkable. Echocardiography uncovered no proof of baffle hole or impediment, the privilege ventricular pit was somewhat enlarged with insignificant pressure of the left ventricular outflow tract. There was moderate tricuspid disgorging and follow mitral spewing forth. There was moderate worldwide defection of right ventricular capacity. Preoperative electrocardiograph uncovered the nearness of pacemaker spikes. Late Holter observing uncovered that the patient was pacemaker

subordinate under 5% of the time. Preoperative research centre assessment uncovered a hemoglobin estimation of 14.2 gm/dL, a hematocrit of 40.8%, and a platelet tally of 312,000/mm³. The prothrombin time was 13.8 seconds, the INR was 1.1, and the fractional thromboplastin time was 33 seconds. A dialogue with the pediatric cardiology administration was held in regards to suitable pacemaker care amid the surgery. They were worried that the utilization a magnet and change of the pacemaker into a VOO mode would prompt the danger of arrhythmia acceptance identified with an R on T wonder. Hence, their suggestion was to leave the pacemaker in the VVI mode amid the case. Since the patient was pacemaker subordinate under 5% of the time, they were not concerned with respect to searing impedance. The patient was kept nil per os for 6 hours. Her morning dosages of sotalol and enalapril were held. She transported to the working room where routine screens per the American Society of Anesthesiologists rules were put. Given the patient's nervousness, inhalational acceptance was done with steady increments in the enlivened grouping of sevoflurane in 70% nitrous oxide in oxygen. At the point when a suitable profundity of anesthesia was accomplished, a 16G intravenous cannula was put in the left lower arm. Tracheal intubation was encouraged with sufentanil 50 µg and a solitary portion of rocuronium 50 mg. Anti-toxin prophylaxis was finished with cefazolin (2 gm) IV. Following tracheal intubation, a second fringe intravenous cannula, an outspread blood vessel cannula, and a twofold lumen focal venous catheter were put. The pattern CVP was 10-12 mmHg. Upkeep anesthesia comprised of desflurane (terminated focus 4%) and a sufentanil imbue ment changed in accordance with keep up the mean blood vessel weight at 55-65 mmHg. No extra portions of neuromuscular blocking operators were controlled amid the case. Cathodes were put for neurophysiologic observing (engine and somatosensory evoked possibilities). Test incitement for engine evoked possibilities (MEP's) uncovered that satisfactory possibilities could be gotten with 450 mV without obstruction with the pacemaker. The patient was turned inclined onto a Jackson table for the methodology. The pacemaker module was noted with the patient turned inclined and fitting access guaranteed should it be important to put a magnet on it amid the case. A magnet was kept prepared on the anesthesia truck amid the case. The sufentanil mixture rate was expanded to a limit of 0.7 µg/kg/hr to keep up the MAP at 55-65 mmHg. Amid utilization of the searing for the dismemberment of the paraspinal muscles, there was a periodic obstruction with the pacemaker and loss of capacity; be that as it may, the patient's HR was ≥ 60

beats/minute all through the case. The length of the surgery was 4.5 hours. Blended venous oxygen immersion estimated from the CVP cannula differed from 65-78% amid the case. Intraoperative blood misfortune was roughly 500 ml. The liquid organization incorporated a sum of 2200 ml of Lactated Ringers answer for keep up the CVP at 8-12 mmHg. Pee yield amid the case was 950 ml. A solitary portion of sodium bicarbonate (50 m Eq) was controlled for a base deficit of -6. The final hemoglobin was 10.9 gm/dl with a hematocrit of 32%. An hour prior to the culmination of the surgery, the sufentanil implantation was ceased and an aggregate of 1 mg of hydromorphone was controlled in augmentations of 0.2 mg depending on the patient's respiratory rate. Endless supply of the surgery, the desflurane was stopped, and the patient was turned recumbent. She immediately opened her eyes and moved all furthest points in light of agonizing improvements. Her trachea was extubated and she was exchanged to the post-anesthesia care unit. Postoperative absence of pain was furnished by the patient-controlled absence of pain with hydromorphone. Supplemental oxygen at 2 lit/min was regulated utilizing nasal cannula for the first 24 postoperative hours. Her standard prescriptions of sotalol and enalapril were restarted on the first postoperative day. She was changed to oral analgesics (hydrocodone with acetaminophen) on the third postoperative day and released home on the fourth postoperative day.

DISCUSSION:

TGA, classified as a cyanotic innate heart contortion, exists when the aorta begins from the correct ventricle and the pneumonic trunk emerges from the left ventricle. TGA is partitioned into dextrolooped (D-TGA) and levo-circled (L-TGA) in light of whether the atria and ventricles are concordant or harsh, respectively.³ Although L-TGA, otherwise called inherently revised transposition, is a perioperative administration in transposition of incredible supply routes uncommon imperfection representing roughly 0.5% of all inborn cardiovascular abnormalities, D-TGA is the second most normal inherent heart deformity noted during childbirth influencing 1:3500-500 live births with a male: female proportion of 3.2:1.3 With D-TGA, a morphological right chamber is associated with a morphologically right ventricle which offers ascend to the aorta. A morphological left chamber is associated with a morphological left ventricle which offers ascend to the pneumonic conduit. This mix of concordant atrioventricular and dissonant ventriculoarterial associations makes two parallel dissemination frameworks in which oxygenated

blood recycles inside the pneumonic circuit through the left ventricle and aspiratory trunk while deoxygenated fundamental blood recycles to the body by means of the correct ventricle and aorta. When contrasted with a physiologically ordinary course framework that exists in the arrangement, the parallel circuits that are available with D-TGA result in foundational cyanosis. Postnatal survival of a newborn child with D-TGA is subject to the nearness of intracardiac blending of blood through an atrial septal imperfection (ASD), a patent foramen ovale (PFO), or a ventricular septal deformity (VSD). Infants with flawless atrial and ventricular septa are cyanotic during childbirth and quickly decompensate because of lacking blending of the blood from the two parallel courses. Palliative treatment with quick inflatable or careful septostomy is important to expand the blending of blood at the arterial level. Prostaglandin E imbuelements might be utilized as an adjunctive treatment to keep up ductal patency, accordingly expanding the left-to-right shunting of blood over the ductus arteriosus. The expanded blood from the aorta over the ductus arteriosus to the aspiratory supply route builds the flow of blood over the lungs into the left chamber. The expanded left atrial return builds left atrial weight, in this manner encouraging the blending of blood at the atrial dimension. Neonates with D-TGA and a related ASD or VSD might be just gently cyanotic during childbirth. Manifestations might be ignored until 2 months and a half of age, at which time changes in the pneumonic vascular obstruction happen, bringing about the improvement of congestive heart failure. Initially, fringe vascular opposition (PVR) is more noteworthy than, or proportional to the foundational vascular opposition, permitting halfway blending of blood between the parallel fundamental and aspiratory circuits by means of the ASD or VSD. As PVR falls, the equalization is irritated consequently permitting expanded shunting from the foundational to the pneumonic circuit bringing about aspiratory clog and congestive heart disappointment. Around then, the newborn children frequently present with tachypnea, tachycardia, and mellow fundamental cyanosis.³ The Senning task for TGA was first announced by Senning in 1959, with later work by Quaegebeur in the late 1970s. It turned into the activity of decision for a careful fix of TGA amid this timespan. The Senning strategy utilizes a 'baffle' between the atria bringing about the redirection of deoxygenated vena cava blood through the mitral valve to the morphological LV and pneumonic courses and oxygenated aspiratory venous blood through the tricuspid valve into the morphological RV and aorta. The 'baffle' made in the Senning methodology is gotten from autologous atrial tissue

that is cut and collapsed in a perplexing way to take into account the redirection of blood flow. Because of the specialized difficulty related to the arrangement of the Senning 'baffle', this technique was not generally grasped. In 1964, Mustard simplified the technique by extracting the atrial septum and putting a 'solitary, engineered pantaloon-molded fix' between the atria to make the 'baffle', again diverting the blood flow in a similar way as the Senning procedure. However, neither the Senning nor the Mustard methodology was without long haul inconveniences. By and large, the survival rate for patients following clinic release was 84-95% at 10 years and 76-89% at 15-20 years, with an actuarial decrease in survival of roughly 0.5% per year. Sinus hub brokenness is a standout amongst the most well-known late difficulties after atrial-level fix with just 40% of patients staying in sinus mood 15-20 years post surgery. The associated etiology with this intricacy is either harm to the sinus hub supply route amid medical procedure or the dynamic improvement of fibrosis at the atrial careful scar lines. A junctional escape beat is basic in patients with sinus hub brokenness along these lines restricting the ordinary increment in a pulse that is seen amid activity. therefore, patients are at expanded hazard for the advancement of atrial arrhythmias and unexpected heart demise. About half of post-careful passings are unexpected and likely identified with the sudden beginning of polymorphic ventricular tachycardia/fibrillation (VT/VF). The administration of sinus hub brokenness and atrial arrhythmias can be testing. Both pacemaker arrangement and radiofrequency removal can be in fact difficult because of changed life structures and improvement of atrial scar tissue, while against arrhythmic drugs can hasten total heart block. Implantation of a pacemaker is suggested for kids with cutting edge second-degree heart square or complete atrioventricular square, patients with wiped out sinus disorder who are symptomatic or have utilitarian confinements, and those with tachy-bradycardia disorders with intermittent occasions in spite of antiarrhythmic medications. Symptomatic patients and those with an archived history of atrial fibrillation/fl express ought to experience electrophysiologic think about for inducible VT/VF with the situation of an implantable cardioverter-defibrillator (ICD) in those that have a positive EP study. Right ventricular brokenness is another wellrecognized, late intricacy related with the Senning or Mustard atrial-fix, as the morphologic RV fills in as the foundational ventricle. The etiology of RV brokenness isn't known; anyway, there are a few hypothesized speculations. The triangular state of the RV, in contrast with the slug molded LV, may render the RV deficient to support the fundamental

workload. The direction of the myocardial fibers may likewise add to RV brokenness. The RV endocardial and epicardial fibers are situated in a slanted way to the bearing of blood flow bringing about a peristaltic movement with constriction; when contrasted with the LV, wherein endocardial and epicardial fibers are oppositely arranged with a zone of round myocytes in the middle of bringing about a wringing movement with contraction. This separation in myocyte direction and contractile movement regularly enables the LV to work against high-obstruction systems. Widening of the RV with dynamic brokenness and annular enlargement might be one contributing component towards the improvement of RV brokenness; nonetheless, tricuspid spewing forth may hasten it as well. Under high-weight from the fundamental dissemination, the septal mass of the RV bows toward the low-weight LV in this way pulling the septal leaflet of the tricuspid valve. Thus, coaptation of the valve leaflets is lost and the valve winds up bumbling. Aspiratory hypertension can be another late inconvenience of an atrial-level fix with an occurrence announced at 7% in those patients getting by to adulthood. The nearness of a ventricular septal deformity and D-TGA fix following 1 year of age are hazard factors for the improvement of pneumonic hypertension. Signs and side effects of pneumonic hypertension frequently don't show up until adulthood, a consequence of the LV's ability to work against a high-opposition system. Given the significant perioperative hazard forced by pneumonic hypertension, its identification is vital to fruitful perioperative the board. The specific perioperative concerns identified with aspiratory hypertension have been widely investigated elsewhere. As noted above, patients may build up a few late inconveniences from the Senning and Mustard methods. With current cardiology care, a significant number of these patients are living into the second and third decade of life. In that capacity, there are an expanding number of patients who may require soporific consideration amid different sorts of surgeries. Given the mind-boggling nature of these patients and the potential for co-morbidities identified with the essential imperfection or procured amid life, an intensive preoperative assessment is important to guarantee the safe perioperative consideration of such patients. The 3 noteworthy issues identified with the life systems of the Mustard or Senning methodology which ought to be tended to preoperatively include: 1) an assessment of myocardial capacity given that the fundamental ventricle is the correct ventricle, 2) an assessment for the nearness of aspiratory hypertension, and 3) an assessment of the baffle for impediment or breaks. Starting assessment of these patients should start with a total cardiovascular

history, specifically addressing on exercise limit or prejudice, orthopnea or paroxysmal night time dyspnea, the nearness of palpitations, a recorded history of arrhythmia, and earlier syncope events. On physical examination, the precordium will be hyperdynamic attributable to the foundational outstanding burden of the correct ventricle. A holosystolic mumble is characteristic of foundational atrioventricular valve spewing forth or leftover VSD, while the nearness of a systolic launch mumble may demonstrate sub pulmonary steno sis. Basic post-atrial fix ECG findings incorporate sinus bradycardia or a junctional beat, proof of single or double chamber pacing, and indications of right ventricular hypertrophy. RVH can be identified on ECG by the nearness of right-hub deviation, enormous R-waves in leads V1 and AVR, and proof of right atrial amplification. ST-section discouragement and T-wave reversal in the privilege precordial leads might be demonstrative of extreme RVH. If accessible, earlier Holter screen chronicles ought to be assessed for paroxysmal atrial fi fibrillation/ flexpress, visit supra ventricular extra systoles, junctional escape musicality very still, or scenes of non-sustained ventricular tachycardia. In those patients with a pacemaker or ICD set up, extra perioperative administration and assessment is required so as to avert gadget related unfavorable occasions, for example harm to the gadget or leads, gadget disappointment, modifications in pacing conduct, improper conveyance of a defibrillatory stun, and coincidental change to reinforcement pacing modes. Electromagnetic impedence (EMI) is the most normally experienced issue intraoperatively and can happen because of electrocautery use, nerve triggers, evoked potential screens, fasciculations, and shuddering, enormous tidal volumes, outside defibrillation, attractive reverberation imaging, radiofrequency removal, extracorporeal stun wave lithotripsy, and electroconvulsive therapy. Current perioperative rules on heart musicality the board gadgets set up by the American Society of Anesthesiologists (ASA) are sketched out in reference. For the most part, in pacemaker-subordinate patients, the gadget will be reinvented to a nonconcurrent mode amid the perioperative period, and the counter tachyarrhythmia capacity of an ICD will be crippled. In our patient, after a discourse with the pediatric cardiology administration, it was chosen to leave the pacemaker in the VVI mode amid the case. Since the patient was pacemaker subordinate under 5% of the time, there was a constrained worry that EMI obstruction would result in significant bradycardia or no heart yield. The worry with the utilization of a magnet and transformation of the pacemaker into a VOO mode was the danger of

arrhythmia enlistment identified with an R on T marvel. In any case, we ensured that subsequent to situating we approached the pacemaker and a magnet in the room if its utilization ended up essential. The other part of our case that was one of a kind was the utilization of neurophysiologic observing including both somatosensory and engine evoked possibilities (see beneath). As these include electrical incitement, we ran a few trials before turning the patient inclined to check that we could evoke the required reactions at a low boost edge without meddling with pacemaker work. Maybe the most significant part of the preoperative assessment in a TGA persistent status-post atrial switch is an appraisal of the foundational ventricular capacity. Echocardiography, either by transthoracic or transesophageal course, assumes a fundamental job in the appraisal of right ventricular structure, capacity, and physiology. In the event that the patient's body habitués is manageable, Trans thoracic echocardiography (TTE) ought to be utilized to assess for myocardial perfusion deserts, ungraceful myocardial withdrawal, tricuspid valve spewing forth, and the nearness of an atrial baffle release, all of which can add to RV failure [18]. Hoffman et al exhibited that the utilization of subcostal TTE imaging planes, rather than standard apical planes, took into consideration the concurrent evaluation of the RV inflow and outflow tracts, just as right ventricular function. When looking at RV execution amid cardiopulmonary exercise and dobutamine stress testing, Li et al found that activity limit corresponded with foundational ventricle free divider outing very still and amid dobutamine stress. Depressed RV capacity and exercise limit might be the aftereffect of inducible ischemia, RV hypertrophy with expanded myocardial oxygen request or a powerlessness to build stroke volume because of fixed preload made by the obstruction of the intra-atrial baffle. The utilization of heart MRI has significantly decreased the requirement for TEE or cardiovascular catheterization. Gadoliniumenhanced MRI can be utilized to survey fundamental right ventricle work and assess regions of anomalous myocardium. The nearness of irregular myocardial districts on MRI relates with markers of unfavorable results including RV brokenness, poor exercise resilience, nearness of arrhythmias, and clinical disintegration after some time. B-type natriuretic peptide (BNP) has a place with a group of peptide hormones that are associated with the homeostasis of intravascular volume status, vascular obstruction, and myocardial capacity. BNP is delivered in the cardiovascular myocytes, is discharged into the flow because of expanded ventricular divider stress, and makes natriuresis and vasodilation counter the impacts of the rennin-angiotensin-aldosterone

framework (RAAS). BNP dimensions may likewise fill in as a biomarker of ailment seriousness with prognostic relevance in CHD patients. A stepwise increment in BNP has been appeared relate with expanding illness seriousness and diminishing fundamental ventricular capacity, to such an extent that a BNP esteem > 78 pg/ mL had a high prescient precision as far as generally speaking mortality. BNP levels as a pointer for infection seriousness may likewise be appropriate in the forecast of results for patients with aspiratory hypertension. Bernus et al found that changes in BNP after some time connected contrarily with cardiovascular record and emphatically with mean RAP, mean aspiratory vein weight, PCWP, and pneumonic vascular obstruction index. Notwithstanding the assessment of RV work, it is additionally critical to assess these patients for the nearness of aspiratory hypertension (PH) preoperatively. The most widely recognized showing indications of PH incorporate dyspnea on effort, weakness, chest torment, syncope, palpitations and lower furthest point edema. A correct heart catheterization (RHC), which is the highest quality level for assessment of pneumonic hypertension, ought to be considered. A few examinations have exhibited that kids with PH have a significant danger of perioperative cardiovascular difficulties including heart failure, aspiratory hypertensive emergency, and death. The perioperative administration of patients with PH can be testing. Patients on perpetual treatment for PH including intravenous prostacyclins (epoprostenol and trepostinil), phosphodiesterase inhibitors (sildenafil), endothelin foes (bosentan), and breathed in prostacyclin analogs (iloprost) ought to be proceeded on these meds all through the pre operative period. Intraoperatively, hypercarbia, alveolar hypoxia, fundamental hypoxemia, metabolic acidosis, and poisonous upgrades, for example, torment or aviation route instrumentation can trigger a fast ascent in PVR, and even a pneumonic hypertensive crisis. Ventilation techniques to limit aspiratory hypertensive triggers ought to be utilized and incorporate the utilization of high oxygen focuses, low tidal volumes (6 mL/kg anticipated body weight), a marginally raised respiratory rate to take into consideration gentle hypocarbia, and ideal dimensions of positive end-expiratory weight (5-10 cmH₂O) to diminish the level of atelectasis and keep up useful lingering limit. The meds used to initiate and keep up anesthesia intraoperatively can likewise influence PVR and the administration of PH. The unpredictable specialists constrict hypoxic pneumonic vasoconstriction in this manner expanding ventilation-perfusion jumbling. Isoflurane and sevoflurane are related to aspiratory vasodilation and are commonly acknowledged as sheltered

sedative segments in patients with PH. Besides, the utilization of benzodiazepines, narcotics, nitrous oxide, etomidate, neuromuscular blocking operators, and propofol are viewed as safe for use in patients with aspiratory hypertension, as they have next to zero impact on PVR. Moderate hyperventilation with 100% oxygen, alongside the inception of breathed in nitric oxide (iNO), are the first line of medications for intra operative PH. Senning and Mustard fix patients with both aspiratory hypertension and RV brokenness may benefit from the perioperative utilization of milrinone. In patients who have tricuspid disgorging (TR), notwithstanding RV brokenness, milrinone decreases SVR while keeping up a typical high pulse. Perioperatively, the effect of meds with negative inotropic properties including unstable analgesic specialists and β -adrenergic opponents ought to be considered. In the inclined position, there is an expansion in CVP, a decline in left ventricular end-diastolic width, and no general change in systolic function. As such, in patients with poor RV capacity experiencing medical procedure in the inclined position, in which volume the executives is significant for hemodynamic steadiness, the utilization of intraoperative TEE and CVP checking, constant blended venous oxygen immersion (SvO₂) might be utilized as a pointer of heart yield and oxygen consumption [39,40]. The utilization of persistent SvO₂ observing can likewise fill in as a manual for fluid and inotropic the board with the exhibit that SvO₂-guided treatment may decrease both emergency clinic remain and postoperative complications [41,43]. Intraoperative hemodynamic administration may likewise be significantly influenced by the patient's normal perioperative prescriptions, more specifically angiotensin-changing over catalyst inhibitors (ACE-I's) and angiotensin-receptor blockers (ARB's). These prescriptions estrange the renin-angiotensin-aldosterone framework (RAAS) bringing about direct thoughtful barricade, expanded bioavailability of characteristic vasodilators, restraint of angiotensin II, and diminished discharge of aldosterone and anti diuretic hormone. The clinical significance, especially identified with analgesic administration, is the hindrance of angiotensin II, which regularly works as an intense vasoconstrictor and improvement for arginine-vasopressin discharge; two substances which restrict the hypotensive impacts of soporific agents. Accordingly, patients treated with ACE-I's and ARB's in the perioperative setting are at higher danger of growing clinically significant hypotension after sedative induction. Fruitful treatment of RAAS foe hypotension requires satisfactory intravascular volume substitution and conceivably the utilization of vasopressin as the hypotension might be inert to

adrenergic agonists, for example, phenylephrine. Although there stays constrained accord on the continuation or withdrawal of RAAS opponents perioperatively, a few examinations have shown an expanded danger of stubborn hypotension when ACE-I's and ARB's are managed on the morning of surgery. As such, the creators of these investigations, just as a few foundations, prescribe that patients on interminable ACE-I and ARB treatment get their keep going portion of medicine on the day preceding medical procedure. Given these worries, our patient's typical portion of enalapril was not regulated the morning of medical procedure and vasopressin was promptly accessible in the working room in case of hypotension.

Preceding the inception of medical procedure, it is additionally critical to decide whether the patient ought to get bacterial endo carditis (SBE) prophylaxis. National Surgical Infection Prevention Project (NSIPP) and Centers for Disease Control (CDC) prescribe the organization of prophylactic anti-infection agents to diminish the danger of careful site infection. The utilization of intraoperative neuromonitoring, including somatosensory-evoked possibilities (SSEP's) and engine evoked possibilities (MEP's), can further affect the soporific administration of patients experiencing PSF. Since the blood supply to the spinal line originates from two separate supplies (the foremost and back spinal veins), MEP checking is utilized to evaluate the front spinal line for damage while SSEP observing is utilized to survey the dorsal spinal columns. All in all, MEP's are touchier to analgesic operators than SSEP's with longer pathways being the most delicate. Along these lines, the most diffifaction sign to achieve is lower limit MEP's. Our typical practice as exhibited by this patient is to utilize a solitary portion of a transitional acting neuromuscular blocking operator to encourage tracheal intubation and after that take into account unconstrained recuperation to allow MEP checking. Our intraoperative sedative routine included 0.5 MAC desflurane and sufentanil imbuement. Ultimately, techniques to decrease intraoperative blood misfortune, and thusly, difficulties related to enormous volume blood misfortune and allogeneic blood transfusions ought to be tended to. The transfusion of allogeneic blood items isn't without hazard; the potential for transmission of irresistible sicknesses, immunosuppression, transfusion-related intense lung damage, transfusion responses, graftversus-have malady, and expanded the danger of nosocomial diseases and pneumonia have all been perceived as unfavorable events. A few strategies exist to both limit the measure of blood misfortune and cutoff the

requirement for allogeneic transfusions, and they incorporate autologous transfusion treatment; intraoperative blood rescue; pharmacologic control of the coagulation course using epsilon-aminocaproic corrosive, tranexamic corrosive, desmopressin (DDAVP), or recombinant factor VIIa; and controlled hypotension. Proper patient situating, particularly when inclined, to limit the impedance of venous profit because of weight for the stomach area, and support of normothermia, can further decrease blood loss. Given our patient's steady hemodynamic status preoperatively, we utilized sufentanil in dosages as high as 0.5 µg/kg/min to keep up a MAP at 55-65 mmHg. In rundown, we present the perioperative contemplations in thinking about a patient with TGA who had experienced a Senning system amid PSF. Perioperative concerns incorporate not just those identified with the essential cardiovascular sore, yet in addition, those identified with the surgery and intraoperative neurophysiologic observing. Heart and hemodynamic worries of these patients incorporate the potential for RV brokenness, baffle breaks or obstacle, the nearness of aspiratory hypertension, and sinus hub brokenness with arrhythmias. The last may require the position of a pacemaker or ICD gadget. Contingent upon the patient's heart status, the effect of analgesic specialists on myocardial capacity, sinus hub capacity, and PVR ought to be considered. Extra concerns incorporate those identified with PSF including blood protection methods and the requirement for neurophysiologic checking.

REFERENCES:

1. Lo Y, Dan Y, Tan Y, et al. Intra-operative monitoring in scoliosis surgery with multi-pulse cortical stimuli and desflurane anesthesia. *Spinal Cord* 2004; 42:342-5.
2. Tobias J. Strategies for minimizing blood loss in orthopedic surgery. *Semin Hematol* 2004; 41:145-56.
3. Carson J, Altman D, Duff A, et al. Risk of bacterial infection associated with allogeneic blood transfusion among patients undergoing hip fracture repair. *Transfusion* 1999; 39:694-700.
4. Goodnough L, Brecher M, Kanter M, et al. Transfusion medicine: first of two parts - blood transfusion. *N Engl J Med* 1999; 340:438-47.
5. Parashurama, Doyle J, Lau W, et al. Transfusion-associated graft versus host disease. *PediatrCrit Care Med* 2002; 3:57-62.
6. Taylor R, Manganaro L, O'Brien J, et al. Impact of allogeneic packed red blood cell transfusion on nosocomial infection rates in the critically ill patient. *Crit Care Med* 2002; 30:2249-54.

7. Florentino-Pineda I, Blakemore L, Thompson G, et al. The effect of epsilon-aminocaproic acid on perioperative blood loss in patients with idiopathic scoliosis undergoing posterior spinal fusion. *Spine* 2001; 26:1147-51.
8. Schmid H, Kurz A, Sessler D, et al. Mild hypothermia increases blood loss and transfusion requirements during total hip arthroplasty. *Lancet* 1996; 347:289-92.
9. Warnes CA. Congenital heart disease for the adult cardiologist: Transposition of the great arteries. *Circulation* 2006; 114:2699-709.
10. Dos L, Teruel L, Ferreira JJ, et al. Late outcome of Senning and Mustard procedures for correction of transposition of the great arteries. *Heart* 2005; 91:652-6.
11. Kammeraad J, van Deursen C, Sreeram N, et al. Predictors of sudden cardiac death after Mustard or Senning repair for transposition of the great arteries. *J Am Coll Cardiol* 2004; 44:1095-1102.
12. Drago F, Silveti M, Grutter G, De Santis A. Long term management of atrial arrhythmias in young patients with sick sinus syndrome undergoing an early operation to correct congenital heart disease. *Europace* 2006; 8:488-94.
13. Villain E. Indications for pacing in patients with congenital heart disease. *Pace* 2008;31: s17-20.
14. Haddad F, Couture P, Tousignant C, Denault A. The right ventricle in cardiac surgery, a perioperative perspective: I. Anatomy, physiology, and assessment. *AnesthAnalg* 2009; 108:404-21.
15. Nielsen E, Smerup M, Agger P, et al. Normal right ventricular three-dimensional architecture, as assessed with diffusion tensor magnetic resonance imaging, is preserved during experimentally induced right ventricular hypertrophy. *Anat Rec* 2009; 292:640-51.
16. Carmosino MJ, Friesen RH, Doran A, Ivy DD. Perioperative complications in children with pulmonary hypertension undergoing noncardiac surgery or cardiac catheterization. *AnesthAnalg* 2007; 104:521-7. Strumpher J, Jacobsohn E. Pulmonary hypertension and right ventricular dysfunction: physiology and perioperative management. *J CardiothoracVascAnesth* 2011; 25:687-704.
17. Stone M, Apinis A. Current perioperative management of the patient with a cardiac rhythm management device. *SeminCardiothoracVascAnesth* 2009; 13:31-43.
18. Navaratnam M, Dubin A. Pediatric pacemakers and ICD's: how to optimize perioperative care. *PediatrAnaesth* 2011; 21:512-21.
19. Haddad F, Couture P, Tousignant C, Denault A. The right ventricle in cardiac surgery, a perioperative perspective: II. Pathophysiology, clinical importance, and management. *AnesthAnalg* 2009; 108:404-21.
20. Hoffman P, Szymanski P, Lubiszewska B, Rozanski J. Echocardiographic evaluation of the systemic ventricle after atrial switch procedure. The usefulness of subcostal imaging. *Cardiol J* 2008; 15:156-61. Li W, Hornung T, Francis D, et al. Relation of biventricular function quantized by stress echocardiography to cardiopulmonary exercise capacity in adults with mustard (atrial switch) procedure for transposition of the great arteries. *Circulation* 2004;110:1380-6.
21. Budts W, Scheurwegs C, Stevens A, et al. The future of adult patients after Mustard or Senning repair for transposition of the great arteries. *Int J Cardiol* 2006;113:209-14.
22. Giardini A, Lovato L, Donati A, et al. Relation between right ventricular structural alterations and markers of adverse clinical outcome in adults with systemic right ventricle and either congenital complete (after Senning operation) or congenitally corrected transposition of the great arteries. *Am J Cardiol* 2006; 98:1277-82.
23. Tobias J. B-type natriuretic peptide: diagnostic and therapeutic application in infants and children. *J Intensive Care Med* 2011; 26:183-95.
24. Knirsch W, Hausermann E, Fasnacht M, et al. Plasma B-type natriuretic peptide levels in children with heart disease. *Acta Paediatrica* 2011; 100:1213-6.
25. Trojnaraska O, Gwizdala A, Katarzynski S, et al. The BNP concentrations and exercise capacity assessment with a cardiopulmonary stress test in cyanotic adult patients with congenital heart disease. *Int J Cardiol* 2010; 139:241-7.
26. Giannakoulas G, Dimopoulos K, Bolger A, et al. Usefulness of natriuretic peptide levels to predict mortality in adults with congenital heart disease. *Am J Cardiol* 2010; 105:869-73.
27. Bernus A, Wagner B, Accurso F. Brain natriuretic peptide levels in managing pediatric patients with pulmonary arterial hypertension. *Chest*. 2009; 135:745-51.
28. Fischer L, Van Aken H, Burkle H. Management of pulmonary hypertension: physiological and pharmacological considerations for anesthesiologists. *AnesthAnalg* 2003; 96:1603-16.
29. Blaise G, Langleben D, Hubert B. Pulmonary arterial hypertension. *Anesthesiology* 2003; 99:1415-32.

30. Carmosino M, Friesen R, Doran A, Ivy D. Perioperative complications in children with pulmonary hypertension undergoing noncardiac surgery or cardiac catheterization. *AnesthAnalg* 2007; 104:521-7. Friesen R, Williams G. Anesthetic management of children with pulmonary arterial hypertension. *Ped Anesth*. 2008; 18:208-16.
31. Mossad E. Pro: intraoperative use of nitric oxide for the treatment of pulmonary hypertension in patients with congenital heart disease is effective. *J CardiothoracVascAnesth* 2001; 15:259-62.
32. Dietrich C, Tobias J. Intraoperative administration of nitric oxide. *J Int Care Med*. 2003; 18:146-9.
33. Haddad E, Lawson S, Johns R, Rich G. Use of inhaled nitric oxide perioperatively and in intensive care patients. *Anesthesiology* 2000; 92:1821-5.
34. Forrest P. Anaesthesia and right ventricular failure. *Anaesth Intensive Care* 2009; 37:370-85.
35. Priebe H. Effects of halothane on global and regional biventricular performances and on coronary hemodynamics before and during right coronary artery stenosis in the dog. *Anesthesiology* 1993; 78:541-52.
36. Palmisano B, Mehner R, Baker J, et al. Direct effects of halothane and isoflurane in infant rabbit hearts with right ventricular hypertrophy secondary to chronic hypoxemia. *AnesthAnalg*1995; 80:1122-8.
37. Soliman DE, Maslow AD, Bokesch PM, et al. Transoesophageal echocardiography during scoliosis repair: comparison with CVP monitoring. *Can J Anaesth*1998; 45:925-32.
38. Vincent J. Does central venous oxygen saturation accurately reflect mixed venous oxygen saturation? Nothing is simple, unfortunately. *Intensive Care Med* 1992; 18:386-7.
39. Naruns E. Clinical implications of intraoperative continuous mixed venous oxygen saturation monitoring. *AANA J*. 1989; 57:45-52.
40. Della Rocca G, Pompei L. Goal-directed therapy in anesthesia: any clinical impact or just a fashion. *Minerva Anesthesiol* 2011; 77:545-53.
41. Viale J, Annat G, Ravat F, et al. Oxygen uptake and mixed venous oxygen consumption during aortic surgery and the first three postoperative hours. *AnesthAnalg*1991; 73:530-5.
42. Bernard J, Le Penven-Henninger C, Passuti N. Sudden decreases in mixed venous oxygen saturation during posterior spinal fusion. *Anesth* perioperative management in the transposition of great arteries. *Analg*1995; 80:1038-41.
43. Varin R, Mulder P, Tamion F, et al. Improvement of endothelial function by chronic angiotensin-converting enzyme inhibition in heart failure: role of nitric oxide, proteinoids, oxidant stress, and bradykinin. *Circulation* 2000; 102:351-6.
44. Auron M, Harte, B, Kumar A, Michota F. Renin-angiotensin system antagonists in the perioperative setting: clinical consequences and recommendations for practice. *Postgrad Med J* 2011; 87:472-81.
45. Colson P, Ryckwaert F, Coriat P. Renin-angiotensin system antagonists and anesthesia. *AnesthAnalg* 1999; 89:1143-55.
46. Behnia R, Molteni A, Iqic R. Angiotensin-converting enzyme inhibitors: mechanism of action and implications in anaesthesia practice. *Curr Pharm Des* 2003; 9:763-76.
47. Coriat P, Richer C, Douraki T, et al. Influence of chronic angiotensin-converting enzyme inhibition on anaesthetic induction. *Anesthesiology* 1994; 81:299-307.
48. Bertrand M, Godet G, Meersschaert K, et al. Should the angiotensin II antagonists be discontinued before surgery? *AnesthAnalg*2001; 92:26-30.
49. Comfere T, Sprung J, Kumar M, et al. Angiotensin system inhibitors in a general surgical population. *AnesthAnalg*2005; 100:636-44.
50. Wilson W, Taubert K, Gewitz M, et al. Prevention of infective endocarditis: guidelines from the American Heart Association: A guideline from the American Heart Association rheumatic fever, endocarditis, and Kawasaki disease committee, council on cardiovascular disease in the young, and the council on clinical cardiology, council on cardiovascular surgery and anesthesia, and the quality of care and outcomes research interdisciplinary working group. *Circulation* 2007; 116:1736-54.
51. Young B, Ng T, Teng C, et al. Non-concordance with surgical site infection prevention guidelines and rates of surgical site infections for general surgical, neurological, and orthopedic procedures. *Antimicrob Agents Chemother* 2011; 55:4659-63.
52. Deiner S. Highlights of anaesthetic considerations for intra operative neuro monitoring. *SeminCardiothoracVascAnesth*2010; 14:51-3.
53. Sloan T and Heyer E. Anesthesia for intraoperative neurophysiologic monitoring of

- the spinal cord. *J Clin Neurophysiol* 2002; 19:430-43.
54. Balvin M, Song K, Slimp J. Effects of anesthetic regimens and other confounding factors affecting the interpretation of motor evoked potentials during pediatric spine surgery. *Am J Electro neurodiagnostic Technol* 2010; 50:219-44.
 55. Reinacher P, Priebe H, Blumrich W, et al. The effects of stimulation pattern and sevoflurane concentration on intraoperative motor-evoked potentials. *Anesth Analg* 2006; 102:888-95.
 56. Mathews TJ, Miniño AM, Osterman MJK, et al. Annual summary of vital statistics: 2008. *Pediatrics* 2011; 127:146-57.
 57. van der Bom T, Zomer AC, Zwinderman AH, Meijboom FJ, Bouma BJ, Mulder BJ. The changing epidemiology of congenital heart disease. *Nat Rev Cardiol* 2011; 8:50-60.
 58. Anderson R and Weinberg P. The clinical anatomy of transposition. *Cardiol Young* 2005; 15(suppl 1):76-87.
 59. Senning A. Surgical correction of transposition of the great vessels. *Surgery* 1959; 45:966-80.
 60. Quaegebeur JM, Rohmer J, Brom AG. The revival of the Senning operating for treatment of transposition of the great arteries. *Thorax* 1977; 32:517-24.
 61. Love BA, Mehta D, Fuster VF. Evaluation and management of the adult patient with transposition of the great arteries following atrial-level (Senning or Mustard) repair. *Nat Clin Pract Cardiovasc Med* 2008; 5:454-67.