



CODEN [USA]: IAJ PBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES<http://doi.org/10.5281/zenodo.3569665>Available online at: <http://www.iajps.com>

Research Article

**FREQUENCY OF RAISED BODY MASS INDEX (BMI) IN
PATIENTS PRESENTING WITH MYOCARDIAL INFARCTION**

Yusra Iqbal, Saadia Sehrish, Maria Anwar

Abstract:

Objectives: To determine the frequency of raised BMI in patients presenting with myocardial infarction. **Material and methods.** This was a cross sectional study that was conducted at multicentre in Pakistan during May 2018 to December 2018, in which 100 cases of acute myocardial infarction (Diagnosed with ECG changes with raised Troponin T levels) were included. Socio demographic data like age, gender, BMI and relevant clinical data i.e. DM, HTN and type of MI (STEMI or NSTEMI) were taken. Raised BMI was labelled when it was more than 25 kg/m². **Results;** In this study there were total 100 cases out of which 60 (60%) were males and 40 (40%) females with mean age of 53.14±5.76 years. There were 32 (32%) cases with DM and 40 (40%) with HTN while 72 (72%) had ST elevation MI. Raised BMI was seen in 40 (40%) of cases. This was significantly high in females where it affected 24 (60%) of cases as compared to 16 (26.67%) females with $p=0.01$. Raised BMI was common with DM but had equal distribution in cases with or without HTN. There was almost equal distribution of raised BMI in cases of both STEMI and NSTEMI with $p=0.98$. **Conclusion;** Raised BMI is an independent risk factor in cases of acute coronary syndrome. It is significantly high in number in female cases.

Key words; MI, Raised BMI, DM, HTN**Corresponding author:**

Yusra Iqbal,

QR code



Please cite this article in press Yusra Iqbal et al., *Frequency Of Raised Body Mass Index (BMI) In Patients Presenting With Myocardial Infarction.*, Indo Am. J. P. Sci, 2019; 06(12).

INTRODUCTION:

Every year, more than 6 million patients present to the emergency department with chest pain, and the majority are subsequently admitted with concern for Myocardial infarction (MI), which is an acute emergency. MI is diagnosed either immediately with specific ECG changes or needs confirmation with the help of various cardiac enzymes like Troponin T, Troponin I, Creatinine Kinase –MB (CK-MB) or Echocardiography if needed. On the basis of ECG changes and cardiac enzymes, it can be subdivided into two groups, which include ST segment elevation MI (STEMI) and Non ST segment elevation MI (NSTEMI).¹

MI can lead to different complications that can be fatal and include cardiogenic shock, arrhythmias, progressive heart failure, mechanical cardiac complications and repeat MI.^{2, 3} A similar trend was noted in an analysis of data on 2.5 million patients from the National Registry of Myocardial Infarction (NRFMI).⁴

There are numerous well documented risk factors associated with high risk of MI which include DM, HTN, Family history, Smoking, raised BMI.^{5, 6} Few of them like, smoking, HTN, DM and obesity are reversible and to deal with them can decrease the chances of morbidity and mortality in such cases.^{7, 8}

Regarding BMI, it is associated with more chances of DM, high insulin resistance, increase oxygen demand and dyslipidemias which are again risk factors for increase mortality. In contrast to this few studies have shown rather beneficial effect of raised BMI and are known as obesity paradox.⁹

OBJECTIVES:

To determine the frequency of raised BMI in patients presenting with acute myocardial infarction.

MATERIALS AND METHODS:

This was a cross section study that was conducted at multicentre in Pakistan during May 2018 to December 2018 in which 100 cases of acute myocardial infarction (Diagnosed with ECG changes with raised Troponin T levels) were included. Socio demographic data like age (in years), gender (male/female), height (in meters), weight (in kg) and BMI (kg/meter² on admission) were taken and relevant clinical data was also collected i.e. DM (yes/no), HTN (yes/no) and type of MI (STEMI or NSTEMI). Raised BMI was labelled when it was

more than 25 kg/m².

Sample selection;

The cases were selected via non-probability, consecutive sampling with following criteria.

Inclusion criteria:

1. Both genders
2. Age 30 to 60 years
3. STEMI and NSTEMI

Exclusion Criteria:

1. Age less than 30 or more than 60 years
2. Cases with normal ECG.
3. Cases undergoing any surgical intervention during first 30 days.
4. Mortality due to any other cause except for cardiac event (like road traffic accident)

DATA ANALYSIS PROCEDURE:

The data was entered and analysed by using SPSS-17. Quantitative variables like age and BMI were assessed in mean \pm SD. Qualitative variables like gender, DM, HTN, raised BMI (yes/no) and type of MI were presented as frequencies and percentages. Stratification was done on the basis of age, gender, DM, HTN and type of MI to see its effect on outcome variable i.e. raised BMI. Post stratification chi-square test was applied and $p < 0.05$ was taken as significant.

RESULTS:

In this study there were total 100 cases out of which 60 (60%) were males and 40 (40%) females with mean age of 53.14 ± 5.76 years. There were 55 (55%) cases with age more than 50 years. There were 32 (32%) cases with DM and 40 (40%) with HTN while 72 (72%) had ST elevation MI. Raised BMI was seen in 40 (40%) of cases. This was significantly high in females where it affected 24 (60%) of cases as compared to 16 (26.67%) females with $p = 0.01$. This was more common in age group of 35 to 50 years ($p = 0.68$) as shown in table 1. Raised BMI was common with DM but had equal distribution in cases with or without HTN as in table 2. There was almost equal distribution of raised BMI in cases of both STEMI and NSTEMI (Table 3) with $p = 0.98$.

DISCUSSION:

Acute myocardial infarction is a life-threatening condition and commonly encountered entity in the emergency and cardiac settings. There are multiple modifiable and non-modifiable risk factors. Obesity is one of the most common of them and raised BMI is

considered as most widely used tool to label it.

In this study the raised BMI was seen in 40 (40%) of cases. This was similar to studies done by Khan HS et al¹⁰ and Parsa AF et al.¹¹ However, the latter found a high significance in cases with raised BMI and severity of disease with $p=0.001$. Why this difference was found, it may be due to ethnic difference, because the study conducted in Pakistan had similar results to ours while this conducted in Africa may be thought to interfere with racial differences.

Raised BMI was significantly high in females where it affected 24 (60%) of cases as compared to 16 (26.67%) females with $p=0.01$ in cases of MI. This was also observed by the studies done by Walker SP

and Rubinshtein R et al that also found high number of females with raised BMI.¹²⁻¹³ The reason of high number of females can be due to endocrine causes and the life styles of the females in our region, as the males are more active and do the physical activity as compared to females, so it was seen higher in females.

Raised BMI was common with cases of DM. Many studies have reported this in their results.¹⁴⁻¹⁵ The high number of ACS cases with raised BMI having co morbid of DM can be explained by the increased lipogenesis and deposition of fat in the subjects.

CONCLUSION;

Raised BMI is an independent risk factor in cases of acute coronary syndrome. It is significantly high in number in female cases.

TABLE 1: RAISED BMI WITH RESPECT TO DEMOGRAPHICS
n= 100

Gender	Raised BMI		Total	Significance
	Yes	No		
Male	16 (26.67%)	44 (73.33%)	60	$p=0.01$
Female	24 (60%)	16 (40%)	40	
Age groups	Raised BMI		Total	Significance
	YES	NO		
35-50	21 (46.67%)	24 (53.33%)	45	$p=0.68$
>50	19 (34.55%)	36 (65.45%)	55	

TABLE 2: RAISED BMI WITH RESPECT TO COMORBIDITIES
n= 100

DM	Raised BMI		Total	Significance
	Yes	No		
Yes	15 (46.88%)	17 (53.12%)	32	$p=0.43$
No	25 (36.76%)	43 (63.24%)	68	
HTN	Raised BMI		Total	Significance
	YES	NO		
Yes	16 (40%)	24 (60%)	40	$p=1.0$
No	24 (40%)	36 (60%)	60	

TABLE 3: RAISED BMI WITH RESPECT TO TYPE OF MI
n= 100

Type of ACS	Raised BMI		Total
	Yes	No	
STEMI	28 (38.89%)	44 (61.11%)	72 (30%)
NSTEMI	12 (42.86%)	16 (57.14%)	28 (70%)
Total	40 (100%)	60 (100%)	100 (100%)

$p=0.98$

REFERENCES:

- 1) Rogers WJ, Frederick PD, Stoehr E. Trends in presenting characteristics and hospital mortality among patients with ST elevation and non-ST elevation myocardial infarction in the National Registry of Myocardial Infarction from 1990 to 2006. *Am Heart J*. 2008;156(6):1026-29.
- 2) Asif MM, Khan MI, Tareen ZK, Khan SB, Khattak MU, Ahmed M, et al. Frequency of risk factors of coronary artery disease in tertiary care hospital. *Khyber J Med Sci*. 2014;7(1):66-71.
- 3) Faisal A, Ayub M, Waseem T, Shahzad R, Khan AT, Hasnain SS. Risk factors in young patients of acute myocardial infarction. *J Ayub Med Coll Abbottabad*. 2011;23(3):10-13.
- 4) Wu H, Apple FS, Gibler WB, Jesse RL, Warshaw MM, Valder R. National academy of clinical biochemistry standards of laboratory practice; recommendations for the use if cardia markers in coronary artery disease. *Clin chem*. 1999;45(7):1104-21.
- 5) Chen Z, Yang G, Offer A, Zhou M, Smith M, Peto R, et al. Body mass index and mortality in china: a 15-year prospective study of 220000 men. *Int J Epidemiol*. 2012;41(2):472-81.
- 6) Ashwell M, Mayhew L, Richardson J, Rickayzen B. Waist-to-height ratio is more predictive of years of life lost than body mass index. *PLoS ONE*. 2014;9(9):e103483.
- 7) Camprubi M, Cabrera S, Sans J, Vidal G, Salvad T, Bardaj A. Body mass index and hospital mortality in patients with acute coronary syndrome receiving care in a university hospital. *Obesity*. 2012;5(2):02-05.
- 8) Dooley J, Chang AM, Salhi R, Judd E. Relationship between body mass index and prognosis of patients presenting with potential acute coronary syndromes. *Acad Emerg Med*. 2013;20(9):904-10.
- 9) Angeras O, Albertsson P, Karason K, Ramunddal T, MatejkaG, James S, et al. Evidence for obesity paradox in patients with acute coronary syndromes: a report from the Swedish coronary angiography and angioplasty registry. *Eur Heart J*. 2013;34(3):345-53.
- 10) Khan HS, Javed A, Aziz S, Ali J. Relationship between BMI and severity of coronary artery disease in female population of Pakistani origin. *Pak Heart J*. 2011;44:1-2.
- 11) Parsa AF, Jahanshahi B. Is the relationship of body mass index to severity of coronary artery disease different from that of waist-to-hip ratio and severity of coronary artery disease? Paradoxical findings. *Cardiovasc J Afr*. 2015 Jan-Feb;26(1):13-6.
- 12) Walker SP, Rimm EB, Ascherio A, Kawachi I, Stampfer MJ, Willett WC: Body size and fat distribution as predictors of stroke among US men. *Am J Epidemiol*. 1996, 144: 1143-1150.
- 13) Rubinshtein R, Halon DA, Jaffe R, Shahla J, Lewis BS. Relation between obesity and severity of coronary artery disease in patients undergoing coronary angiography. *Am J Cardiol*. 2006;97(9):1277-1280.
- 14) Alpert JS, Thygesen K, Antman E, Bassand JP. Myocardial infarction redefined- a consensus document of the joint European society of Cardiology/ American college of cardiology committee for the redefinition of MI. *J Am Coll Cardiol*. 2000;36(3):959-69.
- 15) Khosla T, Lowe CR. Indices of obesity derived from body weight and height. *Br J Prev Soc Med*. 1967;21(4):122-28.