



CODEN [USA]: IAJPBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

Available online at: <http://www.iajps.com>

Review Article

REVIEW OF APPROPRIATE APPROACH IN PREVENTION OF CARDIOVASCULAR DISEASE

¹Dr. Thamer Matoog Alsufyani, ²BANDAR OTHMAN ALZAHIRANI, ³Alaa Adel Hussen Alsabbagh, ⁴Mazen saeed takroony, ⁵Omar Mohammad Ahmed Hinaidi, ⁶Abdullah Faisal Al-Hazmi, ⁷Saud Abdulaziz Salh Zainy., ⁸Safar Othman Ahmad Alzahrani, ⁹NABEEL FAWZI SHAIKH., ¹⁰Ahmed Nawwar Al-otaibi.

¹General practitioner, Medicine and medical sciences, Prince Ahmed scheme health center.

²EPIDEMIOLOGY, public health, Prince Ahmed scheme health center

³Technician-Nursing, Chronic Diseases, Prince Ahmed scheme health center

⁴Nursing, Primary health care, Prince Ahmed scheme health center.

⁵Nursing, Chronic Diseases, Prince Ahmed scheme health center.

⁶Pharmacy Technician, Prince Ahmed Primary Health Care Center.

⁷Nurse, Environmental Health Department in Makkah Health Cluster / medical waste. Health Center Batha Quraish.

⁸ANASTHESIA, Operation Department, KING FAISAL HOSPITAL

⁹ANASTHESIA, Operation Department, KING FAISAL HOSPITAL.

¹⁰Specialist in Management of health Services and Hospitals. Community Participation Program, Specialist in Management of Health Services and Hospitals, Makkah.

Article Received: December 2019 **Accepted:** December 2019 **Published:** December 2019

Abstract:

Background: Cardiovascular disease (CVD), disease that affects the heart and vessels, includes raised blood pressure, coronary cardiovascular disease (CHD), heart failure.

Objectives: This narrative review was aimed to discuss the diagnostic and management approaches to cardiac diseases, and the ability to prevent it in primary healthcare setting.

Methodology: literature search through most important electronic medical databases for all studies published up to Nov, 2019, discussing the preventive measures taking in primary healthcare.

Conclusion: Huge number of CVD patients as noncontagious chronic diseases are treated in primary health center. Primary healthcare provider have very important role in cardiovascular risk management. Considering that in our country preventive programs are not at acceptable level, more comprehensive guidelines should be implemented through healthcare setting to increase the preventive levels of cardiovascular diseases. Number of CVD patients in primary healthcare increases the importance of healthcare teams to decrease the mortality and comorbidities in future.

Corresponding author:

Dr. Thamer Matoog Alsufyani,
General practitioner,
Medicine and medical sciences,
Prince Ahmed scheme health center.

QR code



Please cite this article in press Thamer Matoog Alsufyani et al., *Review Of Appropriate Approach In Prevention Of Cardiovascular Disease.*, Indo Am. J. P. Sci, 2019; 06[12].

INTRODUCTION:

Cardiovascular disease (CVD), disease that affects the heart and vessels, includes raised force per unit area, coronary disorder (CHD), heart condition (1). Roughly 82 million Americans have several kinds of CVD. In medical aid 0.7% to 2.7% of client encounters are because of pain (2). However, the frequency of severe heart illness in these patients, e.g., chronic steady CHD or intense coronary syndrome (ACS), is low. In unselected patients providing with hurting in medical aid, the final frequency of coronary heart problem is in between 12.8 and 14.6% (2,3).

In Europe area CVD are liable for 4, 3 million death by year, which suggests 48% of all death triggers (54% female deaths and 43% male deaths). In EU Country (ECC) are in command of 42% deaths. CVD are the first cause in male's death in Europe too apart from Spain, France and Netherlands (4). but half CVD deaths is brought on by ischaemic hart diseases and one third by cerebrovascular illness. Last thirty years in an exceedingly lot of European countries, West and South Europe mortality, incidence and lethality of CVD are reduced rather in East and Middle Europe where are increased. Nations with greater CVD are countries with bad financial advancement (4,5).

Heart failure may be a progressively important problem for medical care doctors during a lot of healthcare systems in industrialized nations. The condition is nearly as common as diabetes in older adults, occurring in an exceedingly minimum of twenty-two of the adult population and increasing to three in those aged over 75 years (6). The incidence of the many cardiovascular diseases has actually declined over the past 20 years, the occurrence of cardiopathy has continued to extend, due partially to the actual fact that more individuals are enduring after intense myocardial infarctions and likewise to the increasing number of senior individuals (6). failure is difficult to detect properly on clinical premises. Only 26% of clients with suspected cardiopulmonary arrest described a fast gain access to center for echocardiography had the diagnosing verified after examination. Clinical diagnosis by treatment room physicians is just as poor (7).

Precise and early diagnosis is extremely important in medical aid, but single signs and signs are rarely sufficient to dependably diagnose CHD. This issue could also be gotten eliminate by establishing a forecast rule that mixes numerous symptoms,

indications, and other patient qualities like sex, age and coronary risk elements.

Objective: the aim of this narrative review was to discuss the ability of primary healthcare provider and general practitioner to prevent the main cardiovascular diseases such as coronary heart disease (CHD), and heart failure in primary care setting, through reviewing most evidence-based studies in the related topic.

METHODOLOGY:

A literature search was conducted in November 2019 using PubMed, CINAHL Plus, and PsycINFO. In PubMed the following search terms were used: prevention, primary care, healthcare setting, heart disease, chronic heart diseases, cardiovascular diseases. The reference list of articles was also searched, identified by the search strategy and those selected that were relevant. Selected review articles and meta-analyses were included because they provide comprehensive overviews that may be beyond the scope of this article.

DISCUSSION:

Cardiovascular diseases are preventable in a great deal of patients. World Health Organization (WHO) approximated that middle phase of blood pressure decrease, obesity, cholesterol and tobacco usage, in population level, more than half decreased CVD incidence (4). Action frame for avoidance CVD area is based on great recognized aims, select by management method and capacity advancement for their realization. Understood approaches for leading CVD programs contain practically frame objectives: hearth health promotion in entire population, symptoms of CVD avoidance, life promotion without tobacco, routine body exercise, hypertension danger aspects obliteration, reducing greater lipids level, battle against abdominal type weight problems and diabetes, specific diagnostically procedures, CVD control and therapy, program versus regression of CVD, without tension life, healthy food and healthy and comfy environment (8). Global technique is the final goal of these actions to achieve less frequency in morbidity and mortality from CVD with lower CVD risks and their determinants, to develop logical and equivalent health developments for CVD handling, CVD trend follow up and their threat aspects from modern-day world (4,8).

Rules of early diagnosis and management:

A potential diagnostic art failure of aid in primary care is the evaluation of clients by determining plasma concentrations of brain natriuretic peptide. Data on the credibility of brain natriuretic peptide are

contrasting. Brain natriuretic peptide testing had a sensitivity of 97%, an uniqueness of 84%, a favorable predictive value of 70%, and a negative predictive worth of 98% in 106 clients with signs of recent onset who were described a fast access cardiac arrest center (9). Similar predictive efficiency for the peptide was reported both in clients with validated left ventricular systolic dysfunction (the commonest cause of heart failure) in a population of 1653 adults aged between 25 and 75 years who were evaluated in Glasgow, a location with high rates of heart disease, (10) and in another paper in the BMJ in which 155 clients aged over 75 were evaluated in medical care (11). However, these research studies were small, and a research study of 134 patients who were stable after a myocardial infarction discovered that brain natriuretic peptide might not anticipate the existence of mild to moderate left ventricular systolic dysfunction compared with regular function (McClure SJ *et al*, 20th Congress of the European Society of Cardiology, Vienna, 1998). For 126 patients in basic practice who were referred to an echocardiography center, in another recent paper in the BMJ, there was just a small diagnostic benefit in including brain natriuretic peptide to standard investigations of electrocardiography and chest radiographs, although 1 in 7 patients was offered a false negative diagnosis.¹⁵ However, these latter findings contrast with a negative predictive worth of 98% reported in an accompanying paper on brain natriuretic peptide that appeared in the very same issue.¹⁴

Cardiovascular diseases are avoidable in large number of clients. World Health Organization (WHO) estimated that middle phase of high blood pressure reduction, weight problems, cholesterol and tobacco usage, in population level, over half lowered CVD incidence (7). Action frame for avoidance CVD area is based on good recognized aims, pick by management strategy and capability development for their realization. Understood methods for leading CVD programs include practically frame objectives: hearth health promo in entire population, symptoms of CVD prevention, life promotion without tobacco, routine body workout, hypertension risk elements elimination, reducing higher lipids level, battle against stomach type weight problems and diabetes, exact diagnostically treatments, CVD control and treatment, program versus regression of CVD, without stress life, healthy food and healthy and comfortable environment (4). Global technique is the final objective of these actions to accomplish less frequency in morbidity and mortality from CVD with lower CVD dangers and their determinants. To develop rational and equal health innovations for

CVD managing, CVD trend follow up and their risk factors from modern world (8,12).

Availability of diagnostic methods in primary healthcare setting:

To positively establish a diagnosis of heart failure in primary care, most patients need to be referred for cardiac imaging. A recent study of a random sample of primary care doctors throughout six European nations reported only 5% (Netherlands) to 37% (United Kingdom) of general practitioners had direct access to echocardiography (9). This poor access is partly due to an absence of skilled personnel to run clinics and partially to issues that providing primary care physicians with direct gain access to would result in inappropriate usage (although a trial of open access echocardiography found just 12% of recommendations to be "inappropriate") (7,8).

Are there alternatives to echocardiography in primary care? A regular electrocardiogram usually omits left ventricular dysfunction (10). Nevertheless, changes might be subtle and medical care physicians' lack of ability in interpreting electrocardiograms might suggest that referral for specialist opinion is still needed.

A prospective diagnostic help in primary care is the assessment of clients by determining plasma concentrations of brain natriuretic peptide. However, information on the credibility of brain natriuretic peptide are conflicting. Brain natriuretic peptide screening had a sensitivity of 97%, a specificity of 84%, a positive predictive worth of 70%, and an unfavorable predictive worth of 98% in 106 clients with signs of current beginning who were described a fast gain access to heart failure clinic. Comparable predictive performance for the peptide was reported both in clients with confirmed left ventricular systolic dysfunction (the commonest cause of heart failure) in a population of 1653 adults aged in between 25 and 75 years who were screened in Glasgow, an area with high rates of heart disease, and in another paper in the BMJ in which 155 clients aged over 75 were screened in primary care (6,9,11). However, these studies were little, and a research study of 134 clients who were steady after a myocardial infarction discovered that brain natriuretic peptide might not predict the presence of moderate to moderate left ventricular systolic dysfunction compared with regular function. For 126 patients in general practice who were described an echocardiography clinic, in another recent paper in the BMJ, there was only a little diagnostic benefit in adding brain natriuretic peptide to standard investigations of

electrocardiography and chest radiographs, although 1 in 7 patients was given a false negative medical diagnosis (9,11). However, these latter findings contrast with a negative predictive value of 98% reported in an accompanying paper on brain natriuretic peptide that appeared in the exact same issue.

In one research study all CVD signs of medical services management are filled with higher portion of very little level which showed that in family physician group, quality medical services management is great however could be even much better. Brand-new patients with angina pectoris percentage sent out to ergometry or/and cardiologist remained in 51.72%, high blood pressure values \leq 140/90 achieved in 58.62% clients. Overall cholesterol values \leq 5mmol/l was achieved just in 27.58% patients, what is very little filled level and it is essential to operate in clients education about healthy life style, healthy food and much better specific health care. In British nationwide research study about CVD, high blood pressure and stroke patients quality care in primary health care, 71.4% -88.6% clients with brand-new diagnose of AP were sent out to ergo test or cardiologist, what was better identify verification quality management of AP (13).

Clients with CVD assessment with patient's file registration Aspirin treatment administration in last 15 months or alternative anticoagulant therapy (other than with clear contraindications) remained in 68.96% (according to recognized standards minutes.25%, max level was 90%). Comparable outcomes existed in research study (14) about CVD risk quality management realized in primary healthcare in some European nations which revealed that 80% clients with cardiac failure were with anticoagulant treatment. Clients with CVD percentage treated with β -blockers in last 15 months remained in 79.31%. Indicator is filled what revealed excellent quality management treatment of CVD (min. 25%, max levels was 80%). According to Brenan and others study, revealed that even half of severe myocardial infarct patients did not get Aspirin or β -blocers, and mistakes could be fatal and could be priceless and cause more costs than life and could be result of irresponsible practice (15).

CONCLUSION:

Huge number of CVD patients as noncontagious chronic diseases are treated in primary health center. Primary healthcare provider has very important role in cardiovascular risk management. Considering that in our country preventive programs are not at acceptable level, more comprehensive guidelines

should be implemented through healthcare setting to increase the preventive levels of cardiovascular diseases. Number of CVD patients in primary healthcare increases the importance of healthcare teams to decrease the mortality and comorbidities in future.

Primary healthcare physicians need structured approaches to diagnosing heart failure; these approaches should involve stratifying patients into risk groups and assessing them with objective tests. In many cases, there is no immediate alternative to echocardiography to confirm the diagnosis and determine the etiology. However, the brain natriuretic peptide assay holds an exciting potential to determine who should be listed for echocardiography, with the possibility that it can aid in treatment and predicting prognosis.

REFERENCES:

1. Imes CC, Lewis FM. Family history of cardiovascular disease, perceived cardiovascular disease risk, and health-related behavior: a review of the literature. *J Cardiovasc Nurs.* 2014;29(2):108-129. doi:10.1097/JCN.0b013e31827db5eb.
2. Verdon F, Herzig L, Burnand B, Bischoff T, Pecoud A, Junod M, Muhlemann N, Favrat B. Chest pain in daily practice: occurrence, causes and management. *Swiss Med Wkly.* 2008;138(23-24):340-347.
3. Bösner S, Becker A, Haasenritter J, Abu Hani M, Keller H, Sönnichsen AC, Karatolios K, Schaefer JR, Seitz G, Baum E. et al. Chest pain in primary care: epidemiology and pre-work-up probabilities. *Eur J Gen Pract.* 2009;15(3):141-146. doi: 10.3109/13814780903329528.
4. Beganlic A, Pavljasevic S, Kreitmayer S, et al. Qualitative Evaluation of Cardiovascular Diseases Management in Family Medicine Team in One Year Level. *Med Arch.* 2015;69(3):140-144. doi:10.5455/medarh.2015.69.140-144.
5. Anonymous. Copenhagen: WHO-Regional Office for Europe; 2006. Gaining Health, The European Strategy for the Prevention and Control of Noncommunicable Diseases. WHO.
6. Hobbs R. Can heart failure be diagnosed in primary care?. *BMJ.* 2000;321(7255):188-189. doi:10.1136/bmj.321.7255.188
7. Friedlander Y, Siscovick DS, Arbogast P, et al. Sudden death and myocardial infarction in first degree relatives as predictors of primary cardiac arrest. *Atherosclerosis.* 2002;162(1):211-216.
8. Lieshout van J, Grol R, Campbell S, Falcoff H, Capell EF, Glehr M, Goldfracht M, Kumpusalo E, Künzi B, Ludt S, Petek D, Vanderstighelen V,

- Wensin M. Cardiovascular risk management in patients with coronary heart disease in primary care: variation across countries and practices. An observational study based on quality indicators. *BMC Family Practice*. 2012;13:96.
9. Cowie MR, Struthers AD, Wood DA, Coates AJS, Thompson SG, Poole-Wilson PA, et al. Value of natriuretic peptides in assessment with patients with possible new heart failure in primary care. *Lancet*. 1997;350:1349–1353.
 10. McDonagh TA, Robb SD, Murdoch DR, Morton JJ, Ford I, Morrison C, et al. Biochemical detection of left ventricular systolic dysfunction. *Lancet* 1998;9-13
 11. Smith H, Pickering RM, Struthers A, Simpson I, Mant D. Biochemical diagnosis of ventricular dysfunction in elderly patients in general practice: observational study. *BMJ*. 2000;320:906–908.
 12. Saxen S, Car J, Eldred D, Soljak M, Majeed A. Practice size, caseload, deprivation and quality of care of patients with coronary heart disease, hypertension and stroke in primary care: national cross-sectional study. *BMC Health Services Research*. 2007;7:96.
 13. Anonymous. Akreditacijski standardi za timove Porodične/Obiteljske medicine. Sarajevo: AKAZ; 2012. Agencija za kvalitet i akreditaciju u zdravstvu u FBiH.
 14. Lieshout van J, Grol R, Campbell S, Falcoff H, Capell EF, Glehr M, Goldfracht M, Kumpusalo E, Künzi B, Ludt S, Petek D, Vanderstighelen V, Wensin M. Cardiovascular risk management in patients with coronary heart disease in primary care: variation across countries and practices. An observational study based on quality indicators. *BMC Family Practice*. 2012;13:96.
 15. Brennan TA, Leape LL, Laird NM, et al. Incidence of adverse events and negligence in hospitalized patients. Results of the Harvard Medical Practice Study I. *N Engl J Med*. 1991;324:370–376.