



CODEN [USA]: IAJ PBB

ISSN : 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4316421>Available online at: <http://www.iajps.com>

Research Article

**PREVALENCE OF CANDIDA ISOLATES FROM URINE OF
PATIENTS**Dr Misbah Arshad¹, Dr Naveed Irshad¹, Dr Kashif Javed²¹DHQ hospital Faisalabad, ²DHQ Teaching Hospital Sargodha**Article Received:** October 2020 **Accepted:** November 2020 **Published:** December 2020**Abstract:**

Background: Worldwide, urinary tract infection is found most common problem caused by microbial invasion to different tissues of urinary tract. There has been a marked increase in fungal infections involving the urinary tract, of which candida species are the most prevalent.

Objective: To determine the prevalence of candida isolates from urine of patients reporting to Hospital.

Methods: A cross-sectional study was carried out in DHQ Hospital, Faisalabad during June 2019 to March 2020, among 4069 urine cultures of patients aged 18 years and above of either gender. Urine samples were taken in a sterilized container and were cultured using standard techniques. Candida isolates were identified by Lactophenol blue wet films and Gram stain. SPSS-17 was used for statistical analysis. Descriptive statistics were calculated and chi square test was applied to see the associations.

Results: Among 4069 patients, 37.6% were male and 62.4% were female. Female patients revealed higher incidence of candida species in urine (7%) than that of male patients (3.2%) while there was no difference of Candida species isolations among ward and outdoor female (7.4% and 6.3%) and male patients (3.3% and 3%). The results shows significant association of urine culture result with gender, age and setting among patient <2 years.

Conclusion: Urinary tract infections caused by Candida species were often encountered in both out-patients and in-patients in this study. Females were more commonly affected than males.

Corresponding author:**Dr Misbah Arshad**

DHQ hospital Faisalabad.

QR code



Please cite this article in press Misbah Arshad *et al*, *Prevalence Of Candida Isolates From Urine Of Patients.*, *Indo Am. J. P. Sci.*, 2020; 07(12).

INTRODUCTION:

Worldwide, urinary tract infection is found most common problem caused by microbial invasion to different tissues of urinary tract. [1] Gender, diabetes, hypertension, genetic predisposition, behavioral factors, urologic structural abnormalities, immune-suppression, pregnancy, stone formation, nosocomial acquired infections and instrumentation are known as most predisposing factors of UTIs. [2-5] There has been a marked increase in opportunistic fungal infections involving the urinary tract, of which candida species are the most prevalent. [6]

Candida is the one of most common opportunistic fungal infection, responsible for 90% of vaginitis infectious.[7,8] Incidence of lower urinary tract infections caused by yeasts is fourfold more common in women than in men.[9-11] Multiple studies indicate that at least 10 to 15 percent of hospital acquired UTIs are caused by candida species.[12-17] All common Candida species are capable of causing urinary tract infections. Candida species in measurable quantities in the urine are found in 1% of clean voided specimens in healthy persons while account for 5% of all urine culture results in the general hospital setting and 10% of urine isolates in tertiary care facilities. [6]

Most UTIs due to candida occurs in hospitalized patients with indwelling bladder catheters. [13,18-19] Within the hospital setting, candida is especially common in intensive care units (ICUs) and may represent the most frequent UTIs encountered in adult surgical ICUs. [13,18,19]

A study from Spain reported 22% of critically ill patients, hospitalized for more than 7 days in an ICU developed candida. [19] While study from France reported mean incidence of candida as 27.4/1000 among ICU admissions. [15] 25,000 cases per year was estimated by shay and miller while approx one-third of hospitalized patients with urine cultures yielding candida were in the ICU where bladder catheter use was high.¹⁷ Overall percentage of nosocomial UTIs among catheterized individuals was as high as 37%, of whom 16.4% had infections due to candida species. [20]

Best to our knowledge there have been very few studies on candida isolates in the asian population in general, and in the Pakistani population in particular. The aim of the present study is to reveal the prevalence of candida isolates from urine of patients reporting to Hospital.

MATERIAL & METHOD:

A cross-sectional study was carried out in DHQ Hospital, Faisalabad during June 2019 to March 2020 among 4069 sample of patients aged 18 years and above of either gender reported and or admitted to hospital for urinary complaints and advised urine for culture. This study was conducted after approval of hospital ethical review committee. Written and verbal informed consent was taken from the patient. Urine samples were taken in a sterilized container and were cultured quantitatively on CYSTINE LACTOSE ELECTROLYTE DEFICIENT (CLED) and blood agar using standard techniques. *Candida* isolates were identified by Lactophenol blue wet films and Gram stain.

Data was analyzed by using statistical package for social sciences (SPSS) version 17. Mean and standard deviation were computed for quantitative variable and frequency and percentage were calculated for qualitative variables. Stratification was done and post stratification chi square test was applied to check association between variables. P value ≤ 0.05 were considered as significant for all analysis.

RESULTS:

Among 4069 patients, 1529 were male and 2540 were female. For 2540 female patients, 1477 urine samples were collected from ward and 1063 urine samples were collected from outdoor. For 1529 male patients 936 urine samples were collected from ward and 593 urine samples were collected from outdoor. Among total study cases, 2441 were aged < 2 years and 1628 were aged ≥ 2 years. Most of patients (59.3%) were from Wards. Positive urine culture were found for 226(5.6%) of patients as presented in Table-1.

Table-1: Descriptive statistics of study Population

	n(%)
Gender	
Male	1529(37.6)
Female	2540(62.4)
Age Group	
<2 years	2441(60)
≥2 years	1628(40)
Setting	
OPD	1656(40.7)
Ward	2413(59.3)
Urine Culture Result	
Positive	226(5.6)
Negative	3843(94.4)

Female patients revealed higher incidence of *Candida* species in urine (7%) than that of male patients (3.2%). There was no difference of *Candida* species isolations among ward and outdoor female (7.4% and 6.3%) and male patients (3.3% and 3%). The results also showed significant association of urine culture result with gender ($p=0.000$), age ($p=0.022$), and setting among patient with age <2 years ($p=0.008$). Detailed results are presented in Table-2.

Table-2: Association of Urine Culture Result with other factors

	n(%)		P-Value
	Positive	Negative	
Gender			
Male	49(3.2)	1480(96.8)	0.000
Female	177(7)	2363(93)	
Age Group			
<2 years	152(6.2)	2289(93.8)	0.022
≥2 years	74(4.5)	1554(95.5)	
Setting			
Admitted	141(5.8)	2272(94.2)	0.331
OPD	85(5.1)	1571(94.9)	
Setting For Male			
Admitted	31(3.3)	905(96.7)	0.765
OPD	18(3)	575(97)	
Setting For Female			
Admitted	110(7.4)	1367(92.6)	0.264
OPD	67(6.3)	996(93.7)	
Setting for <2 years			
Admitted	90(6.9)	1205(93.1)	0.008
OPD	51(4.5)	1095(95.5)	
Setting for ≥2 years			
Admitted	51(4.6)	1067(95.4)	0.077
OPD	34(6.7)	476(93.3)	

Chi Square Test was applied.
 $P \leq 0.05$, considered as significant.

DISCUSSION:

Urinary tract fungal infections caused by *Candida* species are becoming most common and important problem over the past decades.[2] Among urinary tract infections, *Candida* ranked as seventh among all nosocomial pathogens and fourth among nosocomial pathogens of. [22] *Candida* is often diagnosed on the basis of clinical features alone as well as with other conditions.[3,4] The rate of misdiagnosed vulvovaginal candidia on clinical presentation alone is high and can be wrongly subjected to treatment. [24]

In our study, total of 4069 urine samples from patients were received, out of which 5.6% urine samples were found positive for candida which in nearly same to study reported by Ali JA et al⁷ while is lower than the 13.27% reported for candidia by Fernandez-Limia et al. [5]

In current study, 53% of the patients were females while 43% were males and the female-to-male ratio was 1.66:1. The results of the present study also showed a higher prevalence of *Candida* in females 7% than in males 3.2% which is similar to results reported by other studies. [12] This high incidence in females may reflect vaginal candidia.

Study limitation:

The study has certain limitations, like small sample size. Other the cases were those which presented to a public sector hospital, thus cases from private sector were not included and also the sample cannot represent the entire population of the community overall.

CONCLUSION:

Urinary tract infections caused by *Candida* species were often encountered in both out-patients and in-patients in this study. Females were more commonly affected than males. Considering the changing pattern of the disease, it has become important to monitor infections of urinary tract as the management would depend on the isolates.

REFERENCES:

1. Ellepola AN, Khan ZU, Joseph B, Chandy R, Philip L. Prevalence of *Candida dubliniensis* among oral *Candida* isolates in patients attending the Kuwait University Dental Clinic. *Med Princ Pract.*2011;20(3):271-6.
2. Khan ZU, Ahmad S, Mokaddas E, Chandy R. Simplified sunflower (*Helianthus annuus*) seed agar for differentiation of *Candida dubliniensis* from *Candida albicans*. *Clin Microbiol Infect.* 2004 Jun;10(6):590-2.
3. Ahmad S, Mokaddas E, Al-Sweih N, Khan ZU.

Phenotypic and molecular characterization of *Candida dubliniensis* isolates from clinical specimens in Kuwait. *Med Princ Pract.* 2005;14(Suppl. 1):77-83.

4. Wang H, Wang Y, Chen J, Zhan Z, Li Y, Xu J. Oral yeast flora and its ITS sequence diversity among a large cohort of medical students in Hainan, China. *Mycopathologia.* 2007;164(2):65-72.
5. Thaweboon S, Thaweboon B, Srithavaj T, Choonharuangdej S. Oral colonization of *Candida* species in patients receiving radiotherapy in the head and neck area. *Quintessence Int.* 2008;39(2).
6. Sobel JD, Fisher JF, Kauffman CA, Newman CA. *Candida* urinary tract infections—epidemiology. *Clin Infect dis.* 2011;52(suppl_6):S433-6.
7. Alli JA, Okonko IO, Odu NN, Kolade AF, Nwanze JC. Detection and prevalence of *Candida* isolates among patients in Ibadan, Southwestern Nigeria. *World J Microbiol Biotechnol.* 2017;1(3):176-84.
8. Nwadioha S, Egesie JO, Emejuo H, Iheanacho E. Prevalence of pathogens of abnormal vaginal discharges in a Nigerian tertiary hospital. *Asian Pac J Trop Med.* 2010;3(6):483-5.
9. Behzadi P, Behzadi E. The microbial agents of urinary tract infections at central laboratory of Dr. Shariati Hospital, Tehran, Iran. *Turk Klin Tip Bilim.* 2008;28(4):445.
10. Behzadi P, Behzadi E, Yazdanbod H, Aghapour R, Cheshmeh MA, Omran DS. A survey on urinary tract infections associated with the three most common uropathogenic bacteria. *Maedica.* 2010;5(2):111.
11. Behzadi P, Behzadi E, Yazdanbod H, Aghapour R, Cheshmeh MA, Omran DS. Urinary tract infections associated with *Candida albicans*. *Maedica.* 2010;5(4):277.
12. Kauffman CA, Vazquez JA, Sobel JD, Gallis HA, McKinsey DS, Karchmer AW, et al., Prospective multicenter surveillance study of funguria in hospitalized patients. *Clin Infect Dis.* 2000;30(1):14-8.
13. Kauffman CA. Candiduria *Clin Infect Dis* 2005; 41:S371–6.
14. Kobayashi CC, Fernandes OD, Miranda KC, de Sousa ED, Silva MD. Candiduria in hospital patients: a study prospective. *Mycopathologia.* 2004;158(1):49-52.
15. Bougnoux ME, Kac G, Aegerter P, d'Enfert C, Fagon JY, CandiRea Study Group. Candidemia and candiduria in critically ill patients admitted to intensive care units in France: incidence, molecular diversity, management and outcome. *Intensive Care Med* 2008;34(2):292-9.