



CODEN [USA]: IAJPB

ISSN : 2349-7750

**INDO AMERICAN JOURNAL OF  
PHARMACEUTICAL SCIENCES**

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4338045>Available online at: <http://www.iajps.com>

Research Article

**PRACTICABILITY OF CONDENSED PORT LAPAROSCOPIC  
COLORECTOMY WITH NATURAL ORIFICE SPECIMEN  
REMOVAL SURGERY FOR COLORECTAL CANCER**<sup>1</sup>Dr Aniqa Rahman, <sup>2</sup>Dr. Tehreem Fatima, <sup>3</sup>Dr Maryam Mumtaz<sup>1</sup>DHQ Teaching Hospital Sahiwal**Article Received:** October 2020**Accepted:** November 2020**Published:** December 2020**Abstract:**

*The aim of this study is to review the practicability and benefits of reduced port laparoscopic surgery with NOSES. Over the past years rectal cancer surgery has been advanced with the addition of different techniques to reduce the invasiveness with better outcome. Minimal invasive surgery is the finest addition in the treatment with reduced post-operative pain, less wound complication, earlier return of bowel function, less adhesions, better cosmetic and possibly shorter hospital stay. In conventional technique of Specimen extraction, we need "mini-laparotomy" which needs a long cut on the abdomen for the extraction of specimen which mask the benefits of laparoscopic surgery. Natural orifice specimen extraction (NOSE) is a better technique to overcome this shortcoming in which we extract the specimen through the hollow viscus that already communicates with the outside world such as anus or vagina. The notion of this procedure is to remove the specimen with decrease trauma required for the removal of specimen with better outcome such as reduced post-operative pain, less use of analgesics, early healing with less wound complications and most important the excellent cosmesis effect compared to conventional specimen extraction. While the feasibility of NOSE has been demonstrated but there are also some limitation of NOSE which effect the application of this technique in colorectal surgery. Certain factor like BMI, ASA Score, Specimen size, and gender are main factors for effectiveness of this technique.*

**Keywords:** *Colorectal cancer surgery, natural orifice specimen extraction, rectum, feasibility of NOSE, colorectal, minimally invasive surgery.*

**Corresponding author:**

**Dr. Aniqa Rahman,**  
DHQ Teaching Hospital Sahiwal.

QR code



Please cite this article in press Aniqa Rahman et al, *Practicability Of Condensed Port Laparoscopic Colorectomy With Natural Orifice Specimen Removal Surgery For Colorectal Cancer., Indo Am. J. P. Sci, 2020; 07(12).*

**INTRODUCTION:****History and Feasibility**

The treatment strategy for colorectal cancer, especially low rectal cancer has been changed and improved dramatically in the last century. Colorectal Cancer is one of the major cause of death worldwide and the number of colorectal patients has significantly increased in the last few years. Abdominoperineal resection subsequently became the standard surgical treatment for rectal cancer which was described around 1900, however the attendant risk of operative mortality was very high[1]. Preservation of the anal sphincter is a major aspect of colorectal cancer surgery which still exist today and achieved successfully in mostly cases. Anterior resection and the pull-through technique were developed and progress has been made in these procedures. In 1980s, Laparoscopic surgery was introduced and total mesorectal excision (TME) advocated for the thorough removal of cancer [2,3]. Laparoscopic colectomy and proctectomy has become the standard technique for rectal cancer treatment now a days. Laparoscopic surgery is more beneficial and advantageous in term of enhanced visuals with high resolution, less postoperative pain, better quality of life with equivalent oncologic outcome. Clinical trials demonstrated the benefit of laparoscopic surgery over open surgery for rectal cancer [4,5,6].

The evolution of laparoscopic surgery has recently led to the introduction of Single Incision Laparoscopic Surgery (SILS) and Reduced Port Laparoscopic Surgery (RPLS), despite the limited laparoscopic handling space. The conventional multi-port laparoscopic colorectal surgeries (CMLS) are performed through multi-ports allowing variation of scope and other instruments placement for easy and safe dissection. Certain studies previously compared single-incision laparoscopic colectomy to conventional multi-port laparoscopic colectomy for CRC in terms of operative procedure and outcome. There were no differences in operative time, open conversion, number of harvested lymph nodes, length of hospital stay, postoperative complications and mortality but with certain technical difficulties and surgeon fatigue [7-9]. So a better technique is developed which combined the advantages of conventional laparoscopic surgery and SILS. This technique is known as Reduce Port Laparoscopic Surgery (RPLS) in which 3 or 4 ports are introduced in the body according to surgical technique (right hemicolectomy, left hemicolectomy or anterior resection). Additional ports may be added according to surgeon decision and operative difficulties. Results for RPLS are comparable to large clinical trials and results is not inferior to conventional laparoscopic

surgery. Several studies of RPLS reported that operation time of RPLS was comparable to CMLS [10,11]. This technique must therefore prove to be equal to conventional techniques with regard to benefits and oncologic safety. Laparoscopic surgery can meet social needs towards quality of life (QOL) in cancer treatment by enhancing patient satisfaction with reduced abdominal wall trauma and post-surgical scarring. So RPLS is better technique which alleviate the glitches of SILS and open surgery and become a bridge between conventional laparoscopic surgery and SILS which provide a mid-way to for performing surgery with ease. Specimen extraction is one of the basic and challenging process in colorectal surgery. A mini-laparotomy is needed in which extra incision is given on the abdomen for the removal of specimen extraction which compromise the benefits of laparoscopic surgery. The length of incision depend on the size of specimen and associated with increased wound infection, hernia, and postoperative pain. [12-16] Natural orifice specimen extraction (NOSE) eliminates this extra incision which is otherwise indispensable for specimen removal. NOSE is defined as the removal of specimen from the body via natural route which is already communicated with outside world i.e. vagina and gastrointestinal tract. NOSE remove the extra incision from the abdomen thus relieve the patient from comorbidities caused by mini-laparotomy. Feasibility of NOSE for colorectal surgery is acknowledged in literature. Stewart et al [17] and Nezhat [18] extract the colectomy specimen through the vagina reported in 1991 and 1992. In Nezhat's series, 9 patient were treated for endometriosis with transvaginal specimen extraction, however this technique is successfully used now for treatment of malignancy, diverticulitis and inflammatory bowel disease [19]. Successful colon and rectum cancer removal are described in various case reports and studies. Franklin et al [20] first describe the partial colectomy with NOSE via anus.

Intra corporeal anatomies are after the removal of distal site such as distal descending colon and sigmoid colon cancers is pre-requisite skill for the adoption of NOSES. NOSE technique is difficult due to steep learning curve so need more technical skills. Removal of more proximal specimen as in right hemicolectomy need an expert surgeon because of the length by which specimen passed and tortuous sigmoid colon made it difficult. Posterior colpotomy is needed in those patient in which specimen is removed by the vagina and this technique is normally not performed by colorectal surgeons. There is no standardization of NOSE technique which also intensified the technical difficulties.

Practically left sided colon pathology is more easy and feasible to perform than right side colon pathologies. Long length of ascending colon, transverse colon, descending colon and narrow tortuous sigmoid colon made it more difficult to pass the large size specimen of right colon cancer. So there is strict criteria of specimen size for the right side colon pathologies than left side colon lesions which limit its use for right sided colon cancer. The vagina has been established as a specimen extraction route in NOSE for gynecological and colorectal pathologies. First vaginal specimen extraction is noted in 1990s. This technique has been used mostly for right side colon pathologies or bulky size

specimen [21]. Vagina is better option because of its improved healing and elasticity but only possible in females [22-24]. A greater volume of literature focus on the extraction of sigmoid and rectal cancer via NOSE. Wound protection is needed to avoid from implantation of cancer cell and bacterial infection. To date, many studies are published for successful extraction of specimen via colon, rectum, anus and vagina to cure both benign and malignant tumor [19]. Many studies reports NOSE benefits over conventional specimen extraction in term of Post-operative analgesic use, time of first bowel function, cosmesis and length of hospital stay [25-27].

N o.	Study Group	Successful NOSES/ Total Attempts	Success Rate (%)	Sex (M/F)	Age Mean(Range) (Years)	BMI Mean(Range)
1	Wolthuis et al, <sup>41</sup> 2015	19/20	95	5/15	54 (median) (31–72)	23.5 (18–29)
2	Saurabh et al, <sup>25</sup> 2017	77/82	93.9	47/35	63.3±13.9	24.4±4.2
3	Karagul et al <sup>16</sup> ,2017	49/67	73.1	40/27	57.9±13.4	Unk
4	Wolthius et al, <sup>42</sup> 2015	17/17	100	3/15	Unk	Unk
5	Pai et al, <sup>43</sup> ,2015	19/19	100	13/6	48 (median) (range:23–78)	Unk
6	Wang et al, <sup>44</sup> , 2013	21/21	100	4/17	62 (median) (range:50–80)	23.6 (mean) (range:18–30)
7	Zhang et al, <sup>45</sup> , 2014	18/18	100	10/8	56.6 (mean) (range:48–69)	22.6 (mean) (range:19.7–26.4)
8	Hisada et al, <sup>46</sup> 2014	20/20	100	12/8	63.7±9	25.47±3.02
9	Cotantino et al, <sup>47</sup> ,2012	16/17	94.1	6/11	60.1±9.42	25.47±3.02
10	Zhang et al, <sup>48</sup> ,2014	24/27	88.9	16/11	54.8 (mean) (range:37–77)	22.11 (mean) (range:18.4–30.2)
11	Han et al, <sup>49</sup> ,2013	21/21	100	12/9	45.4±3.6	23.1±2.8
12	Wolthius et al, <sup>50</sup> , 2015	110/110	100	13/97	38 (median) (range:32–56)	23 (median) (range:21–25)
13	Xingmao et al, <sup>27</sup> 2014	65/65	100	32/33	56.1±9.3	23.7±2.9
14	Franklin et al, <sup>51</sup> ,2013	303/303	100	Unk	Transvaginal: 69.9±14.8 Transanal: 65.3±11.6	Unk
15	Akamatsu et al, <sup>52</sup> , 2009	16/16	100	Unk	Unk	Unk
16	Nishimura et al, <sup>53</sup> ,2011	17/18	94.4	14/4	65.5 (mean) (range:52–89)	21.3 (mean) (range:16.1–24.9)
17	Wolthuis et al, <sup>54</sup> , 2011	21/21	100	2/19	41 (median) (range:34–66)	23 (median) (range:22–26)
18	Huang et al, <sup>55</sup> ,	32/32	100	17/15	68±13 (range: 43–90)	23.3±2.2 (range:

	2016					18–27)
19	Leung et al, <sup>56</sup> , 2013	35/35	100	13/22	62 (median) (range:51–86)	Unk
20	Kim et al, <sup>57</sup> , 2013	57/58	95	0/58	62.8±9	23.5±2.9
21	Denost et al, <sup>58</sup> , 2015	122/122	94.1	70/52	63 (median) (range:20–90)	24.3 (median) (17.3–33.6)
22	Awad et al, <sup>59</sup> , 2014	19/20	95	0/20	66.9±8.9	25.1±6.65
23	Park et al, <sup>60</sup> , 2011	32/34	94.1	0/34	61.0±11.2	23.9±3.1
24	Wolthius et al, <sup>61</sup> , 2011	21/21	100	Unk	35 (median) (range:30–38)	23 (median) (22–25)
25	Marks et al, <sup>62</sup> , 2017	193/193	100	135/58	59 (mean)	27 (mean)

No.	Comorbidity	Viserectomy Site	Location	Specimen Size
1.	ASA I=5 ASA II=15	Transanal=20	Left/sigmoid colon	Unk
2	ASA I=59 ASA II/III=23	Transanal=82	Rectosigmoid=69 Prox. Rectum=13	Largest diameter 2.9±1.6
3	Unk	Transanal=37 Transvaginal=12	Colon:Right=17 Transverse=1 Total=4 Rectosigmoid=30	4.6±3.4 (mean)
4	ASA I=4 ASA II=11 ASA III=2	Transanal=17	Left colon=17	Length: Camera sleeve: 23 (median)(range:20–26) Bag: 21 (median)(range: 15–25)
5	Unk	Transanal=19	Rectum=19	Unk
6	Unk	Transanal=16 Transvaginal=5	Rectum=21	2.8 (mean) (range:1.8–6.0)
7	Unk	Transanal=18	Midrectum=12 Low rectum=6	4.2 (mean) (range:3.5–6.5)
8	Unk	Transanal=20	Upper rectum/ sigmoid=20	2.7±0.9
9	1.47±0.51	Transanal=17	Sigmoid=17	Unk
10	Unk	Transanal=27	Sigmoid=13 Rectum=14	Unk
11	Unk	Transanal=21	Rectum=21	4.6±1.7
12	ASA I=45 ASA II=56 ASA III=9	unk	Left colon/sigmoid/ rectum=110	Unk
13	ASA I=10 ASA II=50 ASA III=5	Transanal=65	Sigmoid=27 Rectum=38	2.9±1.5
14	Unk	Transanal=277 Transvaginal=26	Unk	Unk

15	Unk	Transanal=16	Sigmoid/ Rectosigmoid=16	Unk
16	Unk	Transanal=18	Sigmoid=18	1.84 (mean) (range:0–4.0)
17	Unk	Transanal=21	Sigmoid=21	Length of specimen: 20 (median) (range:13– 25)
18	ASA I=4 ASA II=20 ASA III=8	Transanal=32	Sigmoid=15 Rectosigmoid=9 Upper rectum=6 Mid rectum=2	3.3±1.8 (range:1.3–6.2)
19	Unk	Transanal=35	Left colon=35	2 (median) (range:2–4)
20	ASA I=20 ASA II=32 ASA III =6	Transvaginal=58	Sigmoid=21 Rectosigmoid=8 Rectum=29	3.4±1.8
21	Unk	Transanal=122	Low rectum=122	3.9 (median) (range:1–10)
22	ASA II=4 ASA III=15 ASA IV=1	Transvaginal=20	Right colon=20	4.735±3.61
23	ASA I=12 ASA II=18 ASA III=4	Transvaginal=34	Cecum=10 Ascending Colon=16 Proximal transverse colon=8	3.8±1.3
24	ASA I=13 ASA II=8	Transvaginal=21	Sigmoid=21	Length of specimen: 21 (median) (range:17–24)
25	ASA I=4 ASA II=92 ASA III=92 ASA IV=5	Transanal=193	Low rectum=193	Unk

(ASA American Society of Anesthesiologist Score; Unk Unknown)

**Table 1:** NOSE surgeries related to Criteria of patient selection

### MATERIAL AND METHODS:

Randomized and controlled clinical trials or cohort observational studies were considered for inclusion. Medline (PubMed) was systematically searched for 2009 to 2019 for laparoscopic colorectal surgery with natural orifice specimen extraction. First, the title was screened and then studies were selected by reading abstract only. Of these text only English text were acquired. Relevant data was extracted according to predefined parameters of age, gender, BMI, operative approach, location of tumor, specimen size in the excel database. Case reports and studies without any proper method were excluded.

### Potential benefits:

Within laparoscopic surgery Reduced Port Laparoscopic surgery (RPLS) is the better in term of accessible movement of hands and less vulnerable port site infection. There are certain benefits of laparoscopic surgery over conventional open surgery

like less blood loss, less need of blood transfusion, Reduced Surgical Trauma, less use of epidural anesthesia, lower rate of postoperative complications after laparoscopic surgery, WBC and CRP are lower after laparoscopic surgery, less hospital stay, more lymph-node are harvested with CRT in low rectal cancer, lower rate of positive circumferential resection margin (CRM) due to better and magnified visuals by laparoscope [4,28].

Additional ports have been recommended for the safe completion of SILS. More trocar sites means more vulnerable sites for infection and more scars on body. So by decreasing the ports we have possibly less infection sites with less scars on the abdomen. While Single Incision Laparoscopic Surgery (SILS) is also comparable with RPLS but there is no additional ports exist and maneuvering is greatly restricted by nearby instruments. Therefore SILS requires an experienced surgeon to overcome the technical difficulties of triangulation, instruments crowding

and pneumoperitoneum leakage [29-31]. SILS technique is good in term of cosmetic score but increase the technical difficulty and fatigue of surgeon [32-33].

NOSE is more feasible in female than male patients. Many clinical studies shows successful NOSE without any difficulty and complication. Specimen extraction is a key factor with masks the benefits of laparoscopic surgery when specimen is extracted from abdominal site by giving an incision on the body. To avoid any incision we have the better option of Natural Orifice Specimen Extraction (NOSE) by which specimen extracted without giving any extra incision. The common focus of this technique is to reduce the incisional complication of pain, infection and hernia [34-36]. In male patients anus is main site for specimen extraction which can limit its application but in female vagina is better option especially in old female patients because of its elasticity and early healing properties.

#### **Pitfalls:**

The transanal specimen extraction (TASE) procedure may not be possible in patients with bulky tumors, a thick mesentery, a narrow rectum, or anal stenosis [37]. Other shortcoming of NOSES is Contamination caused by trans-anal specimen extraction (TASE) which can be classified into two categories. **Bacterial contamination** by bacterial colonies present in colon and **Tumor cell contamination** such as peritoneal dissemination, port-site recurrence, and delivery-site metastasis. Intra-corporeal contamination by tumor cells, which may arise during NOSE, must be avoided. McKenzie et al. reported that the risk of tumor seeding during transvaginal delivery is no higher than that associated with transabdominal extraction providing proper oncologic principles are followed and specimen handling is performed using a specimen retrieval bag [38-40]. Although rectal washout is performed, there is a risk of remnant tumor cells remaining. NOSE technique limitation is T4 stage cancer, large size specimen may injure the rectal wall or extra stretching of anal sphincter. Transvaginal specimen extraction (TVSE) procedure may not be possible for patients with bulky tumors, previous pelvic surgery or radiation, or a narrow vagina [38]. But overall this technique is feasible and beneficial for the specimen extraction. Downsides are present in surgical procedures as same in the case of NOSES. There are some limitation which bounds the patient selection for using this technique. In which the most important is size and stage of the tumor.

#### **DISCUSSION:**

NOSES is comparable to conventional specimen extraction surgery in selected patient population. Short term retrospective studies and randomized control trials shows benefits of NOSE over conventional specimen extraction surgery with mini-laparotomy. But we still need more randomized trials to compare the potential benefits of two techniques including early return of bowel function, less post-operative analgesic use, possible shorter hospital stay and wound complication significantly. The favorable outcomes motivate the use of this technique with caution for strictly selected criteria for patients. Several factors such as sex, BMI, pathology, anatomic location, size of tumor (width and length), distance from anal verge and specimen extraction site determines the feasibility of NOSE.

Fat patients mean more visceral fat which may relate to bulk of mesentery which impedes the specimen removal from natural orifice. A few studies shows the role of NOSE in patient with BMI>30 is feasible especially related to wound complication but feasibility of NOSE is not well proven in such patients. Standardization of this technique help to improve the research on feasibility outcome for colorectal cancer surgeries. Usually laparoscopic surgery is not common practice in fat patients. One more population can take benefits from NOSE are elderly and ill patient due to advantages of less blood loss, less susceptible to infection, increased inflammatory markers and wound complication which result in early healing and fast recovery. In literature most of studies include patient with fewer comorbidities mostly ASA classes I and II. Further investigations are needed to explore the benefits in sick patients with more comorbidities. NOSE provide other means of specimen removal in case of complication such as first priority is by anal route if specimen is large or any other complication we can use vaginal route even if this is not possible we can extract specimen by conventional surgery. Standardizing NOSE is the main issue due to variation in technique and procedure followed by surgeons. Many studies have different techniques for anastomosis (single-stapled versus double- or triple-stapled anastomosis) and removal of specimen (with plastic bag or wound protector).

#### **CONCLUSION:**

Reduced port surgery with NOSE is better technique than conventional open surgery results better outcome without compromising oncological safety of

the patient. Feasibility and safety of NOSE technique is shown by many clinical studies in the literature. Full effectiveness of this technique depends upon the patient selection. Criteria for patient selection should be BMI<30, ASA ≤3, T2 tumor stage and less bulky specimen. This technique is more effective in female because of vaginal specimen extraction. Sigmoid Colon Cancer and Low rectal cancer patient are most benefited from this technique.

#### Weakness:

We review articles from last ten years and randomly select only those articles which have proper method and results. Our study has several weaknesses. This is not a systematic literature review and did not include all literature on internet.

#### REFERENCE:

1. Lim SW, Kim HJ, Kim CH, Huh JW, Kim YJ, Kim HR. Umbilical incision laparoscopic colectomy with one additional port for colorectal cancer. *Tech Coloproctol* 2013;17:193-199.
2. Huscher CG, Mingoli A, Sgarzini G, Mereu A, Binda B, Brachini G and Trombetta S: Standard laparoscopic versus single incision laparoscopic colectomy for cancer: Early results of a randomized prospective study. *Am J Surg* 204: 115-120, 2012.
3. Kim SJ, Ryu GO, Choi BJ, Kim JG, Lee KJ, Lee SC and Oh ST: The short term outcomes of conventional and single port laparoscopic surgery for colorectal cancer. *Ann Surg* 254: 933-940, 2011.
4. M P Dziwiatr, M Pisarska, M Kisielewski, et al. Is ERAS in laparoscopic surgery for colorectal cancer changing risk factors for delayed recovery? *Med Onco* 2016; 33:25.
5. Hendrik J Bonjer. Survival after laparoscopic surgery versus open surgery for colon cancer: long-term outcome of a randomized clinical trial. *Lancet Oncol* 2009; 10: 44–52.
6. Kang SB, Park JW, Jeong SY, et al. Open versus laparoscopic surgery for mid or low rectal cancer after neoadjuvant chemoradiotherapy (COREAN trial): short-term outcomes of an open-label randomised controlled trial. *Lancet Oncol*. 2010;11:637–645.
7. Van der Pas MH, Haglind E, Cuesta MA, et al. Laparoscopic versus open surgery for rectal cancer (COLOR II): short-term outcomes of a randomised, phase 3 trial. *Lancet Oncol*. 2013;14:210–218.
8. Jacobs M, Verdeja JC, Goldstein HS. Minimally invasive colon resection (laparoscopic colectomy). *Surg Laparosc Endosc*. 1991;1:144–150.
9. Heald RJ, Husband EM, Ryall RD. The mesorectum in rectal cancer surgery: the clue to pelvic recurrence? *Br J Surg*. 1982;69:613–616.
10. Miles WE. A method of performing abdomino-perineal excision for carcinoma of the rectum and of the terminal portion of the pelvic colon. *Lancet*. 1908;2:1812–1813.
11. Franklin ME Jr, Ramos R, Rosenthal D, Schuessler W. Laparoscopic colonic procedures. *World J Surg*. 1993;17(1):51–56.
12. Wolthuis AM, de Buck van Overstraeten A, D’Hoore A. Laparoscopic natural orifice specimen extraction-colectomy: a systematic review. *World J Gastroenterol*. 2014;20(36):12981–12992.
13. Stewert EA, Liau AS, Friedman AJ. Operative laparoscopy followed by colpotomy for resecting a colonic leiomyosarcoma: a case report. *J Reprod Med*. 1991;36(12):883–884.
14. Karagul S, Kayaalp C, Sumer F, et al. Success rate of natural orifice specimen extraction after laparoscopic colorectal resections. *Tech Coloproctol*. 2017;21(4):295–300.
15. Winslow ER, Fleshman JW, Birnbaum EH, Brunt LM. Wound complications of laparoscopic vs open colectomy. *Surg Endosc*. 2002;16(10):1420–1425.
16. Ihedioha U, Mackay G, Leung E, Molloy RG, O’Dwyer PJ. Laparoscopic colorectal resection does not reduce incisional hernia rates when compared with open colorectal resection. *Surg Endosc*. 2008;22(3):689–692.
17. Kaminski JP, Pai A, Ailabouni L, et al. Role of epidural and patient controlled analgesia in site-specific laparoscopic colorectal surgery. *JSLs*. 2014;18(4):e2014.00207.
18. Nezhat F. Laparoscopic segmental resection for infiltrating endometriosis of rectosigmoid colon: a preliminary report. *Surg Laparosc Endosc Percutan Tech*. 2001;11(1):67–68.
19. Hackert T, Uhl W, Buchler MW. Specimen retrieval in laparoscopic colon surgery. *Dig Surg*. 2002;19(6):502–506.
20. Kawamata F, Homma S, Minagawa N, Kawamura H, Takahashi N, Taketomi A. Comparison of single-incision plus one additional port laparoscopy-assisted anterior resection with conventional laparoscopy-assisted anterior resection for rectal cancer. *World J Surg* 2014;38:2716-23.
21. Monclova JL, Targarona EM, Vidal P, Peraza Y, Garcia F, Otero CR, Pallares L, Balague C, Trias M (2013) Single-incision versus reduced-port splenectomy: searching for the best alternative to conventional laparoscopic splenectomy. *Surg Endosc* 27:895–902

22. Ross S, Rosemurgy A, Albrink M, Choung E, Dapri G, Gallagher S, Hernandez J, Horgan S, Kelley W, Kia M, Marks J, Martinez J, Mintz Y, Oleynikov D, Pryor A, Rattner D, Rivas H, Roberts K, Rubach E, Schwaitzberg S, Swanstrom L, Sweeney J, Wilson E, Zemon H, Zundel N (2012) Consensus statement of the consortium for LESS cholecystectomy. *Surg Endosc* 26:2711–2716
23. Koya H, Ryosuke O, Yoshiharu S, Open versus Laparoscopic Surgery for Advanced Low Rectal Cancer. *Ann Surg.* 2017; DOI: 10.1097/SLA.0000000000002329.
24. Xingmao Z, Haitao Z, Jianwei L, Huirong H, Junjie H, Zhixiang Z. Totally laparoscopic resection with natural orifice specimen extraction (NOSE) has more advantages comparing with laparoscopic-assisted resection for selected patients with sigmoid colon or rectal cancer. *Int J Colorectal Dis.* 2014;29(9):1119–1124.
25. Ma B, Huang XZ, Gao P, et al. Laparoscopic resection with natural orifice specimen extraction versus conventional laparoscopy for colorectal disease: a meta-analysis. *Int J Colorectal Dis.* 2015;30(11):1479–1488.
26. Saurabh B, Chang SC, Ke TW, et al. Natural orifice specimen extraction with single stapling colorectal anastomosis for laparoscopic anterior resection: feasibility, outcomes, and technical considerations. *Dis Colon Rectum.* 2017;60(1):43–50.
27. McKenzie S, Baek JH, Wakabayashi M, Garcia-Aguilar J, Pigazzi A (2010) Totally laparoscopic right colectomy with transvaginal specimen extraction: the author's initial institutional experience. *Surg Endosc* 24:2048–2052
28. Franklin ME Jr, Kelley H, Kelley M, Brestan L, Portillo G, Torres J (2008) Transvaginal extraction of the specimen after total laparoscopic right hemicolectomy with intracorporeal anastomosis. *Surg Laparosc Endosc Percutan Tech* 18:294–298
29. Choi GS, Park IJ, Kang BM, Lim KH, Jun SH (2009) A novel approach of robotic-assisted anterior resection with transanal or transvaginal retrieval of the specimen for colorectal cancer. *Surg Endosc* 23:2831–2835
30. Karagul S, Kayaalp C, Sumer F, et al. Success rate of natural orifice specimen extraction after laparoscopic colorectal resections. *Tech Coloproctol.* 2017;21(4):295–300.
31. McKenzie S, Baek JH, Wakabayashi M, Garcia-Aguilar J, Pigazzi A (2010) Totally laparoscopic right colectomy with transvaginal specimen extraction: the author's initial institutional experience. *Surg Endosc* 24:2048–2052
32. Palanivelu C, Rangarajan M, Jategaonkar PA, Anand NV (2008) An innovative technique for colorectal specimen retrieval: a new era of “natural orifice specimen extraction” (NOSE). *Dis Colon Rectum* 51:1120–1124
33. Atsushi Nishimura, Mikako Kawahara, Keisuke Honda et al. Totally laparoscopic anterior resection with transvaginal assistance and transvaginal specimen extraction: a technique for natural orifice surgery combined with reduced-port surgery.
34. Atsushi Nishimura, Mikako Kawahara, Kazuyoshi Suda et al. Totally laparoscopic sigmoid colectomy with transanal specimen extraction. *Surg Endosc*, 2011; 25:3459–3463.
35. Choi GS, Park IJ, Kang BM, Lim KH, Jun SH (2009) A novel approach of robotic-assisted anterior resection with transanal or transvaginal retrieval of the specimen for colorectal cancer. *Surg Endosc* 23:2831–2835
36. McKenzie S, Baek JH, Wakabayashi M, Garcia-Aguilar J, Pigazzi A (2010) Totally laparoscopic right colectomy with transvaginal specimen extraction: the author's initial institutional experience. *Surg Endosc* 24:2048–2052
37. Palanivelu C, Rangarajan M, Jategaonkar PA, Anand NV (2008) An innovative technique for colorectal specimen retrieval: a new era of “natural orifice specimen extraction” (NOSE). *Dis Colon Rectum* 51:1120–1124
38. Yun JA, Kim HC, Park JS, Cho YB, Yun SH, Lee WY. Perioperative and oncologic outcomes of single-incision laparoscopy compared with conventional laparoscopy for colon cancer: an observational propensity score-matched study. *Am Surg* 2015;81:316–323.
39. Markar SR, Wiggins T, Penna M, Paraskeva P. Single-incision versus conventional multiport laparoscopic colorectal surgery systematic review and pooled analysis. *J Gastrointest Surg* 2014; 18:2214–27.
40. Makino T, Milsom JW, Lee SW (2012) Feasibility and safety of single-incision laparoscopic colectomy. *Ann Surg* 255:667–676
41. Wolthuis AM, de Buck van Overstraeten A, Fieuws S, Boon K, D'Hoore A. Standardized laparoscopic NOSE-colectomy is feasible with low morbidity. *Surg Endosc.* 2015;29(5):1167–1173.
42. Han FH, Hua LX, Zhao Z, Wu JH, Zhan WH. Transanal natural orifice specimen extraction for laparoscopic anterior resection in rectal cancer. *World J Gastroenterol.* 2013;19(43):7751–7757.
43. Zhang X, Zhou H, Hou H, Hu J, Wang H, Zhou Z. Totally laparoscopic resection with natural orifice specimen extraction for carcinoma of

- sigmoid colon and rectum: a feasible and innovative technique. *J Clin Gastroenterol.* 2014;48(7):e57–e61.
44. Costantino FA, Diana M, Wall J, Leroy J, Mutter D, Marescaux J. Prospective evaluation of peritoneal fluid contamination following transabdominal vs. transanal specimen extraction in laparoscopic left sided colorectal resections. *Surg Endosc.* 2012;26(6):1495–1500.
  45. Hisada M, Katsumata K, Ishizaki T, et al. Complete laparoscopic resection of the rectum using natural orifice specimen extraction. *World J Gastroenterol.* 2014;20(44):16707–16713.
  46. Zhang XM, Wang Z, Hou HR, Zhou ZX. A new technique of totally laparoscopic resection with natural orifice specimen extraction (NOSE) for large rectal adenoma. *Tech Coloproctol.* 2015;19(6):355–360.
  47. Wang Q, Wang C, Sun DH, Kharbuja P, Cao XY. Laparoscopic total mesorectal excision with natural orifice specimen extraction. *World J Gastroenterol.* 2013;19(5):750–754.
  48. Pai VD, Desouza A, De Menezes JL, Saklani AP. Laparoscopic intersphincteric resection and hand-sewn coloanal anastomosis: a natural orifice specimen extraction technique. *J Laparoendosc Adv Surg Tech A.* 2015;25(5):396–400.
  49. Wolthuis AM, De Buck Van Overstraeten A, D’Hoore A. Laparoscopic NOSE colectomy with a camera sleeve: a technique in evolution. *Colorectal Dis.* 2015;17(5):O123–O125.
  50. Wolthuis AM, Fieuws S, Van Den Bosch A, de Buck van Overstraeten A, D’Hoore A. Randomized clinical trial of laparoscopic colectomy with or without natural-orifice specimen extraction. *Br J Surg.* 2015;102(6):630–637.
  51. Marks JH, Myers EA, Zeger EL, Denittis AS, Gummadi M, Marks GJ. Long-term outcomes by a transanal approach to total mesorectal excision for rectal cancer. *Surg Endosc.* 2017;31(12):5248–5257.
  52. Wolthuis AM, Meuleman C, Tomassetti C, et al. Laparoscopic sigmoid resection with transrectal specimen extraction: a novel technique for the treatment of bowel endometriosis. *Human Reprod.* 2011;26(6):1348–1355.
  53. Park JS, Choi GS, Kim HJ, Park SY, Jun SH. Natural orifice specimen extraction versus conventional laparoscopically assisted right hemicolectomy. *Br J Surg.* 2011;98(5):710–715.
  54. Awad ZT, Griffin R. Laparoscopic right hemicolectomy: a comparison of natural orifice versus transabdominal specimen extraction. *Surg Endosc.* 2014;28(10):2871–2876.
  55. Kim HJ, Choi GS, Park JS, Park SY, Ryuk JP, Yoon SH. Transvaginal specimen extraction versus conventional minilaparotomy after laparoscopic anterior resection for colorectal cancer: mid-term results of a case-matched study. *Surg Endosc.* 2014;28(8):2342–2348.
  56. Leung AL, Cheung HY, Fok BK, Chung CC, Li MK, Tang CN. Prospective randomized trial of hybrid NOTES colectomy versus conventional laparoscopic colectomy for left-sided colonic tumors. *World J Surg.* 2013;37(11):2678–2682.
  57. Denost Q, Adam JP, Pontallier A, Celerier B, Laurent C, Rullier E. Laparoscopic total mesorectal excision with coloanal anastomosis for rectal cancer. *Ann Surg.* 2015;261(1):131–143.
  58. Huang CC, Chen YC, Huang CJ, Hsieh JS. Totally Laparoscopic colectomy with intracorporeal side-to-end colorectal anastomosis and transrectal specimen extraction for sigmoid and rectal cancers. *Ann Surg Oncol.* 2016;23(4):1164–1168.
  59. Wolthuis AM, Pennickx F, D’Hoore A. Laparoscopic sigmoid resection with transrectal specimen extraction has a good short term outcome. *Surg Endosc.* 2011;25(6):2034–2038.
  60. Nishimura A, Kawahara M, Suda K, Makino S, Kawachi Y, Nikkuni K. Totally laparoscopic sigmoid colectomy with transanal specimen extraction. *Surg Endosc.* 2011;25(10):3459–3463.
  61. Akamatsu H, Omoro T, Oyama T, Masayuki T, Ueshima S, Masaaki N. Totally laparoscopic sigmoid colectomy: a simple and safe technique for intracorporeal anastomosis. *Surg Endosc.* 2009;23(11): 2605–2609.
  62. Franklin ME Jr, Liang S, Russek K. Natural orifice specimen extraction in laparoscopic colorectal surgery: transanal and transvaginal approaches. *Tech Coloproctol.* 2013;17(Suppl 1):S63–S67.