



CODEN [USA]: IAJPBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4384515>Available online at: <http://www.iajps.com>

Research Article

STUDY TO COMPARE THE ANTIBIOTICS TREATMENTS VERSUS APPENDECTOMY PATIENTS WITH UNCOMPLICATED ACUTE APPENDICITIS

¹Dr Sara Imam, ²Dr Muhammad Sallah, ³Dr Frieha Raza

¹Fatima Jinnah Medical University Lahore

²Chandka Medical College [LUMHS] Jamshoro

³Avicenna Medical College Lahore

Article Received: October 2020

Accepted: November 2020

Published: December 2020

Abstract:

Objective: To comparison between antibiotic therapy and appendectomy in patients presented with uncomplicated acute appendicitis.

Study Design: Randomized controlled trial

Place and Duration of Study: This study was conducted at the Department of Surgery Jinnah Hospital Lahore from August 2019 to July 2020.

Materials and Methods: One hundred and thirty patients of both genders presented with uncomplicated acute appendicitis were included and divided all the patients equally into two groups A and B. Each group comprised 65 patients. Group A received appendectomy while group B received antibiotic treatment. Outcomes such as hospital stay, complications, recurrence within one year and patient's satisfaction were examined.

Results: There were 40 (61.54%) male and 25 (38.46%) female patients in group A with mean age 30.06 ± 8.72 years while in group B 36 (55.38%) male and 29 (44.62%) were females with mean age 29.42 ± 9.65 years. Significant difference was observed regarding hospital stay between group A and B (2.02 ± 0.85 days Vs 6.28 ± 2.44 days) p -value 0.001. Regarding complications group B had overall lower rate of complications as compared to group A [(2 (3.08%) Vs 10 (15.38%)] a significant difference was observed between both group ($p=0.02$). In group A 1 (1.54%) patient develop recurrence while in group B 16 (27.69%) developed recurrence within 1 year. In group A 63 (96.92%) patients were satisfied while in group B 44 (67.69%) patients were satisfied with treatment modality.

Conclusion: Surgical treatment is safe and effective treatment modality with higher patients satisfaction and successful treatment rate as compare to antibiotics treatment.

Key Words: Uncomplicated acute appendicitis, Antibiotic drugs, Appendectomy.

Corresponding author:**Dr Sara Imam,**

Fatima Jinnah Medical University Lahore

QR code



Please cite this article in press Sara Imam et al, Study To Compare The Antibiotics Treatments Versus Appendectomy Patients With Uncomplicated Acute Appendicitis., Indo Am. J. P. Sci, 2020; 07(12).

INTRODUCTION:

Appendicitis is the most well-known earnest condition by and large careful practice with a frequency ~100/100,000/year, and higher pervasiveness in men than ladies (8.6% versus 6.7%).^{1,2} Standard treatment is appendectomy (i.e., open and laparoscopic appendectomy). Around 310,000 appendectomies are performed/year in the United States, of which 250,000 have positive an infected appendix³, giving a negative appendectomy pace of about 15% to 30%.⁴

Appendectomy itself is related with intra and post-usable morbidities including vascular wounds, urinary tract inconveniences, hematomas, colonic fistulas, careful site diseases, bonds, inside deterrents, and huge length of medical clinic stay.^{5,6} The post-employable complexity rate ranges from 2% to 23% and over 3% of patients are readmitted with intestinal obstacle and post-usable grip.^{7,8}

Traditionalist treatment with anti-infection agents is an elective decision for an infected appendix; in spite of the fact that the danger of disappointment is about 13% higher, yet the danger of confusions is lower. Traditionalist treatment with anti-infection agents for simple a ruptured appendix is a topical issue by and large medical procedure and there are numerous investigations in kids, grown-ups and blended populaces (kids and grown-ups) individually.⁹⁻¹¹ Among 13 precise surveys in grown-ups, 10 audits considered just randomized controlled preliminaries (RCT) distributed during 1995 to 2015 with the quantity of included RCTs running from 3 to 6. Among them, all aside from one audit¹² pooled adequacy and confusions among anti-infection agents and appendectomy applying meta-examination. Albeit different anti-infection agents (for example third era of cephalosporin, metronidazole, penicillin, and beta-lactamase) had been utilized, they were crumpled into one classification when contrasted and appendectomy.¹⁰ We conducted present study with aimed to examine the effectiveness of antibiotic treatment and compare it with surgical treatment (appendectomy).

MATERIALS AND METHODS:

This randomized controlled trial was conducted at Department of surgery Jinnah Hospital Lahore from August 2019 to July 2020. During this period 130 patients of both gender presented with uncomplicated acute appendicitis were analyzed. Patients ages were ranging between 15 to 45 years. We divided all the patients equally into two groups A and B and each group consist of 65 patients. Patients demographics including age, sex, body mass index, and symptoms were recorded after taking informed consent. Patients with recurrence, complicated appendectomy and those with refusal of consent were excluded. Group A patients were received surgical treatment (open or laparoscopic appendectomy) while group B received antibiotic treatment. In antibiotic group 250 mg of ciprofloxacin and 500 mg of metronidazole were given three times a day for 5 days. Treatment outcomes were examined in term of hospital stay, complications such as wound infection, perforation, nausea vomiting, and abdominal abscess. Recurrence rate was examined within 1 year after treatment between both groups. All the data was analyzed by SPSS 24. Chi-square test was applied to compare the outcomes between both groups. P-value <0.05 was taken as statistically significant.

RESULTS:

There were 40 (61.54%) male and 25 (38.46%) female patients in group A with mean age 30.06 ± 8.72 years while in group B 36 (55.38%) male and 29 (44.62%) were females with mean age 29.42 ± 9.65 years. Mean BMI in group A was 26.42 ± 3.6 kg/m² and in group B it was 25.54 ± 4.7 kg/m². Pain in lower quadrant was found in all the patients of both groups 100%. Loss of appetite found in 18 (27.69%) in group A and 20 (30.77%) in group B, 15 (23.08%) in group A and 18 (27.69%) in group B patients had nausea and vomiting, fever found in 15 (23.08%) in group A and in group B it was found in 13 (20%) patients. We found no significant difference regarding age, gender, BMI and symptoms between both groups ($p > 0.05$) (Table 1). We found a significant difference regarding hospital stay between group A and B (2.02 ± 0.85 days Vs 6.28 ± 2.44 days) p-value 0.001.

Table No.1: Age, sex, BMI and symptoms between both groups

Variable	Group A	Group B	P-value
Age (Yrs)	30.06±8.72	29.42±9.65	>0.05
Gender			
Male	40 (61.54%)	36 (55.38%)	N/S
Female	25 (38.46%)	29 (44.62%)	
BMI (kg/m²)	26.42±3.6	25.54±4.7	N/S
Symptoms			
Pain in lower quadrant	65 (100%)	65 (100%)	>0.05
Loss of Appetite	18 (27.69%)	20 (30.77%)	
Nausea & Vomiting	15 (23.08%)	18 (27.69%)	
Fever	15 (23.08%)	13 (20%)	

Table No.2: Hospital stay and overall complications between both groups

Variable	Group A	Group B	P-value
Hospital stay (days)	2.02±0.85	6.28±2.44	0.001
Complications			
Yes	10 (15.38%)	2 (3.08%)	0.002
No	55 (84.62%)	63 (96.92%)	

Table No.3: Treatment success rate and patients satisfaction between both groups

Variable	Group A	Group B	P-value
Treatment successful			
Yes	64 (98.46%)	49 (75.38%)	0.001
No	1 (1.54%)	16 (27.69%)	
Patients satisfaction			
Yes	63 (96.92%)	44 (67.69%)	0.001
No	2 (3.08%)	31 (47.69%)	

Regarding complications group B had overall lower rate of complications as compared to group A [(2 (3.08%) Vs 10 (15.38%)] a significant difference was observed between both group (p=0.02). Regarding complications in group A 3 patients developed wound infection, 3 had perforation, 3 had nausea vomiting and 1 with fever and in group B 2 patients had abdominal abscess (Table 2). Regarding recurrence with 1 year after treatment, we found that in group A 1 (1.54%) patient develops recurrence while in group B 16 (27.69%) developed recurrence. A significant difference was observed between both groups A and B with p-value <0.001.

Overall successful treatment rate was high in group A as compared to group B 64 (98.46%) Vs 49 (75.38%), a significant difference was observed between both groups with p-value 0.024. According to patients satisfaction in group A 63 (96.92%) patients were satisfied while in group B 44 (67.69%) patients were satisfied with treatment modality (Table 3).

DISCUSSION:

Appendicitis is one of the most common painful disorders found all over the world and appendectomy is the common surgical intervention performed in surgical settings.¹³ Many of treatment modalities have been used for the treatment of appendicitis in which antibiotic therapy and surgical treatment have been widely used. However surgical treatment

considered as treatment of choice because of better efficacy, low recurrence and cost effectiveness.^{14,15} In present study 130 patients were analyzed and divided into two groups, one group received antibiotic therapy and one received surgical treatment (appendectomy). Overall male patients were high in numbers 76 (58.46%) as compared to females 41.54% with mean age 29.8±8.42 years.

These results showed similarity to many of previous study in which males were predominant 55% to 68% as compared to females and average age of patients were 25 years.^{16,17}

In our study we found that appendectomy treatment group had less hospital stay 2.02 ± 0.85 days (ranges 1 to 3 days) as compared to antibiotics treatment group 6.28 ± 2.44 days (ranges 3 to 10 days), a significant difference was observed between both groups with p-value < 0.001 . A meta-analysis conducted by Salminen et al¹⁸ reported that median hospital stay in antibiotic and appendectomy groups was 3 days. However patients treated with antibiotics had hospital stay ranges 3 to 20 days. Another study conducted by Ibrahim et al¹⁹ reported that surgical treatment group had longer hospital stay 3.6 ± 1.2 as compared to medical treatment group 2.2 ± 0.3 days with p-value < 0.05 .

In present study we found that overall complications rate was high in appendectomy group 15.38% as compared to antibiotics group 3.08%. Some of studies demonstrated that appendectomy had high complications rate 10 to 25% as compared to antibiotic treatment.^{19,20} However, some of studies reported no significant difference regarding complications between both treatment modalities.^{18,21} In this study regarding recurrence with 1 year after treatment, we found that in group A 1 (1.54%) patient develops recurrence while in group B 16 (27.69%) developed recurrence. A significant difference was observed between both groups A and B with p-value < 0.00 . These results were comparable to many of previous studies in which surgical treatment had less recurrence rate and high efficacy rate as compared to antibiotics groups.^{22,23} We found that overall successful treatment rate was high in group A as compared to group B 64 (98.46%) Vs 49 (75.38%), a significant difference was observed between both groups with p-value 0.024. According to patients satisfaction in group A 63 (96.92%) patients were satisfied while in group B 44 (67.69%) patients were satisfied with treatment modality. These results showed similarity to many of previous studies in which appendectomy resulted better efficacy with higher patients satisfaction rate as compared to antibiotic treatment.^{24,25}

CONCLUSION:

Surgical treatment is safe and effective treatment modality with higher patients satisfaction and successful treatment rate as compare to antibiotics treatment. However complications rate was high in appendectomy treated patients. We found no major complication in both groups.

REFERENCES:

1. Di Saverio S, Sibilio A, Giorgini E, et al. The NOTA Study (Non-Operative Treatment for Acute Appendicitis): prospective study on the efficacy and safety of antibiotics (amoxicillin and clavulanic acid) for treating patients with right lower quadrant abdominal pain and long-term follow-up of conservatively treated suspected appendicitis. *Ann Surg* 2014;260(1):109-117.
2. Huang L, Yin Y, Yang L, Wang C, Li Y, Zhou Z. Comparison of antibiotic therapy and appendectomy for acute uncomplicated appendicitis in children a metaanalysis. *JAMA Pediatrics* 2017;171(5):426-34.
3. Wang HT, Sax HC. Incidental appendectomy in the era of managed care and laparoscopy. *J Am Coll Surg* 2001;192:182-8.
4. Omundsen M, Dennett E. Delay to appendectomy and associated morbidity: a retrospective review. *ANZ J Surg* 2006;76(3):153-5.
5. Emil S, Duong S. Antibiotic therapy and interval appendectomy for perforated appendicitis in children: a selective approach. *Am Surg* 2007; 73(9):917-22.
6. Malik AA, Bari SU. Conservative management of acute appendicitis. *J Gastrointest Surg* 2009;13(5): 966-70.
7. Vons C, Barry C, Maitre S, et al. Amoxicillin plus clavulanic acid versus appendectomy for treatment of acute uncomplicated appendicitis: an openlabel, non-inferiority, randomised controlled trial. *Lancet* 2011;377(9777): 1573-9.
8. Styruud J, Eriksson S, Nilsson I, et al. Appendectomy versus antibiotic treatment in acute appendicitis. a prospective multicenter randomized controlled trial. *World J Surg* 2006;30:1033-7.
9. Ansaloni L, Catena F, Coccolini F, et al. Surgery versus conservative antibiotic treatment in acute appendicitis: a systematic review and meta-analysis of randomized controlled trials. *Dig Surg* 2011; 28(3):210-21.
10. Ehlers AP, Talan DA, Moran GJ, Flum DR, Davidson GH. Evidence for an antibiotics-first strategy for uncomplicated appendicitis in adults: a systematic review and gap analysis. *J Am Coll Surg* 2016;222(3):309-14.
11. Harnoss JC, Zelienska I, Probst P, et al. Antibiotics versus surgical therapy for uncomplicated appendicitis: systematic review and meta-analysis of controlled trials

- (PROSPERO 2015: CRD42015016882). *Ann Surg* 2017;265(5): 889-900.
12. Paajanen H, Gronroos JM, Rautio T, Nordstrom P, Aarnio M, Rantanen T, et al. A prospective randomized controlled multicenter trial comparing antibiotic therapy with appendectomy in the treatment of uncomplicated acute appendicitis (APPAC trial). *BMC Surg*. 2013;13:3.
 13. Kirkil C, Yigit MV, Aygen E. Long-term results of nonoperative treatment for uncomplicated acute appendicitis. *Turk J Gastroenterol* 2014;25(4): 393-7.
 14. Chang DC, Shiozawa A, Nguyen LL, et al. Cost of inpatient care and its association with hospital competition. *J Am Coll Surg* 2011;212:12-19.
 15. Dieleman JL, Baral R, Birger M, et al. US spending on personal health care and public health, 1996-2013. *JAMA* 2016;316:2627-46.
 16. Zaidi MA, Nazeer T, Abdul Aziz O, Ahmed A, Asad T, Dar ZS. Antibiotic Therapy versus appendectomy in uncomplicated acute appendicitis in terms of efficacy. *Pak Armed Forces Med J* 2017; 67(1):98-101.
 17. Malik AA, Bari S. Conservative management of acute appendicitis. *J Gastrointest Surg* 2009; 13: 966-70.
 18. Salminen P, et al. Five-year follow-up of antibiotic therapy for uncomplicated acute appendicitis in the APPAC randomized clinical trial". *JAMA* 2018; 320(12):1259-65.
 19. Mason RJ, Moazzez A, Sohn H, Katkhouda N. Meta-analysis of randomized trials comparing antibiotic therapy with appendectomy for acute uncomplicated (no abscess or phlegmon) appendicitis. *Surg Infect* 2012;13(2):74-84.
 20. Minneci PC, Sulkowski JP, Nacion KM, et al. Feasibility of a nonoperative management strategy for uncomplicated acute appendicitis in children. *J Am Coll Surg* 2014;219(2):272-9.
 21. Mahida JB, Lodwick DL, Nacion KM, et al. High failure rate of nonoperative management of acute appendicitis with an appendicolith in children. *J Pediatr Surg* 2016;51(6):908-11.
 22. Flum DR. Acute Appendicitis Appendectomy or the "Antibiotics First" Strategy. *N Engl J Med* 2015;372:1937-43.
 23. Varadhan KK, Humes DJ, Neal KR, Lobo DN. Antibiotic therapy versus appendectomy for acute appendicitis: a meta-analysis. *World J Surg* 2010;34(2):199-209.
 24. Sippola S, et al. Quality of life and patient satisfaction at 7-year follow-up of antibiotic therapy vs appendectomy for uncomplicated acute appendicitis: A secondary analysis of a randomized clinical trial. *JAMA Surg* 2020
 25. Talan DA, Saltzman DJ, Mower WR, et al, for the Olive View-UCLA Appendicitis Study Group. Antibiotics-first versus surgery for appendicitis: a US pilot randomized controlled trial allowing outpatient antibiotic management. *Ann Emerg Med* 2017; 70(1):1-11.