

CODEN [USA]: IAJPBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

http://doi.org/10.5281/zenodo.2552634

Available online at: <u>http://www.iajps.com</u>

Research Article

FREQUENCY OF GASTROESOPHGEAL REFLUX DISEASE SYMPTOMS AMONG ADULTS IN SAUDI ARABIA

Abdulrahman Nasser Alshahrani^{1*}, Ayman Obaid Alrashidi², Raneem Mishal Almandhour³, Mashael Belal Alrubaian³, Nouf Najem Aldafeery³

¹ Albashaer General Hospital, Saudi Arabia, ² King Abdulaziz University, Jeddah, Saudi Arabia, ³Almaarefa University, Riyadh, Saudi Arabia.

Abstract:

Background: Gastroesophageal reflux disease (GERD) is the most common upper gastrointestinal disorder encountered in the elderly patient. It is highly prevalent worldwide with a prevalence of 10-20% in the western world. The health-related quality of life (HRQL) is lower in individuals with GERD than in the general population and is comparable to that in individuals with other chronic diseases. It has a considerable impact on the quality of the patient's life through its symptoms and economically by following consultation procedures and medical care. A few studies have been done in Saudi Arabia using general population as subject and have reported a very high prevalence.

Objective: (1) Estimation of gastro GERD prevalence among schoolteachers in Saudi Arabia. (2) To assess the impact of GERD symptoms on teacher's daily life activity. (3) To compare prevalence and risk factors of GERD between age groups and gender. **Methodology:** A cross-sectional study that was conducted among 200 schoolteachers selected by multi-stage stratified random sample method in Saudi Arabia during 2018. A reliable and valid self-administered GERD questionnaire for diagnosis of GERD was used. GERD-HRQL questionnaire was used to assess the impact of GERD on the patient's life quality. Data were analyzed using Statistical Package for Social Sciences Version 20.0; Chi-square was used to test the association between GERD and sociodemographic data.

Results: The total number of the participants was 200 with an equal male to female ratio. 55% (116/200) of the participants reported with GERD. 53 point 5% of these (62/116) were female and 46.6% (54/116) were male. The commonest age group was 31-40 years with 45.5% (91/200) participants. 13/200 (6.5%) participants were smokers, of which only 15.9% were female. 41 point 3% (48/116) of the GERD +ve participants were having blood group O +ve 7.8% (9/116) of GERD participants reported symptoms which affected their daily life activity.

Conclusion: This study revealed a prevalence of GERD symptoms among 58% of schoolteachers. 7 point 8% of GERD participants reported symptoms, which affected their daily life activities. These data indicate a need for a comprehensive approach to managing the GERD and related diseases and a more intensified level of awareness about GERD symptoms and its complications. In addition, a health care and preventive measure may be implemented to tackle the problem among schoolteachers.

Keywords: Gastroesophageal, reflux, GERD, prevalence, teacher.

Corresponding author:

Abdulrahman Nasser Alshahrani, Albashaer General Hospital, Saudi Arabia. *E-mail: 123alfalleh@gmail.com*



Please cite this article in press Abdulrahman Nasser Alshahrani et al., **Frequency Of Gastroesophgeal Reflux** Disease Symptoms Among Adults In Saudi Arabia., Indo Am. J. P. Sci, 2019; 06(01).

INTRODUCTION:

Gastroesophageal reflux disease (GERD) is the most common upper gastrointestinal disorder encountered in patients. It is highly prevalent worldwide with a prevalence of 10-20% in the western world [1-4]. GERD is a general term for many symptoms that range from intermittent heartburn or acid regurgitation to end by most serious complication as Barrett's esophagus [5] and stricture [6]. The experience of GERD symptoms, at least, 2 times weekly is considered sufficient to cause an impaired quality of life [7]. Between adult patients with GERD who get medical care, up to 20% have serious complications [8].

Health-related quality of life (HRQL) is lower in individuals with GERD than in the general population and is comparable to that in individuals with other chronic diseases, such as diabetes, arthritis or chronic heart failure [9]. It usually has a considerable impact on the quality of the patient's life not only by the symptoms but also economically by following consultation procedures and medical care [1,10]. During the recent decade, several Asian studies about the prevalence of symptom-based GERD and endoscopic reflux esophagitis have revealed a higher number of patients compared to the previous studies [11]. A few studies have been done in Saudi Arabia using general population as subject and have reported a very high prevalence [12]. This study aimed to determine the prevalence and impact of GERD on the educated people and has used school teachers as the representative population.

METHODOLOGY:

A cross-sectional study conducted in Saudi Arabia. The study population included elementary, middle and high school teachers from the schools located in the main cities. Counting up of the sample size was depend on the assumption that the prevalence of GERD in Saudi Arabia will range between 37.5% and 52.5% based on the result reported by some earlier studies.12-14 The estimated sample size was 169. Sample size increased to 200 keeping in view the non-responders. A maximum period of 12-month for their call of symptoms was considered to reduce the chance of recall bias [15].

A multistage stratified random sample method was used. The sample was first stratified into different educational stages. The second stratification was for male and female schools with a ratio of 1:1, while the third stratification was for selecting teachers from selected schools with a male to female ratio of 1:1. These steps were done using randomizer research software.

We used the GERD questionnaire (GERDQ) for making the diagnosis of GERD [16]. The GERDQ

was developed as a patient-centered, self-assessment questionnaire to assist health-care professionals in the diagnosis of GERD. It has a sensitivity of 65% and a specificity of 71% [16]. Those with a score of ≥ 8 have a diagnosis of having GERD, while those with <8 don't have GERD [16]. To assess the impact of GERD on the patient's quality of life, we used GERD-HRQL questionnaire [17], information about age, gender, smoking habit, and blood grouping were also collected from each participant. Teachers were asked about any history and frequency of heartburn, epigastric pain, regurgitation of food, sleep interference from GERD symptoms, and the use of over the counter antacids for the control of their symptoms. They were also asked about the effect of the GERD symptoms on their social interaction.

- Group A 21-30 year
- Group B 31-40 year
- Group C 41-50 year
- Group D 51-60 year.

Statistical analysis

Statistical Package for Social Sciences Version 20.0 was used to analyze the data Pearson's Chi-square (χ^2) test was used to observe and quantify an association between the categorical outcome and the different variables. All calculated *P* values were two-tailed, with *P* < 0.05 considered as statistically significant.

RESULT:

The total numbers of the participants were 200 with an equal male to female ratio. According to age groups, 45.5% (n = 91) of participants fell in Group B, 28.5% (n = 57) in Group A, 24.5% (n = 44) in Group C and 1.5% (n = 3) in Group D. 6.5% (13/200) participants were smokers, of which 15.9% (2/13) were female 58% (116/200) of the participants reported with GERD. Among those diagnosed as GERD patients, 53.4% (62/116) were female and 46.6% (54/116) were male. GERD was the most common among blood group AB+ patients as 70.5% of them reported such symptoms.

GERD association with age

Around 47% (27/57) participants from Group A were found to have GERD. 51.8% (14/27) of them were male and 48.2% (13/27) were female. 65.9% (60/91) participants from Group B were found to have GERD. 36.6% (22/60) were males and 63.3% (38/60) were females. 59% (29/49) participants from Group C were found to have GERD. 62% (18/29) of them were males and 37.9% (11/29) were female. None of the participants from Group D reported GERD. 7.8% (9/116) of GERD participants reported symptoms which affected their daily life activities. Three of them were from Group A, two from Group B and four were from Group C. All of them from Groups A and B were female while in Group C three were female and one male (Table 1).

GERD association with smoking

Nearly 11.2% (13/116) GERD participants were smokers. 15.4% (2/13) of them were females and aged from 31- 40 years and 11 of them were males (1, 3, 6) from Groups A, B, and C, respectively. Daily life activities were affected in 7.7% (1/13) of smokers who were +ve with GERD. 4.3% (8/187) who reported the GERD symptoms affected their daily life activities were non-smokers. However, the relationship between smoking and GERD is significant statistically as the *P* value was 0.003 (Table 2).

GERD association with the educational level

About 67.6% (46/68) participants from elementary school level were found to have GERD. 50% of them were males have GERD. 35.2% (12/34) were males and 22 (64.7%) were female, but the relationship

between educational level and GERD is not statistically significant as the P value was 0.071 (Table 3).

About 11.11% (1/9) participants who reported with GERD symptoms which affected their daily life activities were from elementary school (one female), 11.11% (1/9) from middle school (one female), and 77.77% (7/9) were from high school (one male+six female).

GERD association with blood group

The frequency of GERD among the blood groups can be seen from Table 4. GERD was the most common among AB+ patients as 70.5% of them reported such symptoms. 11.11% (1/9) participants who reported with GERD symptoms which affected their daily life activities were one from blood group A+ (one female), 22.22% (2/9) from blood group B+ (two female), 33.33% (3/9) were from blood group AB+ (three female) and 33.33% (3/9) were from blood group O+ (one male + two female). However, the relationship between blood group and GERD is not statistically significant as the *P* value was 0.424.

Age groups	Number of participants	GERD present (%)	GERD pres ge	ent divided by nder	GERD affect their life		
			Male (%)	Female (%)	Male	Female	
A	The total=57 were 30 male and 27 female	47 (27/57)	51.8	48.2	Nil	3	
В	The total=91 were 38 male and 53 female	65.9 (60/91)	36.6	63.4	Nil	2	
С	The total=49 were 29 and male and 20 female	59 (29/49)	62	38	1	3	
D	The total=3 were 3 male	0	0	0	Nil	Nil	

Table 1: Age and gender distribution among study participants

GERD: Gastroesophageal reflux disease

Grandinaria	A ~~ ~~~~~	Number of	CEDD	CEDD	and disided	CEDD offeet	
Smokers	Age group	Number of	GEKD	GERD pro	esent aivided	GER	D affect
		participants	present (%)	by gender		their life	
				Male (%)	Female (%)	Male	Female
No	A=56	The	55.6	42.3	57.6	Nil	8
		total=187	(104/187)				
		were 89					
		male and 98					
		female					
	B=85						
	C=43						
	D=3						
Yes	A=1	The total=13 were 11 male and 2 female	92 (12/13)	83.3	16.6	1	Nil
	B=6						
	C=6						
	D=0						

Table 2:	Association	of smoking	with age and	gender distribution	among study participants
		· · · · · ·		G	

GERD: Gastroesophageal reflux disease

Table 3: Educational level with age and gender distribution among study participants

School levels	Age group	Number of participants	GERD present (%)	GERD resent (%)GERD present divided by genderGERD affect the		fect their life	
				Male (%)	Female (%)	Male	Female
Elementary school	A=27	The total=68 were 34 male and 34 female	67.6 (46/68)	50	50	Nil	1
	B=29						
	C=12						
	D=0						
Middle school	A=18	The total=66 were 33 male and 33 female	54.5 (36/66)	52.7	47.2	Nil	1
	B=33						
	C=14						
	D=2						
High school	A=12	The total=66 were 33 male and 33 female	51.5 (34/66)	35.2	64.7	1	6
	B=29						
	C=23						
	D=1						

GERD: Gastroesophageal reflux disease

Blood group	Age group	Number of participants	GERD present (%)	GERD present divided by gender		GERD affect their life	
				Male (%)	Female (%)	Male	Female
A ⁺	A=17	The total=50 were 27 male and 23 female	60 (30/50)	30	30	Nil	1
	B=26						
	C=7						
	D=0						
A	A=0	The total=4 were 1 male and 3 female	25 (1/4)	0	25	Nil	
	B=2						
	C=2						
	D=0						
B+	A=4	The total=28 were 16 male and 12 female	64 (18/28)	32.1	32.1	Nil	2
	B=13						
	C=11						
	D=0						
B-	A=0	The total=3 were 2 male and 1 female	66.6 (2/3)	33.3	33.3	Nil	
	B=1						
	C=2						
	D=0						
AB ⁺	A=4	The total=17 were 7 male and 10 female	70.5 (12/17)	23.5	47	Nil	3
	B=9						
	C=4						
	D=0						
AB ⁻	A=1	The total=2 were 2 male	50 (1/2)	50	0	Nil	
	B=0						
	C=0						
	D=1						
0+	A=29	The total=86 were 40 male and 46 female	55.8 (48/86)	23.2	32.5	1	2
	B=35						
	C=20						
	D=2						

Table 4: Blood group association with age and gender distribution among study participants

0-	A=2	The total=10	40 (4/10)	40	0	Nil	
		were 5 male					
		and 5 female					
	B=5						
	C=3						
	D=0						

GERD: Gastroesophageal reflux disease

DISCUSSION:

Our study found a higher prevalence of GERD among the Saudi school teachers as compared to the results reported by Almadi et al., who have reported a prevalence of 45.4% among the general population in Saudi Arabia [12]. A similar study done in Nigerian public elementary school teachers by Akande and Fadupin have reported a prevalence of 13.4% [18]. Fluctuating prevalence has been reported by various researchers across the world in general population including, 31.6% in Spanish [19]. 40.0% in Swedish, 25.7% in Iran, and 12% in Taiwan. Many factors have shown an association with GERD but still controversial. Male gender, hiatus hernia, and chronic obstructive pulmonary disease are three independent risk factors for the development of reflux esophagitis [20-22]. Nearly all epidemiologic studies have found a relationship between increasing body mass index due to obesity and changes in gastroesophageal anatomy and physiology. These include an increased prevalence of diminished pressure of lower esophageal sphincter, esophageal motor disorders, intragastric pressure, and the development of a hiatal hernia. Central obesity may be considered the most important risk factor for the development of reflux and related complications [23,24]. This study finds a higher prevalence of GERD among study population with age range 31-40 year though statistically nonsignificant. The association between GERD and age is controversial. Some studies have reported a direct association [25-27] while others an inverse [28-30] and still others no association whatsoever [31-35]. Our study has shown a higher prevalence of GERD in females. The effect of GERD on females was most severe with the fact that eight of nine teachers who reported the GERD symptoms affected their daily life activities were females.

Our study has shown a significant correlation (P = 0.003) of smoking with 6.5% of our study participants were smokers, and 92% (12/13) of them reported GERD. 56% (104/187) of non-smoking participants were also having GERD in our study. Several studies have revealed a direct relationship between GERD and smoking [36-37]. However,

others did not find any significant relationships [19,38]. Our study also finds an inverse relationship between educational levels with GERD. Elementary school teachers have shown highest prevalence as compared to high school teachers. A similar finding has been reported by Diaz-Rubio et al. and El-Serage et al. [19,39] The impact of symptoms on patients' daily life is one of the most common reasons for consultation for GERD [40]. Studies conducted among Swedish general population measure the impact of the severity and frequency of GERD symptoms on quality of life have found that even symptoms rated as mild are associated with a clinically meaningful reduction in well- being [41,42]. The highest numbers of our study participants were having blood group O +ve while the highest prevalence of GERD was in participants with blood group AB +ve. Type O has shown an association with naturally high stomach acid production and is much more likely to develop GERD, But when type A contracts GERD, it is more likely to develop Barrett's esophagus, and even esophageal cancer suggesting a genetic susceptibility [43].

CONCLUSION:

This study showed a high prevalence of GERD symptoms among school teachers which has an impact on their daily life activities. These data indicate a need for acomprehensive approach to GERD management in the health-care system. In addition, a health-care program may be implemented to address the problem among the teachers.

REFERENCES:

- 1. El-Serag HB. Time trends of gastroesophageal reflux disease: A systematic review. Clin Gastroenterol Hepatol 2007;5:17-26.
- Fujiwara Y, Higuchi K, Watanabe Y, Shiba M, Watanabe T, Tominaga K, *et al.* Prevalence of gastroesophageal reflux disease and gastroesophageal reflux disease symptoms in Japan. J Gastroenterol Hepatol 2005;20:26-9.
- 3. Shaheen NJ, Hansen RA, Morgan DR,

Gangarosa LM, Ringel Y, Thiny MT, *et al.* The burden of gastrointestinal and liver diseases, 2006. Am J Gastroenterol 2006;101:2128-38.

- Wong WM, Lai KC, Lam KF, Hui WM, Hu WH, Lam CL, *et al.* Prevalence, clinical spectrum and health care utilization of gastrooesophageal reflux disease in a Chinese population: A population- based study. Aliment Pharmacol Ther 2003;18:595-604.
- Vakil N, van Zanten SV, Kahrilas P, Dent J, Jones R; Global Consensus Group. The Montreal definition and classification of gastroesophageal reflux disease: A global evidence-based consensus. Am J Gastroenterol 2006;101:1900-20.
- 6. McDougall NI, Johnston BT, Kee F, Collins JS, McFarland RJ, Love AH. Natural history of reflux oesophagitis: A 10 year follow up of its effect on patient symptomatology and quality of life. Gut 1996;38:481-6.
- Dent J, Brun J, Fendrick A, Fennerty MB, Janssens J, Kahrilas P, *et al*. An evidence-based appraisal of reflux disease management the Genval workshop report. Gut 1998;44 Suppl 2:S1-16.
- 8. Chait M. Gastroesophageal reflux disease in the elderly. Pract Gastroenterol 2005;29:52-60.
- 9. Wiklund I. Review of the quality of life and burden of illness in gastroesophageal reflux disease. Dig Dis 2004;22:108-14.
- Chen MJ, Wu MS, Lin JT, Chang KY, Chiu HM, Liao WC, *et al.* Gastroesophageal reflux disease and sleep quality in a Chinese population. J Formos Med Assoc 2009;108:53-60.
- 11. Fock KM, Talley NJ, Fass R, Goh KL, Katelaris P, Hunt R, *et al.* Asia-Pacific consensus on the management of gastroesophageal reflux disease: Update. J Gastroenterol Hepatol 2008;23:8-22.
- Almadi MA, Almousa MA, Althwainy AF, Altamimi AM, Alamoudi HO, Alshamrani HS, *et al.* Prevalence of symptoms of gastroesopahgeal reflux in a cohort of Saudi Arabians: A study of 1265 subjects. Saudi J Gastroenterol 2014;20:248-54.
- 13. El-Serag HB, Sweet S, Winchester CC, Dent J. Update on the epidemiology of gastrooesophageal reflux disease: A systematic review. Gut 2014;63:871-80.
- 14. Al-Humayed SM, Mohamed-Elbagir AK, Al-Wabel AA, Argobi YA. The changing pattern of upper gastro-intestinal lesions in southern Saudi

Arabia: An endoscopic study. Saudi J Gastroenterol 2010;16:35-7.

- Dent J, El-Serag HB, Wallander MA, Johansson S. Epidemiology of gastrooesophageal reflux disease: A systematic review. Gut 2005;54:710-7.
- 16. Jones R, Junghard O, Dent J, Vakil N, Halling K, Wernersson B, *et al.* Development of the GerdQ, a tool for the diagnosis and management of gastro-oesophageal reflux disease in primary care. Aliment Pharmacol Ther 2009;30:1030-8.
- Wiklund IK, Junghard O, Grace E, Talley NJ, Kamm M, Veldhuyzen VZ, *et al.* Quality of life in reflux and dyspepsia patients. Psychometric documentation of a new diseasespecific questionnaire (QOLRAD). Eur J Surg Suppl Acta Chir Suppl 1997;583:41-9.
- 18. Akande K, Fadupin G. The prevalence of gastro-oesophageal reflux disease and its association with selected local food and beverages items among public schools teachers in a Nigerian city. Trial 2003;18:146-52.
- Diaz-Rubio M, Moreno-Elola-Olaso C, Rey E, Locke GR 3rd, Rodriguez-Artalejo F. Symptoms of gastro-oesophageal reflux: Prevalence, severity, duration and associated factors in a Spanish population. Aliment Pharmacol Ther 2004;19:95-105.
- 20. Chen TS, Chang FY. The prevalence and risk factors of reflux esophagitis among adult Chinese population in Taiwan. J Clin Gastroenterol 2007;41:819-22.
- 21. Ronkainen J, Aro P, Storskrubb T, Johansson SE, Lind T, Bolling- Sternevald E, *et al.* High prevalence of gastroesophageal reflux symptoms and esophagitis with or without symptoms in the general adult Swedish population: A Kalixanda study report. Scand J Gastroenterol 2005;40:275-85.
- 22. Vossoughinia H, Salari M, Mokhtari Amirmajdi E, Saadatnia H, Abedini S, Shariati A, *et al.* An epidemiological study of gastroesophageal reflux disease and related risk factors in urban population of mashhad, iran. Iran Red Crescent Med J 2014;16:e15832.
- 23. De Marco S, Passaglia C. Obesity and gastroesophageal reflux disease. Recenti Prog Med 2010;101:106-11.
- 24. Friedenberg FK, Xanthopoulos M, Foster GD, Richter JE. The association between gastroesophageal reflux disease and obesity. Am J Gastroenterol 2008;103:2111-22.
- 25. Bolin TD, Korman MG, Hansky J, Stanton R.

Heartburn: Community perceptions. J Gastroenterol Hepatol 2000;15:35-9.

- Isolauri J, Laippala P. Prevalence of symptoms suggestive of gastro-oesophageal reflux disease in an adult population. Ann Med 1995;27:67-70.
- Stanghellini V. Three-month prevalence rates of gastrointestinal symptoms and the influence of demographic factors: Results from the domestic/international gastroenterology surveillance study (DIGEST). Scand J Gastroenterol Suppl 1999;231:20-8.
- Drossman DA, Li Z, Andruzzi E, Temple RD, Talley NJ, Thompson WG, *et al.* US householder survey of functional gastrointestinal disorders. Dig Dis Sci 1993;38:1569-80.
- 29. Farup C, Kleinman L, Sloan S, Ganoczy D, Chee E, Lee C, *et al.* The impact of nocturnal symptoms associated with gastroesophageal reflux disease on health-related quality of life. Arch Intern Med 2001;161:45-52.
- Locke GR 3rd, Talley NJ, Fett SL, Zinsmeister AR, Melton LJ 3rd. Prevalence and clinical spectrum of gastroesophageal reflux: A population-based study in Olmsted County, Minnesota. Gastroenterology 1997;112:1448-56.
- Haque M, Wyeth JW, Stace NH, Talley NJ, Green R. Prevalence, severity and associated features of gastro-oesophageal reflux and dyspepsia: A population-based study. N Z Med J 2000;113:178-81.
- 32. Ho KY, Kang JY, Seow A. Prevalence of gastrointestinal symptoms in a multiracial Asian population, with particular reference to reflux-type symptoms. Am J Gastroenterol 1998;93:1816-22.
- 33. Kennedy T, Jones R. The prevalence of gastrooesophageal reflux symptoms in a UK population and the consultation behaviour of patients with these symptoms. Aliment Pharmacol Ther 2000;14:1589-94.
- 34. Louis E, DeLooze D, Deprez P, Hiele M, Urbain D, Pelckmans P, *et al.* Heartburn in Belgium: Prevalence, impact on daily life, and utilization of medical resources. Eur J Gastroenterol Hepatol 2002;14:279-84.
- 35. Ruth M, Månsson I, Sandberg N. The prevalence of symptoms suggestive of esophageal disorders. Scand J Gastroenterol 1991;26:73-81.
- 36. Nilsson M, Johnsen R, Ye W, Hveem K,

Lagergren J. Lifestyle related risk factors in the aetiology of gastro-oesophageal reflux. Gut 2004;53:1730-5.

- 37. Nocon M, Labenz J, Willich SN. Lifestyle factors and symptoms of gastro-oesophageal reflux -- A population-based study. Aliment Pharmacol Ther 2006;23:169-74.
- 38. Sharma PK, Ahuja V, Madan K, Gupta S, Raizada A, Sharma MP. Prevalence, severity, and risk factors of symptomatic gastroesophageal reflux disease among employees of a large hospital in Northern India. Indian J Gastroenterol 2011;30:128-34.
- 39. El-Serag HB, Petersen NJ, Carter J, Graham DY, Richardson P, Genta RM, *et al.* Gastroesophageal reflux among different racial groups in the United States. Gastroenterology 2004;126:1692-9.
- 40. Jones R, Armstrong D, Malfertheiner P, Ducrotté P. Does the treatment of gastroesophageal reflux disease (GERD) meet patients' needs? A survey-based study. Curr Med Res Opin 2006;22:657-62.
- Ronkainen J, Aro P, Storskrubb T, Lind T, Bolling-Sternevald E, Junghard O, *et al.* Gastro-oesophageal reflux symptoms and health- related quality of life in the adult general population - The Kalixanda study. Aliment Pharmacol Ther 2006;23:1725-33.
- 42. Wiklund I, Carlsson J, Vakil N. Gastroesophageal reflux symptoms and wellbeing in a random sample of the general population of a Swedish community. Am J Gastroenterol 2006;101:18-28.
- 43. Torrado J, Ruiz B, Garay J, Asenjo JL, Tovar JA, Cosme A, *et al.* Blood-group phenotypes, sulfomucins, and *Helicobacter pylori* in Barrett's esophagus. Am J Surg Pathol 1997;21:1023-9.