



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.1299293>Available online at: <http://www.iajps.com>

Research Article

**ANALYSIS OF CONDITIONS ASSOCIATED TO CHRONIC
UROTHELIAL INFLAMMATION IN LOCAL PAKISTANI
POPULATION**¹Dr. Mafia Akbar, ²Dr. Tayyab Mahmood Ali, ³Dr. Hafsa¹Woman Medical Officer at RHC Khudian Khas, Kasur²Medical Officer at RHC RHC Khudian Khas, Kasur³Woman Medical officer at Pir Mahal, Toba Tek Singh**Abstract:**

Introduction: Recurrent urinary tract infection (UTI) is a very bothersome and a popular problem in the urogynecology clinical practice. According to the IUGA/ICS joint report on the terminology for female pelvic floor dysfunction, recurrent UTI is defined as at least three symptomatic and medically diagnosed UTI in the previous 12 months. **Objectives of the study:** The basic aim of the study is to analyze the conditions associated to chronic urothelial inflammation in local Pakistani population. **Material and methods:** The data was collected from the hospitals of Toba tek singh and Kasur during Jan 2018 to March 2018. The sample size was 50 patients from both genders. Bladder symptoms and lower urinary tract conditions were investigated. The analytical variables included gender, CKD or ESRD, anuria or nonanuria, history of UCC of the bladder, the presence of recurrent UTIs, and clinical symptoms of bladder pain. **Analysis of results:** The analysis of data shows the differences in urodynamic variables between the control subjects and the ESRD/CKD patients. Overall, the ESRD/CKD patients had significantly lower FS, US, and CBC than did the controls. Patients with ESRD/CKD with DU had significantly lower FS, CBC in ESRD/DU patients with BO. **Conclusion:** It is concluded from our study that chronic inflammation, urothelial cell apoptosis and impairment of barrier function of urothelial cells could be the underlying pathophysiology of recurrent UTI in women.

Corresponding author:**Dr. Mafia Akbar,**Woman Medical Officer at RHC Khudian Khas,
Kasur

QR code



Please cite this article in press Mafia Akbar et al., *Analysis of Conditions Associated To Chronic Urothelial Inflammation in Local Pakistani Population*, Indo Am. J. P. Sci, 2018; 05(06).

INTRODUCTION:

Recurrent urinary tract infection (UTI) is a very bothersome and a popular problem in the urogynecology clinical practice. According to the IUGA/ICS joint report on the terminology for female pelvic floor dysfunction, recurrent UTI is defined as at least three symptomatic and medically diagnosed UTI in the previous 12 months. The previous UTI(s) should have resolved prior to a further UTI being diagnosed. Recurrent UTI is one of the most common diagnoses for female pelvic floor dysfunction [1].

The prevalence of urinary tract infections (UTIs) and the incidence of urothelial cell carcinoma (UCC) are also higher in patients with ESRD. Among patients with ESRD, the capacity and compliance of the bladder decrease significantly with the duration of dialysis [2]. In one study, abnormal storage function was noted in up to 71% of ESRD patients and bladder outlet obstruction in 51.6%. Vesicoureteral reflux and high postvoid residual (PVR) urine volumes were observed in 110 of 622 (17.5%) and 83 of 62 patients (13.6%), respectively [3]. Although the bladder capacity increased after kidney transplantation and LUTS remained present in only 31 of 622 of patients (4.9%), patients who did not receive kidney transplants still experienced bothersome bladder symptoms [4]. Patients with end-stage renal disease (ESRD) usually have a small bladder capacity and increased bladder sensation. Patients on chronic dialysis usually have lower urinary tract symptoms (LUTS), such as bladder oversensitivity (BO) or difficulty in urination [5].

OBJECTIVES OF THE STUDY:

The basic aim of the study is to analyze the conditions associated to chronic urothelial inflammation in local Pakistani population.

MATERIAL AND METHODS:

The data was collected from the hospitals of Toba tek singh and Kasur during Jan 2018 to March 2018. The sample size was 50 patients from both genders. Bladder symptoms and lower urinary tract conditions were investigated. The analytical variables included gender, CKD or ESRD, anuria or nonanuria, history of UCC of the bladder, the presence of recurrent UTIs, and clinical symptoms of bladder pain. Each patient was informed of the study rationale and procedures and written informed consent was obtained before cystoscopy and bladder biopsy procedures. All experimental methods were performed in accordance with relevant guidelines and regulations. The urinary bladder specimens were immediately fixed in ice cold 4% formaldehyde phosphate buffered saline (PBS) (pH, 7.4) solution for 1 hour. Next, they were rinsed overnight with ice-cold PBS containing 15% sucrose at 4°C. Then, the specimens were embedded in optimal cutting temperature medium (Miles) and stored at -80°C in liquid nitrogen.

Analysis of data

The collected data were analyzed using SPSS software (version 17). The results are presented as a mean with 95% confidence interval limits or standard deviations. The significant value for P <.05 was accepted as statistically significant.

Analysis of results

The analysis of data shows the differences in urodynamic variables between the control subjects and the ESRD/CKD patients. Overall, the ESRD/CKD patients had significantly lower FS, US, and CBC than did the controls. Patients with ESRD/CKD with DU had significantly lower FS, CBC in ESRD/DU patients with BO (table 01).

Table 01: Analysis of socio demographical variables of patients and control group

Variable	Controls (n = 50)	ESRD/CKD		
		Total	With DU	With BO
Age (yr)	57.9 ± 11.7	59.1 ± 15.0	54.8 ± 11.1	61.0 ± 16.3
FSF (mL)	180.1 ± 65.8	140.2 ± 94.2	63.5 ± 31.8	154.2 ± 95.6
FS (mL)	322.1 ± 81.7	178.3 ± 136.1	66.3 ± 49.6	206.3 ± 137.3*
US (mL)	403.5 ± 104.0	195 ± 133.9	79.3 ± 62.4*	223.9 ± 132.4*
CBC (mL)	404.8 ± 113	204.5 ± 149.1*	79.3 ± 62.4*	235.8 ± 149.2*
Pdet (cm H ₂ O)	24.4 ± 15.7	26.9 ± 20.0	10.5 ± 9.19	29.7 ± 20.2
Qmax (mL/sec)	18.2 ± 11.6	11.7 ± 11.3	0	13.7 ± 11.1
PVR (mL)	51.8 ± 84.0	104.8 ± 164.5	95 ± 77.8	106.4 ± 177.3
Volume (mL)	363.9 ± 175.1	145.7 ± 130.1*	3.33 ± 5.77*	181.3 ± 120.9*
Pves (cm H ₂ O)	32.4 ± 17.1	32.8 ± 20.8	28.5 ± 16.3	33.5 ± 22.0

*P<0.05

Figure 01 shows the infected area in chronic urothelial inflammation in the bladder of patient with CKD.




Figure 01: chronic urothelial inflammation in the bladder of patient with CKD

DISCUSSION:

One recent study demonstrated localized production of mast cell interleukin-10 resulted in suppressed humoral and cell-mediated responses and bacterial persistence. Tissue-resident mast cells not only orchestrate the early innate immunity during bladder infection, they subsequently play a tissue-specific immunosuppressive role which might have association with the recurrent UT I [5]. This observation might explain the mast cell mediated inflammation and related urothelial dysfunction in recurrent bladder infection.

The bladder urothelium is considered not only to act as a barrier, but also to transmit signals of bladder stretching and noxious stimuli. A previous study showed that the antiproliferative factor presented by the urothelium induced increased membrane permeability in cell cultures; regulated the expression of cytokines, which are linked to enhanced purinergic signaling; and mediated increased bladder sensation [6]. Another study revealed that apoptosis was present in the urothelium of patients with IC and showed that it was possibly regulated by inflammatory pathways. Apoptotic signaling molecules were more common in the bladder tissues of IC patients [7]. The increased apoptosis in the bladder urothelium of IC patients could be due to the upregulation of inflammatory signals. In this study, we observed the same patterns of inflammation, urothelial apoptosis, and barrier deficits in bladder samples from ESRD/CKD patients, suggesting that chronic inflammation might be a fundamental form of pathophysiology in the bladders of these patients [8].

Bladder mast cell activation has been reported as a representative pathological finding in a subset of IC patients. Normal basal cell proliferation could be inhibited by chronic inflammation, which might affect apical urothelial function. In this study, the results of TUNEL staining were correlated with those of tryptase staining, indicating that chronic inflammation of the suburothelium was significantly associated with higher levels of urothelial apoptosis in the bladders of patients with ESRD/CKD [9]. These associations demonstrate that inflammation caused increased apoptosis and affected urothelial sensory function in ESRD/CKD patients.

In this study, we did not find a significant correlation between bladder compliance and urodynamic parameters in patients with ESRD/CKD. The expression of E-cadherin and ZO-1 likewise was not significantly lower in ESRD/CKD patients with bladder pain. BO and bladder pain symptoms are likely to occur due to a smaller bladder capacity rather than as a result of defective urothelial barrier function [10].

Recommendations

We suggest that chronic pain syndrome in the bladders of patients with ESRD/CKD could be related to central nervous system sensitization and sustained abnormalities in or activation of the sensory afferent nerves in the urinary bladder. Analysis of the functional proteins in the urothelium might help explain the underlying pathophysiology.

CONCLUSION:

It is concluded from our study that chronic inflammation, urothelial cell apoptosis and impairment of barrier function of urothelial cells could be the underlying pathophysiology of recurrent UTI in women.

REFERENCES:

1. Sant GR, Kempuraj D, Marchand JE, Theoharides TC (2007) The mast cell in interstitial cystitis: role in pathophysiology and pathogenesis. *Urology* 69 (4 Suppl): 34–40.
2. Liu HT, Chancellor MB, Kuo HC (2009) Urinary nerve growth factor levels are elevated in patients with detrusor overactivity and decreased in responders to detrusor botulinum toxin-A injection. *Eur Urol* 56: 700–706.
3. Janssen DA, Hoenderop JG, Jansen KC, Kemp AW, Heesakkers JP, et al. (2011) The mechanoreceptor TRPV4 is localized in adherence junctions of the human bladder urothelium: a morphological study. *J Urol* 186: 1121–1127.
4. Jiang P, Du W, Wu M (2007) p53 and Bad: remote strangers become close friends. *Cell Res* 17: 283–285.
5. Wang H, Wang Z, Chen J, Wu J (2007) Apoptosis induced by NO via phosphorylation of p38 MAPK that stimulates NF-kappaB, p53 and caspase-3 activation in rabbit articular chondrocytes. *Cell Biol Int* 31: 1027–1035.
6. Janssen DA, Hoenderop JG, Jansen KC, Kemp AW, Heesakkers JP, Schalken JA. The mechanoreceptor TRPV4 is localized in adherence junctions of the human bladder urothelium: a morphological study. *J Urol* 2011;186:1121-7.
7. Hauser PJ, VanGordon SB, Seavey J, Sofinowski TM, Ramadan M, Abdullah S, et al. Abnormalities in expression of structural, barrier and differentiation related proteins, and chondroitin sulfate in feline and human interstitial cystitis. *J Urol* 2015;194:571-7.
8. Chuang YC, Fraser MO, Yu Y, Chancellor MB, de Groat WC, Yoshimura N. The role of bladder afferent pathways in bladder hyperactivity induced by the intravesical administration of nerve growth factor. *J Urol* 2001;165:975-9.
9. Wu MY, Chang SJ, Hung SC, Chiang IN. Lower urinary tract symptoms are frequent in dialysis patients. *Perit Dial Int* 2011;31:99-102.
10. Wang SM, Lai MN, Chen PC, Pu YS, Lai MK, Hwang JS, et al. Increased upper and lower tract urothelial carcinoma in patients with end-stage renal disease: a nationwide cohort study in Taiwan during 1997-2008. *Biomed Res Int* 2014;2014:149750.
11. Shie JH, Kuo HC. Higher levels of cell apoptosis and abnormal E-cadherin expression in the urothelium are associated with inflammation in patients with interstitial cystitis/painful bladder syndrome. *BJU Int* 2011;108(2 Pt 2):E136-41.