

Indo American Journal of Pharmaceutical Sciences

CODEN [USA]: IAJPBB

Research Article

http://doi.org/10.5281/zenodo.3237037

ROLE OF PROLACTIN: A COMPARATIVE STUDY BETWEEN GRAND MAL EPILEPTIC AND PSEUDO-SEIZURES PATIENTS Dr Avais Ahmad¹, Dr Farhat Ijaz², Dr Ambreen Tauseef³, Dr. Rana Khurram Aftab⁴, Dr Hira Sohail⁵, Dr Rizwana Kamran⁶, Prof Dr Samia Jawed⁷ and Musarrat Ijaz⁸

^{1,7} Department of Physiology, Continental Medical College, Lahore, ^{2,3} Department of Physiology, CMH LMC & IOD, Lahore, ⁴ Department of Physiology, KEMU, Lahore, ⁵ Department of Biochemistry, CMH LMC & IOD, Lahore, ⁶ Department of Medical Education, CMH LMC & IOD, Lahore, ⁸ Islamia University of Bahawalpur, Bahawalpur.

| Article Received: April 2019 | Accepted: May 2019 | Published: June 2019 |
|------------------------------|--------------------|----------------------|
|------------------------------|--------------------|----------------------|

Abstract:

Background & Objectives:-Epilepsy is a condition of the brain characterized by abnormal, recurrent and excessive discharge from living neurons of brain. It comes in attacks or fits which lasts for a few seconds to a few minutes. It is important to distinguish generalized tonic-clonic seizures (GTCS) from either psychogenic non epileptic seizures (PNES). This study was designed to determine the level of prolactin hormone in females suffering from Grand Mal epilepsy and Pseudo seizures.

Methodology:

After getting approval from Ethical Review Board, this cross-sectional study was conducted in a tertiary care medical teaching hospital over a period of 2 years. Ninety female subjects were enrolled in our study and they were divided in three groups, each of which comprised of 30 subjects: Group-1: Patient with confirmed previous history of grand mal seizures and including freshly diagnosed cases of GTCS. (n=30) Group-2: Patient with history of pseudo seizures. (n=30)Group-3: Control group: Females of comparable age (n=30).Statistical analyses was done using SPSS 21. ANOVA was used to compare the results of the different groups. P<0.001was taken as significant

Results:-

The results showed significant difference of serum prolactin levels among three groups (p value 0.0001). Serum prolactin level was higher in epileptic group as compared to pseudo-seizures and control group (p <0.001).

Conclusion:- It is concluded that serum prolactin may be used as a sensitive biochemical marker in the diagnosis of grand mal seizures in females and is also helpful in differentiating it from pseudo-seizures

Key words:

Prolactin, Generalized Tonic-Clonic Seizures, Pseudo-Seizures

Corresponding author: Dr Farhat Ijaz,

Assistant Professor of Physiology, CMH LMC & IOD, Lahore, E-mail: drranakhurram81@gmail.com

| QR CODE: |
|----------|
| |
| 図の思い |
| |
| |

Please cite this article in press: Avais Ahmad et al. Role Of Prolactin: A Comparative Study Between Grand Mal Epileptic And Pseudo-Seizures Patients. Indo Am. J. P. Sci, 2019; 06(06):11344-11348.

INTRODUCTION:

Grand mal epilepsy, also known as generalized tonicclonic seizure (GTCS) — is caused by abnormal electrical activity throughout the brain ^{(1).}

During the attack , the classical epileptic seizure includes a phase of tonicity showing contraction of facial, jaw, limbs and trunk muscles, followed by jerking of the limbs in the clonic phase. At the beginning of the tonic phase, which usually lasts for about 15-30 seconds, the patients may utter a cry as air is forced out of their lungs, bite their tongue, then become cyanosed as apnoea continues in a position of forced expiration. As the tonic phase passes into clonic phase, the patients may pass urine or occasionally open bowels. The rhythmic jerking of the clonic phase gradually subsides, leaving the patient unconscious or in a state of confusion which may last for some minutes or hours, after which the patients may complain of headache or painful back ^{(2).}

Pseudoseizures (also called psychogenic non-epileptic seizures) are involuntary behavioral changes having no organic cause, showing resemblance to epileptic seizures and are not accompanied by the ictal, perictal and interictal electroencephalical changes(spikes and slowing) that characterize epilepsy. Pseudo seizures may be similar to any kind of epileptic seizures and thus can be misguided for generalized tonic-clonic seizures, absence seizures and simple partial seizures. Pseudoseizure symptoms include indifferent staring, slight motor movements, weird behavior and sweeping movements. Prevalence of non-epileptic seizures in general population all over the world ranges from 2 to 33 cases per 100,000 persons. It is noted that 75 to 85% of the patients of psychogenic non-epileptic seizures are found to be females. These seizures have a tendency to begin in middle age, though the seizures may occur in any age. Psychogenic non-epileptic seizures are found to be more in patients suffering from head injuries, problems of learning or isolated neuropsychological insufficiency.(3).

Clinicians are often faced with the challenging task of distinguishing between PNES and epileptic seizures (ES). Clinical features of ictal events correctly differentiate PNES from ES in less than 70% of cases ^(4,5) Artifacts often obscure scalp EEG during forceful motor movements when a patient's typical events are

recorded during video-EEG. Serum prolactin, a pituitary hormone that is elevated during generalized ES, has been reported to differentiate epileptic from hysterical seizures^{.(6)} However, subsequent studies demonstrated that serum prolactin was elevated in temporal lobe epilepsy but not frontal lobe epilepsy, which may lead to erroneous diagnosis of the latter as PNES^{.(7)}

American Academy of Neurology guidelines regarding the usage of serum prolactin in diagnosing ES stated that elevated serum prolactin 10–20 minutes after a clinical event is perhaps beneficial in separating ES from PNES (Level B) ⁽⁸⁾ Misdiagnosis of PNES has substantial morbidity, social effect, and economic burden on the health care system

Literature review revealed that a lot of previous trials have studied the relationship of serum PRL levels with epileptic events but use of serum PRL levels in clinical practice to diagnose epilepsy or to discriminate epileptic seizures from psychogenic seizures is still controversial because of some limitations which have been described by authors of different studies ^(8,9). One of these limitations includes intrapersonal and interpersonal variations in the basal serum PRL levels, so it is difficult to label significant PRL rise in different individuals. Moreover, most of the seizures occurred at home or nonhospital settings where it is not feasible to measure serum postictal PRL levels, while in hospitals settings video EEG monitoring facility is present which is considered a better tool for this purpose ⁽⁹⁾.

Our study was planned to determine and compare PRL in female Patients of Grand Mal epilepsy and Pseudo seizures.

METHODOLOGY:

This cross sectional study was conducted at King Edward Medical University, Lahore in the Department of Physiology over a period of 2 years. Registered subjects were selected from the indoor as well as outdoor, Emergency Department of Internal Medicine and Psychiatry Department after getting approval from the Ethical Committee of King Edward Medical University Lahore and Mayo Hospital Lahore. Healthy females were recruited from general population. We included young females of child bearing age (16 to 35 years), having history of Grand Mal epilepsy and Pseudo seizures.. Patients with history of head trauma, brain tumour, meningitis, cerbro vascular accidents, metabolic disorders (Hypoglycemia, hypocalcaemia), muscle injuries and mental impairment were excluded. Pregnant and lactating females were also excluded.

The written consent was obtained from the subjects and their guardians at the spot. Ninety female subjects were enrolled and they were divided in three groups equally.

Group-1(n=30): Previous and recently diagnosed patients of grand mal seizures

Group-2 (n=30): Known cases of pseudo seizures.

Group-3 (n=30): Healthy Females (Control)

Under aseptic measures, 5ml of blood sample was collected within 30 minutes of a seizure or pseudo seizure. The samples were sent to centre of Nuclear Medicine (CENUM) of Mayo Hospital, and pathology department of King Edward Medical University Lahore for further processing within 24 hours. Serum prolactin was determined by radio immununo assay (RIA) using an automated Gamma Counter machine (Model-Genesys TM 5000 Series LTI U.S.A). 2 to29 ng/mL was taken as normal range for prolactin

Statistical analyses were done using SPSS 21. Different groups were compared by using ANOVA. Statistical significance was set at P < 0.001.

RESULTS:

The mean age of female included in control, pseudoseizure and epileptic group was 23.4 ± 4.7 year, 21.9 ± 5.5 year and 21.6 ± 4.1 years respectively. The age limit in all females was 16 to 35 years. The mean age in both pseudo-seizure group and epileptic group was not significantly different (P=0.792) but was lower than the control group without significant difference (p= 0.25 & 0.098 respectively). (Table 1)

Mean serum prolactin level was found to be 44.5 ± 5.3 ng/100ml in group 1(epileptic), 8.2 ± 4.8 ng/100ml, in group 2(pseudoseizure), and 4.6 ± 3.2 ng/100ml in group 3 (control) respectively. On applying analysis of variance (ANOVA) we found significant difference between the mean levels of prolactin among three groups. Further statistical analysis showed that in epileptic group prolactin was significantly higher as compared to pseudo-seizures groups and controls (p-value < 0.001) and prolactin was also higher in pseudo-seizures when compared to control group (p-value < 0.001).(Table2,Fig1)

| Variable | Study group | | p-value |
|-------------|--|--|---------|
| Age (years) | Pseudo-seizures (21.90 ± 5.54 year) | Epileptic (22.03 ± 4.62 year) | 0.792 |
| | | Control (18± 1.17 year) | 0.25 |
| | Epileptic (22.03 ± 4.62 year) | Pseudo-seizures (21.90 ± 5.54 year) | 0.792 |
| | | Control (18± 1.17 year) | 0.098 |

| Table -1 Pairwise con | nparison of age (r | mean ± SD years) | among different groups |
|-----------------------|--------------------|------------------|------------------------|
| | I O V | • | 8 8 I |

| Variable | Study group | | p-value |
|--------------------------------|-----------------------------|-----------------------------|----------|
| Prolactin (ng/100ml) | Pseudo-seizure (8.2±4.8) | Epileptic (44.5±5.3) | 0.0001** |
| | | Control (4.6±3.2) | 0.001* |
| | Epileptic (44.5±5.3) | Pseudo-seizure (8.2±4.8) | 0.0001** |
| | | Control (4.6±3.2) | 0.0001** |

Table 2 Pairwise comparison serum Prolactin among different groups

^{*} Significant (P< 0.05) ** Highly significant (P< = 0.0001) w w . i a j p s. c o m



Fig-1: Comparison of mean serum Prolactin in study groups

DISCUSSION:

In our study, prolactin levels were significantly higher in epileptic group as compared to pseudo-seizures groups and control (p-value < 0.001) and prolactin was also higher in pseudo-seizures when compared to control group (p-value < 0.001).

Hypothalamus controls the release of PRL from the pituitary through a PRL inhibitory factor, dopamine.⁽¹⁰⁾ It is thought that generalized neuronal discharge of a seizure may stimulate the hypothalamus either directly, through specific neurotransmitter changes, or indirectly through the release of other substances.⁽¹¹⁾

Epilepsy and pseudo- seizures are important medical problems. Our findings were comparable to the findings of various study groups, ^(12,13) who concluded that PRL can be a valuable measure to differentiate between a generalized tonic–clonic seizure and PNES if it is assessed 10–20 min after an epileptic event. However, baseline PRL level can be measured if the serum PRL test is taken five hours after the epileptic episode.

According to our results, a significant difference in the PRL levels between true epileptic patient (Grand mal) and those with pseudo seizers was noted. Levels of PRL were significantly higher in epileptic group than the pseudo seizure group and mean PRL level in both groups was higher as compared to the healthy control group. These results matched with the study conducted by Priyanka et al ⁽¹⁴⁾ in which PRL was 37 ng/ml in patients of generalized clonic seizures, while in patients of psychogenic non-epileptic seizures, the level was 16.8 ng/ml and in control subjects it was 16.5 ng/ml respectively. Epileptic seizures lead to increase the level of serum PRL hormone due to propagation of epileptic action typically from temporal lobe to hypothalamicpituitary axis. PRL raise was observed in 60% of complex partial seizures which is due to strength of epileptic discharges in a study conducted by Bauer Postictal PRL levels can be used to distinguish epileptic from psychogenic seizures as there was no rise of PRL hormone observed after psychogenic seizures. It is also observed that during the course of status epilepticus PRL secretion may decline in repetitive seizures like in cases of status epilepticus possibly due to reduced spread of ictal discharges (15). Malkowicz et al (16) determined the PRL levels were after 24 hours seizures which were monitored by video EEG intracranial monitoring. It was

seen there was a significant postictal PRL elevations in those patients having longer seizures free intervals while the PRL responses were reduced in subjects having shorter seizures free intervals suggesting that amount of PRL hormone released from the hypothalamus is depleted by seizures or by PRL inhibit feedback mechanism (16). In another study conducted in children it was observed that mean PRL level was 28.6ng/ml in epileptic while in pseudo-seizure and in control groups PRL was 10.4 ng/ml and 9.8 ng/ml respectively. It was proposed that abnormal electrical discharge in epileptic seizures passes through the hypothalamic nuclei responsible for elevation of PRL ^{(17).} Ahmad et al ⁽¹⁸⁾ stated parallel results viewing that generalized epileptic seizure cause noticeable early increase of PRL with more increase following prolonged seizures in children. Thus it may be concluded from these studies that serum PRL is a valuable biochemical indicator to differentiate between epileptic and pseudo-epileptic seizures

CONCLUSION:

An elevation of serum prolactin can be taken as a predictor of epilepsy. Maximum elevation of prolactin is seen within 15 to 30 minutes post ictally. Misdiagnosis of PNES has substantial morbidity, social effect, and economic burden on the health care system.

REFRENCES:

- Sugerman DT. Seizures. JAMA. 2013;310(11):1195.doi:10.1001/jama.2013.27 7840
- 2. Taoufik M, Alsaadi, Anna Vinter Marquez, Psychogenic Non-epileptic Seizures. Am Fam Physician. 2005;72(5):849-856.
- Marie Gillig P. Psychogenic nonepileptic seizures. Innov Clin Neurosci.2013;10(11-12):15–18.
- Desai SD, Desai D, Jani T. Role of Short Term Video Encephalography with Induction by Verbal Suggestion in Diagnosis of Suspected Paroxysmal Nonepileptic Seizure-Like Symptoms. Epilepsy Research and Treatment, 2016; Article ID 2801369,1-6.
- 5. Pedro C. G, Renato L.M. A review of the clinical approach and challenges to psychogenic non-epileptic seizures. Molecular & Cellular Epilepsy 2014; 1: e369.
- Trimble MR. Serum prolactin in epilepsy and hysteria. Br Med J. 1978;2(6153):1682. doi:10.1136/bmj.2.6153.1682

- Meierkord H, Shorvon S, Lightman S, Trimble M. Comparison of the effects of frontal and temporal partial seizures on prolactin levels. Arch Neurol. 1992;49(3):225–230.
- Chen DK, So YT, Fisher RS. Use of serum prolactin in diagnosing epileptic seizures. Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. Neurology 2005;65:668–675.
- 9. Fisher RS. Serum prolactin in seizure diagnosis. Neurol Clin Pract. 2016; 6(2): 100–101.
- 10. Grattan DR. 60 YEARS OF NEUROENDOCRINOLOGY: The hypothalamo-prolactin axis. J Endocrinol. 2015;226(2):T101–T122.
- 11. Aminoff MJ, Simon RP, Wiedemann E. The hormonal responses to generalized tonic-clonic seizures. Brain. 1984;107(Pt 2):569–578.
- 12. Willert C, Spitzer C, Kusserow S, Runge U. Serum neuron-specific enolase, prolactin, and creatine kinase after epileptic and psychogenic non-epileptic seizures. Acta Neurol Scand. 2004;109:318–323.
- 13. Fisher RS, Chan DW, Bare M, Lesser RP. Capillary prolactin measurement for diagnosis of seizures. Ann Neurol. 1991;29:187–190.
- Priyanka P, Yadav P, Ranka R, Kaushik G. Study of serum PRL levels and electrolyte levels in patients with new onset seizures. Int J Adv Res (Indore). 2015; 3(6): 781-786.
- 15. Bauer J. Epilepsy and PRL in adults: clinical review. Epilepsy Res1996; 24(1): 1-7
- Malkowicz DE ,Legido A, Jackel RA, Sussman NM, Eskin BA, Harner RN. PRL secretion following repetitive seizures. Neurology. 1995 ;45(3 Pt 1):448-52.
- 17. Singh U.K, Jana U.K, Plasma prolactin in epilepsy and pseudo-seizures, Indian pediatr,1994. 31(6):667-669.
- Ahmed T. Mahmoud and Akram AM El Deghady. Serum PRL and creatine kinase levels in epileptic and non-epileptic seizures. Alex J Pediatr2005; 19: 1-5