



CODEN [USA]: IAJPBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

<http://doi.org/10.5281/zenodo.2587088>

Available online at: <http://www.iajps.com>

Research Article

ANALYSIS OF STUDY ABOUT FACTORS OF DEPRESSION IN PREGNANCY

¹Dr Irfan Ghani, ²Dr Tayyaba Abdul Hai, ³Dr Mehrosh Shahbaz

¹MO at Civil Hospital, Malakand (KPK), ²WMO at THQ hospital, Khairpur Tamewali, ³Holy family Hospital, Rawalpindi

Article Received: January 2019

Accepted: February 2019

Published: March 2019

Abstract:

Introduction: Maternal health is one of the major goals of reproductive health. Changes in pregnancy make women susceptible to mood and psychological disorders.

Objectives: The purpose of this study is to investigate the prevalence of depression and predicting its related factors in pregnant women.

Materials and Methods: This cross-sectional study was conducted at Civil hospital Malakand during 2018. Beck Depression Inventory (BDI) and demographic information were used as measurement tools. The data was analyzed by descriptive statistics, linear regression and Chi-square test.

Results: The mean age of subjects was 26.45 ± 4.49 . The prevalence rate of depression in pregnant women was 80.5%. The score of depression in the first, second and third pregnancy trimesters was 67.9%, 83.5%, 85.8%, respectively. It was found that depression was significantly increased with the progression of pregnancy ($p = 0.001$). The results of regression analysis showed that the risk factors for depression in pregnancy included female education (OR = 4.67 95% .CI = 1.97-11.07), spouse's education (OR = 4.67 95% .CI = 1.97-11.07), self- employed spouse (OR = 4.67 95% .CI = 1.97-11.07), an increase in the number of children (OR = 4.67 95% .CI = 1.97-11.07), an increase in the number of pregnancy (OR = 4.67 95% .CI = 1.97-11.07), history of abortion (OR= 4.67 95% .CI = 1.97-11.07).

Conclusion: It is concluded that the prevalence of depression in pregnancy is very high and the factors of education, occupation, the number of children, and the number of pregnancies and history of abortion are considered as the variables affecting depression in pregnancy.

Corresponding author:

Dr. Irfan Ghani,

MO at Civil Hospital, Malakand (KPK)

QR code



Please cite this article in press Irfan Ghani et al., *Analysis Of Study About Factors Of Depression In Pregnancy.*, Indo Am. J. P. Sci, 2019; 06(03).

INTRODUCTION:

Pregnancy is an important period in women's life during which brings a lot of emotional, physical and social changes takes place (1). According to WHO, depression disorder is one of the most important causes of women's disability in the world today and approximately 10–15% of women experience *depression during pregnancy*, which is a major risk factor for postpartum depression (2). Various physiological, psychological and social changes, hormonal changes, physical discomforts associated with pregnancy such as nausea, vomiting, and anxiety related to sonography, *anxiety* about the *baby's health* and even the type of delivery are affective in the development of mental disorders including depression during pregnancy (3 and 4).

Depression has negative effects on the mother and the developing fetus (5). Depression during pregnancy can lead to increased stress in pregnancy, self-neglect, lack of prenatal care(6), inadequate nutrition, smoking, pre-eclampsia (7), drug use which may cause low weight of child at the time of birth and preterm delivery (8 and 9). The first psychological symptom of depression includes a decline in physical activity and motivation resulting in decreased activity and efficiency, reduced life expectancy and decreased self-care.

Recent studies suggest that women are more susceptible to depression from the postpartum period than during the pregnancy. Therefore, despite of the relative awareness about the effect of postpartum depression on the child health, mother and family, the issue of depression during pregnancy has been relatively neglected. The timely diagnosis and management of pregnancy during depression is of particular importance.

Objectives of the study

Thus, the aim of this study is to investigate the prevalence of depression and its related factors in pregnant women.

Methodology

This study was conducted in Civil Hospital Malakand during 2018. The study was descriptive and analytic which was performed on 500 pregnant women. The data collection was divided into two sections: the first section consists of the personal characteristics of the subjects under study including age, education, occupation etc. Another questionnaire is the Beck Depression Inventory, which is standard one and has been used for pregnant women and its validity has also been confirmed. In this inventory, a score ranging from 0-15 indicates no depression (normal mean), (16-30) mild depression, (31-45) moderate to severe depression, and (46-63) severe depression (12).

Statistical analysis

The data was analyzed by SPSS 18 and Chi square, as well as linear regression was used for prediction of factors associated with depression.

RESULTS:

The pregnant women aged between 17-45 years with a mean of 27.07 ± 4.88 years. The average age of marriage was 21.18 ± 4.08 . 76.2% of the samples lived in urban areas. The majorities of women (42.7%) had high school degree and were housewives (90.3%). 80.5% of the spouses were self-employed, and most of them (39%) had high school degree. 50.4% of women had a child and 47.7% experienced their first trimester of pregnancy, while 45.9% had their third trimester of pregnancy. 86% of women got pregnant naturally and 59.9% had a natural childbirth history. The frequency of depression in pregnant women showed that 46.1% of women had mild depression and (27.2%) moderate and (7.2%) severe and only 19.5% had no depression.

In comparing the results of depression between third trimesters of pregnancy using Chi-square test, it was found that there was a significant difference between the progression of pregnancy and increasing depression (Table 1).

Table 01: Factors Related to Depression in pregnant women

	N (%)	B	Std. Error	p	OR	95% CI
Educational level				0.007	2.12	2.22-3.68
Uneducated	8(0.9)	0.75	0.28			
Primary school	190(20.9)					
Secondary school	387(42.7)					
College or University	322(35.5)					
Husband's Educational level				0.01	1.24	0.3-5.24
Uneducated	13(1.4)	-0.04	0.02			
Primary school	306(33.7)					
Secondary school	353(38.9)					
College or University	235(25.9)					
Husband Employment status				0.000		
Unemployment	731(80.6)	0.73	13.91		15.57	3.68-65.89
Employed	176(19.4)					
Number of child				0.02		
0	470(51.9)					
1-3	414(45.5)	-1.34	0.60		0.7	0.7-1.2
>3	23(2.5)					
Number of pregnancy				0.006		
First pregnancy	440(48.5)	1.25	0.48		2.81	1.37-5.79
Second pregnancy	337(37.2)					
Tried pregnancy	130(14.3)					
Previous abortion				0.000		
Yes	204(22.5)	-16.18	0.24		3.96	2.22-7.06
No	703(77.5)					

DISCUSSION:

The results of this study indicate that the prevalence rate of depression in pregnant women is 80.5%. On the other hand, the findings of the study by Edward et al and Hosseini Sazi were consistent with our study (9). The low depression score in the first trimester may be attributed to the lack of exposure to pregnancy complications and difficulties increased

in the second and third trimesters due to the progression of pregnancy as well as complications and problems of pregnancy and childbirth. However, there are few studies on the prevalence of depression in pregnancy trimesters and it requires further studies. Regarding the correlation of depression with the studied variables, the results indicate that there is a significant relationship between depression and educational status of women and their husbands.

Low educational status of pregnant women increases the risk of depression by 1.2%. Additionally, in the study by Rahmanie et al., (2011), a significant relationship was found between maternal education and depression (19). Also, the highest percentage of depression was reported in women with elementary education in the study by Hosseini (10). These results when compared with the findings of our study were in line. High educational level raises maternal awareness of their social rights, promotes understanding and cooperation, and provided better and easier access to information resources (3). In the present study, mothers with a history of abortion had a higher prevalence of depression (3.9%). Similarly, in a study by Black et al., the history of abortion has a significant relationship with depression in pregnancy (1).

Furthermore, a study by Loloie et al. (2007) discovered, mothers with abortion history were 2.94 times as likely to be depressed in pregnancy compared with mothers without a history (2). Women with a history of abortion are at increased risk of depression. The results of the study also show that the number of children and the number of previous pregnancies increased 0.7 and 2.8 times the risk of depression.

CONCLUSION:

It is concluded that high prevalence of depression was found in this study, and it is imperative paying attention to the diagnosis of mothers at risk of depression by health care providers. Pregnancy screening for mental and psychological issues in pregnant women should be considered as a part of routine care in pregnancy.

REFERENCES:

1. Lara, M.A., Le, H.N., Letechipia, G., & Hochhausen, L. . Prenatal depression in latinas in the U.S. and Mexico, *Matern Child Health J* 2008 ; 379-384.
2. Abdollahzade Rafi M, Hassanzadeh Avval M, Ahmadi Sh, Taheri M, Hosseini MA Relationship between social support with depression and anxiety during third trimester pregnancy. *Iranian Journal of Nursing Research* 2012; 7 (36): 1-10.
3. Cantwell, R., & Cox, J.L. (2003). Psychiatric disorders in pregnancy and the puerperium, *Current Obstetrics and Gynecology*, 13, 7–13.
4. Campagne, D.M.. The obstetrician and depression during pregnancy, *European Journal of Obstetrics and Gynecology and Reproductive Biology*:2004: 116(6), 125– 130.
5. Karacam, Z., & Ancel, G. (2007). Depression, anxiety and influencing factors in pregnancy: a study in Turkish population, *Midwifery* 2007; 10, 1016-1029.
6. Bayrami M, Zahmatyar H, Bahadori KJ. Prediction strategies to coping with stress in the pregnancy women with first experience on based factors hardiness and social support. *Iran J Nurs Res* 2013; 7(27):1-9.(Persian).
7. Moafi F, Dolatian M, Keshavarz Z, Alavi Majd H. ASSOCIATIONS OF DEPRESSION AND PREECLAMPSIA DURING PREGNANCY. *The Journal of Urmia Nursing and Midwifery Faculty*, 2014; 12(4).
8. Jabbari Z, Hashemi H, Haghayegh SA. Survey on effectiveness of cognitive behavioral stress management on the stress, anxiety and depression of pregnant women. *J Health Syst Res* 2013; 8(7):1341-7. (Persian).
9. M. Ranjbar, M. Heydarzadeh, M. Yazdani, M. Mirghafourvand, F. Mirza Ahmadi, Depression in Women during Pregnancy and After Delivery, *journal of advanced psychological research*.2016:10(40):1-15
10. Ghanei R, Golkar F, Aminpoor E. Foot care in depressed and non-depressed diabetic patients. *Modern Cares J* 2013; 10(2):124-31. (Persian).