



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3708946>Available online at: <http://www.iajps.com>

Research Article

**RELATIONSHIP AMONG BLOOD VESSEL STRENGTH AND
PLATELET ACTIVATION IN NORMOTENSIVE,
OVERWEIGHT AND CORPULENT YOUNG ADULTS**

Dr. Alisha Anwar, Dr. Rimsha Anwar, Dr. Amna Noreen

THQ Hospital Kallur Kot Bakkar Pakistan

Article Received: January 2020 Accepted: February 2020 Published: March 2020

Abstract:

In obese people, platelet initiation and blood vessel firmness are high, but the quality and transient nature of connection among those components remains unclear. Authors plan to regulate impact of blood vessel strength on platelet action in overheavy or obese young adults. The current investigation involved 93 members (average age 42 years, 65 females) in SAVE (Slow Contrary Vascular Effects of overabundance weight), a preliminary medical study exploratory impacts of lifestyle intercession with or without sodium limitation on vascular well-being in normotensive/overheavy young grownups. This existing research was conducted at THQ Hospital Kalur Kot Bakhar Pakistan from March 2018 to February 2019. The carotid-femoral (cf), lower brachial leg (ba) and lower femoral leg (fa) were assessed for blood vessel firmness and were estimated at gauge and 6, 12 and two years of development. Platelet movement was estimated in plasma b-thromboglobulin (b-TG) at two years. Higher plasma b-TG was associated through a higher presentation for PWV ($p=0.03$) and PWV ($p=0.05$) over the previous two years. After modification for serum leptin, the increased introduction of elevated b-TG remained substantial ($p=0.05$) and the presentation of raised b-TG to the imperceptibly large PWV ($p=0.056$) in anticipation of increased plasma b-TG. The increased blood vessel strength, particularly the firmness of the focal blood vessels, suggests an increase in platelet production in overheavy and obese people. This association may incompletely clarify the relationship between blood vessel firmness and atherothrombotic episodes.

Keywords: Arterial stiffness, obesity, platelet activation, pulse wave velocity, weight loss.

Corresponding author:

Dr Alisha Anwar,

THQ Hospital Kallur Kot Bakkar Pakistan

QR code



Please cite this article in press Alisha Anwar et al., *Relationship Among Blood Vessel Strength And Platelet Activation In Normotensive, Overweight And Corpulent Young Adults.*, Indo Am. J. P. Sci, 2020; 07(03).

INTRODUCTION:

Platelets play a very significant role in thrombotic vascular actions and in initiation also progression of atherosclerosis. Platelets issue challenge atoms and developmental aspects in addition to endothelial beginning also relocation and multiplication of vascular smooth muscle cells, altogether of these are important procedures in atherosclerosis [1]. In inverted stage, platelet initiation is activated by the disintegration of endothelial cells and causes atherosclerotic plaques to burst, also by high shear pressure. Blood vessel strength is a marker of vascular well-being, and stiffer focal pathways increase divider shear and ductile anguish, accelerate fatigue of the divider parts of the blood vessels, and advance atherosclerotic plaque impotence throughout the vascular system [2]. Therefore, it is logical that increased blood vessel strength may also advance a prothrombotic phenotype, including more remarkable platelet activation, throughout the vascular tree. By estimating beat wave velocity (PWV), blood vessel firmness can be assessed in any area of the blood vessel tree [3]. In the past, small cross-sectional studies in obviously healthy adults have observed the relationship between platelet initiation and carotid-femoral heart beat velocity (cf. PWV), a proportion of aortic strength and an independent indicator of cardiovascular events. Different studies in obviously healthy adults have found a relationship between some markers of platelet activation in vivo and lower brachial-leg beat wave velocity (ba PWV), a proportion of focal (aortic) and marginal blood vessel firmness [4]. These cross-linkages can be clarified either by the impact of activated platelets on the vascular system or by the impact of vascular injury and rupture on platelets during formation. The purpose of this review was to determine the future relationship between the introduction of greater blood vessel firmness, estimated repeatedly during a one-year lifestyle mediation and again one year after the procedure, and the action of ongoing platelet action, as estimated by b-thrombic plasma globulin (b-TG)(15,16) at the last point in time of the survey, in overweight and overweight young adults. Authors estimated that, unlike PWV ba and femoro-lower leg PWV (fa), PWV cf would be more clearly related to platelet action due to impact of aortic firmness on divider stress, blood flow design and atherosclerotic movement along the blood vessel tree. Authors further estimated that relationship between presentation of higher CPAP after a certain time and greater platelet action could remain free of impacts of the introduction of other cardiovascular and metabolic danger aspects throughout two years of research [5].

MATERIALS AND METHODS:

Study population:

This existing research was conducted at THQ Hospital Kallur Kot Bakkar Pakistan from March 2018 to February 2019. The carotid-femoral (cf), lower brachial leg (ba) and lower femoral leg (fa) were assessed for blood vessel firmness and were estimated at gauge and 6, 12 and two years of development. Platelet movement was estimated in plasma b-thromboglobulin (b-TG) at two years. The SAVE (Slow Adverse Vascular Effects of overabundance weight) study is a preliminary randomized controlled study that evaluates the impacts of weight reduction, increased physical movement and decreased dietary sodium intake on vascular well-being. Members were recruited from June 2007 to May 2009 by mass mailing. Qualified members were overweight or corpulent (body mass index (BMI) 26.1 to 37.8 kg/m²) and truly sleepy (58 months of usual physical activity in the previous year), male and female, aged 22 to 46 years, with the exception of 1) diabetes, 2) hypertension or normal circulatory pressure of $\geq 140/90$ mmHg, 3) cholesterol-lowering, antipsychotic or vasoactive medication, and 4) current pregnancy or lactation. Members who gave a blood test for b-TG estimation at the last examination visit were recalled for this investigation (n=494). As this sub-study was initiated more than two years after the start of the preliminary parent clinical study and many subjects were excluded from this sub-study due to their continued use of headache medications or NSAIDs, the sample size for this survey was quite small compared to the total number of preliminary members (n=4351).

Blood and urine testing:

Blood tests were estimated at the Heinz Laboratory at the Graduate School of Public Health, University of Pittsburgh, using the standard strategies described above. Rapid blood tests were performed between 7:52 a.m. and 11:40 a.m. on subjects fasting for at least 10 hours. The intra- and inter-test CV% for insulin were 5.8% and 11.6%, separately. The CV% for the different tests were all 55%. 24-hour pee sets were performed during approximately 14 days of visits to the center where all other estimates were resolved. Assortments considered substantial had a volume of 500 to 4000 mL, a term of ≥ 24 to ≥ 28 of manual creatinine in the normal range. Sodium, potassium and creatinine were resolved as described above.

RESULTS:

Plasma b-TG was estimated in 96 individuals at 26-month visit of preliminary medical parent. The example had a mean duration of 41.3 years (SD 6.8) and included 64 females, 14 African-Americans, and 9 existing and 24 former smokers. In cross-sectional surveys, high plasma b-TG was measurably associated by developed BMI ($r=0.26$, $p=0.03$), higher leptin ($r=0.21$, $p=0.049$), and

higher PPV ba ($r=0.23$, $p=0.05$), and barely fundamentally related to higher PPV cf ($r=0.18$, $p=0.08$) and PPV fa ($r=0.21$, $p=0.07$). Essentially, higher plasma b-TG was related to higher BMI and leptin when these variables were studied as AUC from baseline to the two-year visit (Table 2). In adding, higher plasma b-TG was actually huge with

higher GWP cf and lower GWP ba once measured as AUC (Figure 1). At the time the collaborations between each free factor and the treatment group in the preliminary clinical trials were inspected, none were considered critical in any model for plasma b-TG.

Table 1. Clinical characteristics over the course of the study.

Characteristic	Baseline (n/492)	6 Months (n/490)	12 Months (n/490)	24 Months (n/493)	AUC (n/493)
Waist circumference	96.0 (12.2) *	101.6 (11.0)	98.1 (12.6) *	95.0 (11.7) *	96.4 (10.6)
BMI (kg/m ²)	29.9 (4.2) *	32.4 (3.6)	31.1 (4.2) *	29.7 (4.0) *	30.3 (3.7)
Pulse pressure (mm Hg)	37.8 (7.5)*	37.0 (7.1)*	40.0 (7.6)	38.6 (4.8)	39.3 (7.6)
Mean arterial pressure (mmHg)	85.7 (7.8)	83.3 (7.2)*	84.6 (5.4)	86.4 (7.6)	84.1 (7.6)*
Insulin (mU/mL)y	(9.6, 16.2)	11.3 (9.1)	98.5 (6.1)	11.4 (9.7, 14.3)	12.0 (9.3, 16.0)
Glucose (mg/dL)	98.9 (8.8)	98.2 (9.2)	98.8 (9.2)	98.1 (10.4)	12.114.9)*

Table 2. Pearson associations among b-thromboglobulin at 24-months:

Variable	r	pValue
Waist circumference	0.48	0.07
Mean arterial pressure	0.64	0.06
BMI	0.008	0.29
LDL-C	0.52	0.07
Triglycerides	0.39	0.09
Pulse pressure	0.17	0.14
HDL-C	0.58	0.06




Figure 1. Scatter plots of average pulse wave velocity (PWV) over 24-months:**DISCUSSION:**

The foremost conclusion of the current review was that, in young, overweight and corpulent, but still strong adults, more fluid platelet movement, as estimated by plasma b-TG, was anticipated by greater introduction into stiffer corridors throughout first 2 years [6]. Nevertheless, the more notable introduction of weight abundance and serum leptin over a similar period, when inspected together, expelled the measurable immensity of this affiliation [7]. Our findings recommend that high blood vessel firmness, particularly focal blood vessel strength, may be one of components by which platelet initiation is expanded in overweight and tall individuals [8]. Endothelial microparticles, which discharge at endothelial cell initiation and are firmly identified with a decrease in the auxiliary and useful respectability of the endothelium, are definitely related to BPV ba in DM type-2 and fit grownups. Second, shear pressure plays a significant part in platelet activation and is affected by blood vessel firmness [9]. Pathologically high shear pressure initiates platelets and is an important factor in the disposition of thrombi in blocked vessels. However, even in people with negligible atherosclerosis, as the focal veins solidify, the velocity of blood flow increases, resulting in increased shear pressure and creating a waveform of precarious systolic weight that improves the flexibility of shear concerns in the peripheral vessels. Such oscillatory shear pressure can cause the prothrombotic, prooxidative and pro-inflammatory state in vascular endothelial cells, mainly in less consistent vessels [10].

CONCLUSION:

Overall, in cases of overweight and corpulence, but in any case, in youth, a more remarkable presentation of blood vessel firmness over a two-year period is an indicator of a more remarkable circular movement of platelets, as estimated by plasma b-TG. A more remarkable presentation of extra mass and serum leptin is similarly related to greater platelet action. Those results recommend that high blood vessel firmness, particularly focal blood vessel strength, may be one of systems by whom platelet action remains expanded in overheavy and corpulent persons. Forthcoming research on lifestyle changes also additional procedures to "harden" blood vessels in overheavy in addition overweight adults should examine whether a continued decrease in blood vessel firmness can decrease danger of thrombosis.

REFERENCES:

1. Protogerou AD, Papaioannou TG, Blacher J, et al. Central blood pressures: do we need them in the management of cardiovascular disease? Is it a feasible therapeutic target? *J Hypertens* 2007;25: 265–72.

2. Vlachopoulos C, Aznaouridis K, O'Rourke MF, et al. Prediction of cardiovascular events and all-cause mortality with central haemodynamics: a systematic review and meta-analysis. *Eur Heart J* 2010; 31:1865–71.
3. Weber C. Platelets and chemokines in atherosclerosis: partners in crime. *Circ Res* 2005;96:612–16. Malik AR, Kondragunta V, Kullo IJ. Forearm vascular reactivity and arterial stiffness in asymptomatic adults from the community. *Hypertension* 2008;51:1512–18.
4. Kobayashi K, Akishita M, Yu W, et al. Interrelationship between non-invasive measurements of atherosclerosis: flow-mediated dilation of brachial artery, carotid intima-media thickness and pulse wave velocity. *Atherosclerosis* 2004; 173:13–18.
5. Matzdorff AC, Kemkes-Matthes B, Voss R, Pralle H. Comparison of beta-thromboglobulin, flow cytometry, and platelet aggregometry to study platelet activation. *Haemostasis* 1996;26:98–106.
6. Bernstein AM, Willett WC. Trends in 24-h urinary sodium excretion in the United States, 1957–2003: a systematic review. *Am J Clin Nutr* 2010;92:1172–80.
7. Cooper JN, Tepper P, Barinas-Mitchell E, et al. Serum aldosterone is associated with inflammation and aortic stiffness in normotensive overweight and obese young adults. *Clin Exp Hypertens* 2012; 34:63–70.
8. Jennings LK. Mechanisms of platelet activation: need for new strategies to protect against platelet-mediated atherothrombosis. *Thromb Haemost* 2009;102:248–57.
9. Miyazaki Y, Nomura S, Miyake T, et al. High shear stress can initiate both platelet aggregation and shedding of procoagulant containing microparticles. *Blood* 1996;88: 3456–64.