



CODEN [USA]: IAJPBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

<http://doi.org/10.5281/zenodo.2861571>

Available online at: <http://www.iajps.com>

Research Article

STUDY TO EVALUATE THE TECHNICAL QUALITY OF ROOT CANAL TREATMENT RADIOGRAPHICALLY DONE BY THE HOUSE SURGEONS OF DENTAL DEPARTMENT

* Dr Syeda Kaniz Umul Baneen, **Dr Bakhtawar Tahir, **Dr Hafiz Muhammad Adeel Sarfraz, ** Dr Zafar Iqbal Sajid, **Dr Mohsin Majeed, **Dr Sara Izhar

* De'Montmorency College of Dentistry, Lahore

** Nishtar Institute of Dentistry, Multan

Article Received: March 2019

Accepted: April 2019

Published: May 2019

Abstract:

Objective: The aim of this study was to evaluate the technical quality of root canal treatment by house surgeons in a dentistry hospital.

Study Design: A Pilot Study.

Location and duration: In the Dental Department of Nishtar Hospital, Multan for one year duration from February 2018 to January 2019.

Methods: To determine the adequacy of the treatment, radiographic recordings of 64 teeth treated with root canal were analyzed. The last sample consisted of 60 teeth. Root treatment was considered to be sufficient if the apical length of the filling was 0 to 2 mm from the radiographic apex and there was no clearance on the radiograph.

Results: Of the 60 teeth, 27 (45%) were classified as adequate. Accurate radiographic apical length (<2 mm) 35 (58.3%), 44% (73.3%) on radiography was not found any gap. There were 5 teeth fracture due to instruments and this was significantly correlated with more than 20 ° channel curvature (p = 0.01). There was no correlation between the adequacy of the treatment and the type of the tooth or the adequacy of the treatment and canal curvature.

Conclusion: Sufficient root treatment assessed by radiography was performed in only 45% of the treated teeth.

Key words: radiographic technical quality, root canal fillers.

Corresponding author:

Dr.Syeda Kaniz Umul Baneen

De'Montmorency College of Dentistry, Lahore

QR code



Please cite this article in press Syeda Kaniz Umul Baneen et al Study To Evaluate The Technical Quality Of Root Canal Treatment Radiographically Done By The House Surgeons Of Dental Department., Indo Am. J. P. Sci, 2019; 06(05).

INTRODUCTION:

In Pakistan, the awareness of the preservation of natural dentists has increased and, as a result, more and more people seek endodontic treatment to protect their natural teeth, rather than extracting pain to alleviate pain and treat excessively decayed teeth. In Nishtar Hospital, Multan, preclinical endodontics were taught during a two-month rotation in the third month. Last year, students are expected to perform endodontic treatments on patients in hospital under supervision. During a mandatory one-year housework, patients are expected to perform endodontic treatments once more. If necessary, they can get help from the faculty. The quality of root canal treatment carried out by home surgeons reflects the level of treatment offered to the community, especially since older supervisors cannot be guided. When done correctly, endodontic treatment is very successful, with reported success rates between 85% and 95% (Orstavik et al., 1987) when performed by experts, but decreased by 65% to 75% when performed by experts and general dentists (Eriksen 1991). Friedman 1998). This can be the cause of the standard when working in an experimental environment. Endodontic treatment is much higher. In other configurations, this may not be the case. Boltacz-Rzepakowska and Pawlicka (2002) found that 71% of root-treated teeth were not properly filled in dentistry departments in the Lodz region of Poland. Teeth with sufficient root fill are more likely to have a healthier periapical condition than inadequate fullness (Kirkevang et al., 2000). The gaps in the channel filling and inadequate lateral sealing are also associated with periapical disease in endodontically treated teeth (Chugal et al. 2003, Kirkevang et al. 2000, Dugas et al 2003). Broken instruments have also been shown to result in higher rates of endodontic failure (Crump et al., 1970). Our hypothesis is that root canal treatment is more likely to have a root curvature of at least 20 ° in the teeth and roots, and that molar root canals are performed in a weaker, more independent way. in presence or absence of curvature of the root.

MATERIALS AND METHODS:

This Pilot Study was held in the Dental Department of Nishtar Hospital, Multan for one year duration from February 2018 to January 2019 for one year duration from February 2018 to January 2019. 64


teeth of patients with 119 roots were included in the study. These were the first 64 root canals that were completed. It was carried out by assistant surgeons in the Department of Operative Dentistry of the Nishtar Hospital, Multan. The poor quality of radiographs represented 4 excluded cases. Therefore, 60 teeth with 112 roots were included in the study. The cases performed by other surgeons were pulpal and periapical diseases, elective endodontics and withdrawal. The length of the study was determined by radiographic method. Radiographs were explained with information technology. The channel was prepared using the back-closure method using stainless steel K-files (Mani). The filler was caused by the cold lateral condensation of gutta-percha and the filler used was endomethasone (Septodont). EDTA was used when the pulp chamber was found in the pulp chamber or suspected in the root canals, otherwise irrigation was limited to normal saline use. The following x-ray films were analyzed retrospectively using a magnifying lens (x 2 and x 4) and an X-ray image examining radiographs in a dark room. For measurements, a transparent scale was used with 0.5 mm calibrations. The curvature of the root canal was determined by measuring the internal angle generated by the intersection of two lines through which one draws from the coronal portion of the hole to the hole and the other to the apical part of the apex. Suitable root canals were defined as those in which the apical fill length was <2 mm in the radiographic apex and did not have visible cavities in filling the canal or adjacent to the walls. Inadequate fillings are fillers >> 2 mm below the apex or extended beyond the radiographic apex. The presence of a gap between the filler or the channel wall and the fill made the filling insufficient. The results were analyzed using SPSS. The chi-square test was used to determine whether there was a significant relationship between the fracture instruments, the adequacy and curvature of the treatment, and the adequacy of the channel against the tooth type. Significance was determined at a value of 0.05.

RESULTS:

64 teeth with 119 roots were evaluated. 4, there were X-rays that were not of sufficient quality to be tried, so they were not included. Finally, 60 teeth were evaluated. There were 112 roots. Of the 60 teeth, 27 (45%) were adequate and 33 (55%) were inadequate.

Table No 01: Radiographic Adequacy of Root-canal Treatment According to Tooth Type


<i>Teeth Type</i>	Quantity	Adequate	Inadequate
<i>Maxillary</i>	28	15	13
<i>Incisor</i>	05	03	02
<i>Canine</i>	05	04	01
<i>Premolar</i>	10	04	06
<i>Molar</i>	08	04	04
<i>Mandibular</i>	32	12	20
<i>Incisor</i>	02	02	0
<i>Canine</i>	01	0	01
<i>Premolar</i>	12	06	06
<i>Molar</i>	17	04	13
<i>Total</i>	60	27	33



Of the 60 teeth, 35 were placed correctly, taking into account the apical length of the obstruction. Therefore, 58.33% of all teeth were satisfactorily filled according to the apical length criteria of occlusion. 16 (26.7%) were short and 9 (15%) were filled.

Table No 02: Apical Length of Obturation


Length of obturation	Teeth Frequency	%	Roots Frequency	%
>2mm from radiographic apex	16	26.7	27	24.1
0-2mm from radiographic apex	35	58.3	66	58.9
Over Obturated	09	15	19	17
Total	60	100	112	100



Of the 112 roots, 66 (58.9%) were accepted within 0-2 mm of the radiological apex, 27 (24.1%) were placed below, and 19 (17%) were overdosed. Data on the teeth treated according to the presence or absence of gaps in the filling.

Table No 03: Voids in the Obturation

Length of obturation	Teeth Frequency	%	Roots Frequency	%
Uniform Density of Obturation	44	73.3	91	68.8
Void Present in Obturation	16	26.7	21	25
Total	60	100	112	100



DISCUSSION:

This study consisted of 60 teeth of all types with straight and root canals. 45% of all teeth were considered acceptable. The success criteria were 0-2 mm roots from the radiographic apex without gaps. Most of the studies have used the radiographic criteria of the technical quality radiographic criteria 0-2 mm to demonstrate the appropriateness (E. Boltacz-Rzepakowski and H Pawlicka (2003) peak. 2005) Lambriadis Eleftheriadis license treated by students of 620 dental records were obtained and 55.3% when the apical filling level evaluation reported that sufficient quality seal, roots and holes .. radiographic apex 0-2 mm and leisure 82.6% stake others, for the license in Turkey A similar criterion was added to the continuation of shrinkage. Root filling The resultant qualification rate was 33%, 69% had apical filling, 53.2% had sufficient radioizite and 68.3% had a gradual decrease. a) 6% 4 is filled appropriately in the Dental Institute. The seniority of the operator has not been discussed, so it may have been made by students, assistants and senior professors. The criterion for adequate obturation was only the apical length of the obturation, which could have increased the percentage of teeth that were considered adequate compared to other studies. The gaps in the channel filling material were not used as a criterion for adequate treatment. In this study, 58.9% of the roots and 58.33% of the teeth were filled with 2 mm of radiographic vertex unless the gaps in the filling were taken into account. These results are very similar. While Hayes et al. (2001) believed that more specialized endodontists were needed to improve treatment standards, Dugas et al. (2003) found that specialist training did not significantly improve the

clinical success rate. However, more people need to be included in the study to confirm the results, he said. Although the quality of radiographic technique is considered to affect the clinical outcome, other factors also play a role. For example, debridement and reduction of the bacterial population, placement of an appropriate coronal seal (Eriksen et al., 2002).

CONCLUSION:

Within the limitations of this study, it was found that 45% of all teeth could be considered in an acceptable way when evaluated radiologically. This study did not show a tendency to inadequate obturation in curved roots compared to smooth roots.

REFERENCES:

1. Silveira, Stephanie B., Flávio RF Alves, Marília F. Marceliano-Alves, Júlio César N. Sousa, Victor TL Vieira, José F. Siqueira Jr, Hélio P. Lopes, and José C. Provenzano. "Removal of Root Canal Fillings in Curved Canals Using Either Mani GPR or HyFlex NT Followed by Passive Ultrasonic Irrigation." *Journal of endodontics* 44, no. 2 (2018): 299-303.
2. Pirani, C., Friedman, S., Gatto, M.R., Iacono, F., Tinarelli, V., Gandolfi, M.G. and Prati, C., 2018. Survival and periapical health after root canal treatment with carrier-based root fillings: five-year retrospective assessment. *International endodontic journal*, 51, pp.e178-e188.
3. Keleş, A., Torabinejad, M., Keskin, C., Sah, D., Uzun, İ. and Alçın, H., 2018. Micro-CT evaluation of voids using two root filling techniques in the placement of MTA in mesial root canals of Vertucci type II

- configuration. *Clinical oral investigations*, 22(5), pp.1907-1913.
4. Moinzadeh, Amir-Teymour, Roeland JG De Moor, and Mieke AA De Bruyne. "Influence of a calcium hydroxide-based intracanal dressing on the quality of the root canal filling assessed by capillary flow porometry." *Clinical oral investigations* (2018): 1-7.
 5. Rödig T, Wagner J, Wiegand A, Rizk M. Efficacy of the ProTaper retreatment system in removing Thermafil, GuttaCore or vertically compacted gutta-percha from curved root canals assessed by micro-CT. *International endodontic journal*. 2018 Jan 24.
 6. Orhan, Kaan, Reinhilde Jacobs, BerkanCelikten, Yan Huang, Karla de Faria Vasconcelos, Laura Ferreira Pinheiro Nicolielo, ArdaBuyuksungur, and Jeroen Van Dessel. "Evaluation of Threshold Values for Root Canal Filling Voids in Micro-CT and Nano-CT Images." *Scanning* 2018 (2018).
 7. Connert, T., Truckenmüller, M., ElAyouti, A., Eggmann, F., Krastl, G., Löst, C. and Weiger, R., 2018. Changes in periapical status, quality of root fillings and estimated endodontic treatment need in a similar urban German population 20 years later. *Clinical oral investigations*, pp.1-10.
 8. Kim, Jung-Ae, Yun-Chan Hwang, Vinicius Rosa, Mi-Kyung Yu, Kwang-Won Lee, and Kyung-San Min. "Root canal filling quality of a premixed calcium silicate endodontic sealer applied using gutta-percha cone-mediated ultrasonic activation." *Journal of endodontics* 44, no. 1 (2018): 133-138.
 9. Fong, W., Heidarifar, O., Killough, S., Lappin, M.J. and El Karim, I.A., 2018. An audit on technical quality of root fillings performed by undergraduate students. *International endodontic journal*, 51, pp.e197-e203.
 10. Siqueira Junior, José Freitas, Isabela das Neves Rôças, MaríliaFaguryMarceliano-Alves, Alejandro Ron Pérez, and Domenico Ricucci. "Unprepared root canal surface areas: causes, clinical implications, and therapeutic strategies." *Brazilian oral research* 32 (2018).
 11. Elshinawy, Mohamed I., Hossam A. Eid, Ashraf A. Khalil, Abdelnasser M. Soliman, and Hesham El-Refaey. "TNF- α -mediated inflammation and cytotoxicity of different root canal sealers and root-end fillings in cell line culture." *Tanta Dental Journal* 15, no. 1 (2018): 27.
 12. Saatchi M, Mohammadi G, Sichani AV, Moshkforoush S. Technical Quality of Root Canal Treatment Performed by Undergraduate Clinical Students of Isfahan Dental School. *Iranian endodontic journal*. 2018;13(1):88.
 13. Ribeiro, D. M., J. C. Reus, W. T. Felipe, C. Pachêco-Pereira, K. L. Dutra, J. N. Santos, A. L. Porporatti, and G. De Luca Canto. "Technical quality of root canal treatment performed by undergraduate students using hand instrumentation: a meta-analysis." *International endodontic journal* 51, no. 3 (2018): 269-283.
 14. Costa, F.F.N.P., Pacheco-Yanes, J.A., Siqueira Jr, J.F., Oliveira, A.C.S., Gazzaneo-Rauseo, I.D., Amorim, C.A., Santos, P.H.B. and Alves, F.R.F., 2018. Association between missed canals and apical periodontitis. *International endodontic journal*.
 15. Neelakantan P, Ahmed HM, Wong MC, Matinlinna JP, Cheung GS. Effect of root canal irrigation protocols on the dislocation resistance of mineral trioxide aggregate-based materials: A systematic review of laboratory studies. *International endodontic journal*. 2018 Jan 29.
 16. Hendi, SeyedehSareh, Hamed Karkehabadi, and Amir Eskandarloo. "Iatrogenic Errors during Root Canal Instrumentation Performed by Dental Students." *Iranian endodontic journal* 13, no. 1 (2018): 126.