Saba Sabahat et al



## CODEN [USA]: IAJPBB

ISSN: 2349-7750

# INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

http://doi.org/10.5281/zenodo.4444389

Available online at: <u>http://www.iajps.com</u>

**Research Article** 

# A RESEARCH STUDY TO COMPARE THE EFFECTIVENESS OF VERAPAMIL AND INTRAVENOUS ADENOSINE IN ACUTE PAROXYSMAL SUPRAVENTRICULAR TACHYCARDIA

<sup>1</sup>Dr. Saba Sabahat, <sup>2</sup>Dr. Nida-e-Farah, <sup>3</sup>Dr. Merab Ahsan

<sup>1</sup>Tehsil Headquarter Hospital Sambrial, <sup>2</sup>WMO, RHC Kassoke Hafizabad, <sup>3</sup>Woman Medical Officer, Civil (THQ Level) Hospital, Dinga.

Article Received: March 2019	Accepted: April 2019	Published: May 2019

## Abstract:

**Background:** The most frequently observed arrhythmia is the severe convulsive supraventricular tachycardia. The antibiotics used initially after the letdown of vagal manoeuvres adenosine to lapse severe PSVT. The antibiotics can be selected on the basis of their prices, need of action and contraindication.

*Objective:* The main purpose of the study was to compare the effectiveness of intravenous adenosine and verapamil to transform a severe section of paroxysmal supraventricular tachycardia into the sinus rhythm.

Subjects and Methods: The investigational study was arranged in Jinnah Hospital, Lahore (August 2017 to March 2018). The 200 patients suffering from severe PSVT present in the ICU were added in the study. A dose of 6 mg adenosine 4 boluses was given to the patients added in Group A. Then the dose of 12 mg 4 boluses was given to a similar group of patients. The second option is applicable when the PSVT was not reversed. Verapamil 5 mg was given to the patients present in Group B. Continuous dosage of 5 mg verapamil was given to the patients with the break of about 5 minutes unless complete success was achieved. The time required to rebuild the sinus rhythm from the execution of antibiotics in the body was recorded. The patients were also undergone ECG. ECG was performed on patients prior to and 15 minutes later to the cure. When the treatment was not completed by PSVT, patients were treated with the incorporation of verapamil in Group A.

**Results:** The age limits of the patients present in Group A were between  $(42.8 \pm 13.9)$  years. The age of Group B was about  $(40.5 \pm 15)$  years. The presence of males is slightly greater in group A which was 58%. In group B the males were lesser in number about 49%. The achievement rate was 96% and 95% accordingly. The first dose of adenosine was found to be more successive in finishing PSVT. It was noticed that the adenosine take less time to perform its function than verapamil. The difference was  $(2.1 \pm 2.7)$  to  $(5.1 \pm 3.5)$  minutes.

*Conclusion:* The success rate of adenosine and verapamil is almost equal but the speed was different in both cases. The 12 mg doses of adenosine act rapidly and efficiently and remove the severe segments of paroxysmal supraventricular tachycardia. **Keywords:** Paroxysmal supraventricular tachycardia, Adenosine, Verapamil, Vagal maneuvers.

**Corresponding author:** 

Dr. Saba Sabahat,

Tehsil Headquarter Hospital Sambrial.



Please cite this article in press Saba Sabahat et al., A Research Study to Compare the Effectiveness of Verapamil and Intravenous Adenosine in Acute Paroxysmal Supraventricular Tachycardia., Indo Am. J. P. Sci, 2019; 06(05).

www.iajps.com

Saba Sabahat et al

### **INTRODUCTION:**

PSVT is the frequently observed tachyarrhythmia. It is present in about 2.25 patients out of 1000 patients. Most of the patients suffering from PSVT visit the emergency units of clinics [1 – 5]. All tachyarrhythmia that initiates in supraventricular cells has spontaneous inception and butchery [6]. The two major kinds of PSVT are atrioventricular nodal reentrant tachycardia and atrioventricular re-entrant tachycardia. There exist some other kinds of PSVT which are not too common. These include atrial tachycardia, paroxysmal atrial flutter and paroxysmal atrial fibrillation [6]. The words PSVT is considered to be interrelated with AVNRT and AVRT [7]. Special features like tachycardia with QRS composite of supraventricular in derivation, abrupt inception and execution can characterize the re-entrant tachycardia including atrioventricular nodal tissue. The speed of inception and execution are generally lain between 150 and 250 beats in every minute with a continuous pulse on electrocardiogram (ECG) [7]. We can identify PSVT by continues R-R intermissions, QRS of supraventricular derivation, heart rate more than 150 pulses per minute [8]. Some special characters found in the PSVT. One of these is the Heart beat of the re-entrant tachycardia which is about 150 to 200 bpm, P wave present in QRS composite or brusquely after the ORS composite [9]. The heart beat of the AV re-entrant tachycardia was recorded to be 150-250 bpm. The situation of the QRS was tapered at the time of orthodromic transference and spacious in case of antidromic transference [10, 11]. Sinus tempo can be reestablished by the intensification of vagal attitude by blue-collar carotid inspiration and some other vagal drills. Adenosine dosage of 6-12 mg was considered as the initial protector. Another protector antibiotic was verapamil which was used 5 to 15 mg [7]. Both the antibiotics were found successful in about 90% cases [3]. AV node can be barren rapidly by a comparative of adenosine called purinergic. But when it is governed intravenously wedge PSVT counting the AV node as part of re-entrant path [6]. Verapamil is a calcium conduit. It acts oppositely, it lessens the speed of AV transference and ceases AVNRT [12].

Both adenosine and verapamil can be used for the treatment. But adenosine is more widely used because its speed of achievement is very fast and having lesser half-life [13]. The faster rates of achievements of adenosine were noticed in PSVT cases. On the other hand, the achievements of verapamil in PSVT patients enhanced with lesser rates [14]. We can choose one of the drugs either adenosine or verapamil by considering the variations

#### **SUBJECTS AND METHODS:**

The investigational study was arranged in Jinnah Hospital, Lahore (August 2017 to March 2018). In the study, about 200 cases suffering from severe PSVT were added in the study. The age of the patients was greater than 15 years. There was a total of 200 patients added in the study. The patients were divided into two groups. Both groups A and B contains 100 patients. Some patients like PSVT removed by vagal manoeuvres, the previous record of bronchospasm, greater QRS composite tachycardia, initially suffered from sinus syndrome or heart wedge and patients undergoing heart transplantation. Complete information about patients like narration, corporeal determination, and permission for the usage of antibiotics was taken from their relatives. ECG of both the groups was examined and performed.

Adenosine 6 mg with 20 ml saline flushes to group A was given through three various pathways with the addition of 12 mg 4 boluses with saline flush if the success was not obtained by the ingestion of the first bolus.

5 mg successive dose of verapamil was given to the patients with a break of 5 minutes. This process was carried out continuously until success was achieved. From the injection of antibiotics to the patients and the time when the response occur were noted in both groups. After the 15 minutes of cure ECG was performed on patients to check out the details.

Patients present in group B were also given the adenosine. This was done when the verapamil did not show any special successive rates and vice versa for group B. Data was recorded on papers. Data was entered and detected by SPSS 10 version.

#### **RESULTS:**

Total of 200 cases suffering from severe PSVT was added in the experimentation. The average age of the patients was about 41.06 years. Out of these patients, the number of women was 53.5% and a number of men were 46.5%. Both groups were almost similar. In group A the percentage of men was 58%. In group B the percentage of men was 49%. The average age of the patients in group A was  $(42.8 \pm 13.9)$  years. And the average age recorded in group B patients was about  $(40.5 \pm 15)$  years.

The main drawback observed in most of the patients was shudder. Shudder has recorded in 94% of patients of group A and 100% patients of group B.

Some other common symbols of the disease are squatness of inhalation and plummeting of heart. Very less plummeting of heart and vertigo were also noticed in some cases.

Variables	Patients characteristics	Adenosine	Verapamil	P value
	15-35	33%	46%	
	36-55	46%	37%	
	56-75	21%	16%	
Age	>76	-	01%	
	Mean	$42.8\pm139$	$40.5 \pm 15.1\%$	
	Males	58%	49%	
Sex	Females	42%	51%	P value= $0.26$
	Palpitation and dyspnea	55%	58%	
	Palpitation and sinking	30%	4206	
Clinical	of the heart	3970	42.70	
presentation	Sinking of heart	05%	0%	P value insignificant
	Vertigo	01%	0%	
Mean heart rate		101.3 bpm	179.5 hpm	
before treatment		191.5 opin	179.5 opin	
Short RP		75	73	
Long RP		07	12	
Prexcitation	Absent	99%	96%	P value $= 0.17$
Preexcitation	Present	01%	04%	

Table – I: Baseline characteristics of two groups



Doses	Group A	Group B
1 <sup>ST</sup> bolus	74%	68%
2 <sup>nd</sup> bolus	22%	24%
3 <sup>rd</sup> bolus	0%	3%
Total	96%	95%

 Table – II: Outcome after successive bolus doses in both groups



Syncope was not observed in any of our patients. A hemodynamic compromise was also not identified in any of our cases. With the ingestion of 2 regular doses of 6 mg and 12 mg adenosine, 96% achievement was observed in group A. From total of 100 cases 74% achievement was observed by the use of only first bolus. The second bolus was given to only those 26% patients in which success was not achieved by the first attempt. These were easily reverted by the use of the second bolus.

It was observed that all women can be easily cured by only a single bolus. Most of the men received the second bolus to cure themselves completely. In group A only 5 males were given the second bolus. The rate of reversion of sinus rhythm was more successful in women as compared to men. The achievement rate was 90.47% in males and 62.06% in females. The restoration rate was 100% in women and 81.18% in men in the case when the second bolus was tried. There were only 4 cases in which complete treatment was not observed even after the implementation of both boluses. These patients were then given the verapamil and all of them were safe and healthy completely.

Achievement rate was 95% in group B. Three

successive doses of verapamil were given to each patient. 68% success was gained by the use of the first dose of 5 mg verapamil. Left of the patients was followed by the second bolus of verapamil. 24 out of the remaining 32 patients received the restoration. 8 patients did not show restoration. These include only one female and 4 male who did not revert themselves completely even after the regular injection o f3 doses of verapamil. These 5 patients than received adenosine instead of verapamil. The restoration was found in these 4 patients. But still, deterioration did not occur in one case. He was an exceptional case of Ebstein incongruity and WPW syndrome. He was then given fourth bolus and he received the setback at last.

The time was recorded from the initiation of use of antibiotics first time until the success rate was obtained. More rapid consequences were found by the use of adenosine. The time difference was  $2.1 \pm 2.7$  minutes in group A and  $5.1 \pm 3.5$  minutes in group B. It was also recorded that rapid and more setback was found in group A as compared to group A. it can be shown by the observation that adenosine is more successful than verapamil.

ECG and average heartbeat of the patients prior to

and following the treatment were evaluated. The heart rate of the patients of both groups was determined. The average rate of group A patient was  $191.3 \pm 57.1$ . And in group B the average heart rate was  $179.5 \pm 30$ . No apparent P wave was examined during tachycardia. In these types of patients, RP intermissions should be measured. Left patients were observed to have petite RP intermissions. The example of lesser RP intermissions was 91.47% in group A and 85.89% in group B. Other patients contain greater RP values. After the treatment, the heartbeat was about  $93.6 \pm 10.8$  in group A and 85.3% in group B.

### **DISCUSSION:**

PSVT assaults can be removed or hindered by various non-pharmacological methods. If they cannot be removed or cured by simple processes that we use biological tools like adenosine and verapamil. Both can protect patients from severe PSVT. The selection of adenosine or verapamil depends on the selection of doctor, prices and most wanted commencement of the function. There are only a few chances that patients become hemodynamically concessional and need cardioversion [12].

In this observation adenosine and verapamil were compared with each other and the result was that the adenosine is more effective in reversion of PSVT as compared to verapamil.

In the removal of severe PSVT, both antibiotics were found to be effective. But the action of adenosine in fast and spontaneous in comparison with verapamil. It was observed in those study that adenosine is effective in its two boluses but 3 boluses of verapamil were required to remove the disorder.

Two successive doses of adenosine are affected to terminate the PSVT in group A up to 96%. Similar observations were made in case of verapamil. Our study was also similar to some previous observations and studies in which both the adenosine and verapamil both were found to be effective in removing the PSVT [3, 17].

In group A the average age of the patients was  $42.8 \pm 13.9$  years. The average age of the group B patients was  $40.5 \pm 15$  years. The ages were almost similar in both the groups. In recent generations, PSVT was found to attack more profusely. It was noticed that older people suffering from this disease were found more in number in clinics and emergency wards than the younger ones. The average age of the PSVT patients was  $54 \pm 17$  according to study reports [18].

It was noticed that in group A most of the patients achieved success in their treatment by a single bolus of adenosine 6 mg. A study organized by Davis and his companion showed that 65% achievement rate in PSVT was observed by using single bolus of adenosine. These consequences were similar to our study [10]. 85% achievement rate was also analyzed in another study [19].

The 26 patients who were not reverted by single bolus were reverted to about 84.61% by the second bolus. The reaction in SVT with the injection of various boluses of adenosine was also observed in one study. The adenosine dosage and reactions produced were 3 mg, 6mg, 12 mg and the responses of termination were 35.2%, 62.3%, 80.2% and 91.4% correspondingly [20].

Out of 100 Patients, 96% showed complete success but 4 did not revert. These did not show any reversion even after the injection of the second bolus. These were then subjected to verapamil. 5 mg verapamil showed a notable reaction. 81.3% achievement rate was identified in cases where adenosine failed to respond [21].

Adenosine was found to be more effective in females as compared to males. The rate of success in females was 100% as compared to males 92% in our study. In prior studies, the same consequences were found. 5 mg boluses of verapamil were needed to revert the PSVT in 68%, 75% and 37.5% cases accordingly. By the use of 5 regular boluses of verapamil 95% achievement rate was seen in patients. This was almost similar to the consequences of adenosine in group A which was 96%. 5 patients who did not show a reaction against verapamil were treated with adenosine and 4of which showed a success.

One study compared the effectiveness of adenosine and verapamil in severe PSVT. 62 patients were subjected to verapamil with 5 mg boluses on a regular interval of time for the hindrance of SVT [17]. There was no significant variation between the two drugs were seen. The success rate was 80.6% in the case of adenosine and 87.1% in the case of verapamil [17]. Same consequences were identified in the experiment conducted by Ballo P et al. [22].

Adenosine was found to be more rapid and efficient in our study. Average time of reversion was  $2.1 \pm 2.7$ minutes in group A and  $5.1 \pm 3.5$  minutes in group B. The variation in time was found because of the rapid function of adenosine. Adenosine is fast and more effective as compare to verapamil which was slow and takes more duration of time to perform its action. It was observed that in group A 26% of cases showed an immediate success. Only 1% of patients in group B reverted at a spontaneous time. The time of action of adenosine was lower than verapamil.

In PSVT fast heartbeat showed shudder and squatness of inhalation. The average heartbeat before treatment was  $191.3 \pm 57.1$  in group A and  $179.5 \pm 30$  in group B. The average rate of heartbeat after setback was  $93.6 \pm 10.8$  in group A and 85.3 in group B. Another study recorded heartbeat before setback as  $171 \pm 21$  in every minute [23]. After the setback, the heart rate was decreased in both groups with similar rates. The reduction in heart rate was 120 bpm in group A and 105 bpm in group B. More reduction in heartbeat was recorded in verapamil in comparison with adenosine. This was because of the longer duration of action of verapamil [24].

By ECG a P wave retrograde was assessed in PSVT. This showed that in most of the cases lesser RP hiatus was seen. It was 191.47% in group A and 85.89% in group B. Lesser RP means that most of the patients had AVNRT and AVRT as a method of PSVT. It was also noticed by Porter M J and companions that severe PSVT followed by AVRT was the main method. The choice of patients was performed irregularly. The patients already suffered from PSVT attacks were taken in group B. These patients were smaller in number. Because of that reason, there is less jeopardy of assortment prejudice in our experimentation.

The duration of time from the injection of antibiotic to its response was calculated. The time duration of the response was much lesser in adenosine than verapamil. But it is difficult to calculate the duration of time in tragedy room. So, the time is usually calculated in minutes. The expenditures of adenosine were not analyzed in our study.

#### **CONCLUSION:**

Adenosine is found to be more effective in the execution of PSVT. The adenosine and verapamil are almost similar to each other. The difference lies just in the mechanism of action of adenosine which is more rapid than verapamil.

#### **REFRENCES:**

- Martius H. Uber dies Behan lung von Blasenscheidenfisteln, insbesondere met Hilfe einer Lappenplastik. Geburtshilfe Gynakol 1932; 103:22-34 7.
- Kirchhoff JD, Wechsler M, Romas NA. Urinary fistulas: vaginal repair using a labial fat pad. J Urol 1977; 117:57-59.

- Wall LL, Karshima JA, Kirschner C, Arrowsmith SD. The obstetric vesicovaginal fistula: characteristics of 899 patients from Jos, Nigeria. Am J Obstet Gynecol 2004; 190: 1011-9.
- Evans DH, Madjar S, Politano VA, Bejany DE, Lynne CM, Gousse AE. Interposition flaps in transabdominal vesicovaginal fistula repairs: are they really necessary? Urology 2001 Apr;57(4):670-4.
- 5. Uprety D, Babu S, Sharma M, Jha M. Obstetric genitourinary fistula in the eastern part of Nepal. Asian J Obstet Gynecol 2001; 6: 36-8. 12.
- 6. Tariq M, Ahmed I, Arif M, Ali MA, Khan MI. Comparison of Out Come of Vesico-Vaginal Fistula Repair with and Without Omental Patch. Ann Pak Inst Med Sci 2011; 7(3): 123-6.
- O'Conor VJ Jr. Review of experience with vesicovaginal fistula repair. J Urol 1980; 123:367-9
- Uprety DK, Subedi S, Budhathoki B, Regime MC. Vesicovaginal fistula at a tertiary care center in eastern. Nepal. J Nepal Med Assoc. 2008; 47(171):120-2
- Nawaz H, Khan M, Teareen FM, Khan S. Retrospective study of 213 cases of female urogenital fistula. J Pak 190: 1011-9 Med Assoc 2010; 60(1):28-32 22.
- Davies RJ, Miranda S. Conservative treatment of vesicovaginal fistulas by bladder drainage alone. Br J Urol 1992; 70:339
- 11. Stovsky MD, Ignatoff JM, Blum MD. Use of electro coagulation in the treatment of vesicovaginal fistulas. J Urol 1994; 152(pt 1):1443-4
- Morita T, Tokue A. Successful endoscopic closure of radiation-induced vesicovaginal fistula with fibrin glue bovine collagen. J Urol 1999; 162:168-9
- Dogra PN, Nabi G. Laser welding of vesicovaginal fistula. Int Urogynecol J.2001; 12:69-70
- 14. Blaivas JG, Heritz DM, Romanzi LJ. Early versus late repair of vesicovaginal fistulae--a tenyear repair of vesicovaginal fistulas: vaginal and abdominal approaches. J Urol 1995; 153:111-2
- 15. Wein AJ, Malloy TR, Carpiniello VL. Repair of the vesicovaginal fistula by a suprapubic transvesical approach. Surg Gynecol Obstet 1980; 150:57-60 16.
- Robertson JR. Vesicovaginal fistulas. In: Slate WG, editor. Disorders of the female urethra and urinary incontinence. Baltimore: Williams & Wilkins; 1982. P. 2429.
- 17. Keetel WC. Vesicovaginal and urethrovaginal fistulas. In: Buchsbaum HJ, Schmidt JD, editors.

Gynecologic and obstetric urology. Philadelphia: WB Saunders; 2004; 190: 1011-9 1982. P.318-26

- Latzko W. Postoperative vesicovaginal fistulas: genesis and therapy. Am J Surg 1942; 58:211-28?
- Langkilde NC, Pless TK, Lundbeck F, Nerstrøm B. Surgical repair of vesicovaginal fistulae--a ten-year retrospective study. Scand J Urol Nephrol 1999 Apr; 33(2): 100-3.
- 20. Ray IB. Narrow complex tachycardia: recognition and management in the emergency room. J Assoc Physicians India 2004; 52: 816-24.
- 21. Steinbeck G, Reichmann C. Paroxysmal supraventricular tachycardia: preferential use of either adenosine or verapamil.
- 22. Wall LL, Karshima JA, Kirschner C, Arrowsmith SD. The obstetric vesicovaginal fistula: characteristics of 899 patients from Jos, Nigeria. Am J Obstet Gynecol and Obstet Gynecol 2004; 190: 1011-9.
- 23. Rasool M, Mumtaz F, Tabassum SA. Vesicovaginalv fistulas: fistula repair, urologist's experience at Bahawalpur, Professional Med J 2006; 13(3):445-52 19.
- 24. Jalbani MH, Deenari RA, Sheikh JM. Vesico-Vaginal Fistula- Experience of Surgical Repair at Larkana. Pak J Med Res, 2006;45:3:156-9.