CODEN [USA]: IAJPBB ISSN: 2349-7750

INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES
http://doi.org/10.5281/zenodo.3830141

Available online at: http://www.iajps.com Research Article

# USE OF AN ANTHROPOMETRIC INDICATOR TO RECOGNIZE HYPERTENSION IN YOUNG PEOPLE STUDYING IN BAHAWALPUR, PAKISTAN

**[1]Dr. Umar Farooq, [2]Dr. Aleena Sabir, [3]Dr. Kinza Ayesha**
[1]BVH Bahawalpur.

**Article Received:** March 2020 **Accepted:** April 2020 **Published:** May 2020

**Abstract:**
*Our current cross-sectional survey was led to decide on presumed intensity of anthropometric markers and to prescribe cut-off points to separate hypertension in young people in Bahawalpur, Pakistan. The overall 2,467 persons with ages 13-18 years joined in the survey, which represents an average duration of 15.6±2.52 years. Each anthropometric marker had a critical area below the ROC elbow, with a weight file (BMI) and an abdominal contour reaching from 0.8 to 0.9. The best anthropometric gauges to predict hypertension in young men remained waist circumference, BMI, and abdomen to body weight ratio. For young women, BMI remained best indicator, trailed by body weight and waist circumference. The suggested BMI cut-off for young men was 21 kg/m2 and 21.8 kg/m2 for young women. For WC, suggested cut-off was 68.2 cm for young men and 69.3 cm for young women. The BMI and WC thresholds were prescribed for use in the school setting, where the estimate can be oriented without stretching it too much.*
***Place and duration:*** *the research was conducted in BVH Bahawalpur for one-year duration from Jan 2019 to December 2019.*
**Keywords:** *Anthropometric Pointer, Hypertension, Young People.*

**Corresponding author:**
**Dr. Umar Farooq,**
*BVH Bahawalpur.*

QR code

*Please cite this article in press Umar Farooq et al,* ***Use Of An Anthropometric Indicator To Recognize Hypertension In Young People Studying In Bahawalpur, Pakistan.,*** *Indo Am. J. P. Sci, 2020; 07(05).*

## INTRODUCTION:

Pre-adult overweight has become a growing concern for well-being, a scourge that influences both creative and created nations. Occurrence has reached a level that warrants rapid reflection on the essential and optional anticipation of overweight and weight in youth and adolescents. It is during the period of youth and adolescence that they construct their conception of food and movement that can influence their lifestyle as adults [1]. With the current health shift that includes access to fast foods, soft drinks, stationary lifestyles, physical latency, and the progressive use of innovation-related devices, many youths aged 11 to 20 are considered less energetic also eat more, subsequent in increased BMI and fat [2]. These unfortunate trends have paid to expansion of co-morbidities just like increased blood cholesterol, DM-type 2 and hypertension. Numerous studies have linked hypertension to overweight and corpulence in young people [3]. Despite the fact that the accumulated confirmations have a mixed ending on the connection between hypertension and muscle/fat ratio, estimation of muscle/fat ratio using anthropometric markers has proven to be a powerful methodology for anticipating hypertension, especially in surveys based on a large population and network. The National Morbidity and Health Survey of Pakistan 2013 reported that 45.7% of Pakistan grownup people was inclined by hypertension, an expanded pattern compared to previous years (44.8% rendering to NHMS III 2007 also 34.9% rendering to NMHS II 1997) [4]. Those results demonstrated position of controlling also avoiding hypertension at a younger age. The use of anthropometric markers as a screening instrument in a roundabout way to distinguish high circulatory pressure at school. Since the technique does not include the use of a confusing device and the contribution of particular abilities, the estimation can be done systematically by school specialists. If, by chance, the anthropometric pointers show variations from the normal results, a more thorough pulse assessment can be performed and the managers can be referred to an emergency clinic. Therefore, the purpose of this review was to decide on ubiquity of hypertension and the intensity of preciousness of the anthropometric markers and to prescribe cut-off points to separate hypertension in young people [5].

## METHODOLOGY:

Our current research was the cross-sectional survey of circulatory strain profiling among youth in Bahawalpur, Pakistan. The review remained supported by Basic Research Fellowship Programme of the Pakistan Ministry of Higher Education. Sarawak is Pakistan's main province, positioned on the island of Borneo. According to the 2015 assessment, its population is expected to reach 3,638,500, with more than 45 sub-ethnic gatherings, each with its own language, culture and way of life. The seven key ethnic sets are Iban, Chinese, Malay, Bidayuh, Melanau and Orang Ulu. The examination population included 186 under students from 14-18 year old auxiliary schools in Sarawak, with an enrolment of 205,150 under students, obtained from the Ministry of Education (March 2015). Using the recipe for example size for limited populations [11] s= [X2Np(1-p)]÷[d2(N-1)+ X2p(1-p)], where s is required example size, X is the Z score for 98% provisional certainty (3.59), N is the population size, p is the range of the population (estimated at 0.6 as this could give most dangerous example size), d is accuracy level or margin of error (0.029), and the least required example size was 2200. A share of 19 schools was chosen for each state and deliberate consideration was used to determine schools by size of enrolment, as was stratification by urban area and country. The measures for inclusion were cases with no physical or intellectual debility, no pre-existing hypertension, or no conditions that would cause ancillary hypertension. Moral agreement was obtained from the Medical and Ethical Committee of the University of Pakistan at Sarawak and the Ministry of Education, Pakistan. Informed consent remained gained from guardians/parents of the respondents prior to the day of the information package. An examination form was given to each respondent and their respondents/guardians. In addition, the respondents were informed of the examination data upon arrival of the information package. Cases were assured of privacy of its information and reserved privilege of withdrawing from exam at any time.

## RESULTS:

The overall of 2550 people responded to the survey and agreed to take an interest in it now. However, 82 (4.3%) were banned because they missed school when the information package arrived and a number declined regardless of parental agreement, so the size of the last example was 2,468 (95.7% response rate). There were more young women than young men (59 per cent versus 41 per cent), with an average duration of 15.6±2.60 years, with a greater proportion coming from Iban and Malay ethnic gatherings. Average weight, stature and abdominal circuitry remained higher in young men than in young women, while BMI and GTR were higher in young women. In any case, contrasts in sexual orientation were found only in terms of weight, stature and body mass index. Both sexual orientations have a comparable Coindex. More young males than young females were considered prehypertensive and hypertensive, and this distinction

was enormous. More young women were considered overweight than young men; at least in the overweight class, the reverse was true. BMI showed an upward trend between defendants, particularly in the 14+ age group. Midline circumference also SBP and DBP circulatory pressure were also used as comparative examples, but were not reliably extended to more established age sets (see Table 2). Table 2 and Figures 1 and 2 show the latest models using Stepwise relapse testing and CAR surveys. Altogether anthropometric pointers displayed a critical relationship with hypertension for calculated relapse and CAR curves, with BMI and WC taking precedence over a greater area and reasonable appraisals (0.8-0.9). Founded on the results of thresholds, affectability, and particularities, the preferred anthropometric marker for predicting hypertension in young men remained WC and BMI, trailed by TMI. Though, for young women, BMI remained best predictor, trailed through WHtR and WC.

**Table 1: Sociodemographic and health profile amongst man and woman cases (N=2466).**

| Variables | Mean+ Standard Deviation | | P value |
|---|---|---|---|
| | Females N= 1430 | Males N= 1035 | |
| Age | 14.5 ± 1.51 | 14.4 ± 1.48 | 0.06 |
| Weight (kg) | 52.1 ± 13.81 | 57.6 ± 15.79 | <0.002** |
| Height (m) | 1.5 ± 0.06 | 1.6 ± 0.08 | <0.002** |
| BMI (kg/m2) | 21.6 ± 4.78 | 22.4 ± 5.74 | 0.09 |
| Blood pressure | | | <0.002** |
| Pre-hypertension | 185 (12.9) | 235 (23.6) | |
| Hypertension | 199 (19.3) | 125 (8.8) | |

**Table 2: Anthropometric measurement and blood pressure conferring to age between male also female cases (N=2465).**

| | Gender | 13 years | 14 years | 15 years | 16 years | 17 years | 18 years | P value |
|---|---|---|---|---|---|---|---|---|
| BMI | F | 21.96±4.86* | 22.76±5.06* | 20.86±4.10 | 22.32±4.56* | 20.55±4.27* | 21.38±5.62 | <0.02 |
| | M | 21.57±4.21 | 21.78±4.46 | 21.09±5.30 | 21.85±3.89 | 20.80±4.97 | 20.96±5.03 | 0.162 |
| WC | F | 68.58±8.62 | 72.04±10.70 | 70.75±9.82 | 67.68±9.20* | 73.73±49.48* | 70.21±9.71 | 0.008 |
| | M | 70.42±13.30 | 70.66±12.08 | 72.81±10.04 | 72.92±10.34 | 69.77±12.59 | 71.80±10.08 | 0.046 |
| WHt | F | 0.46±0.07 | 0.45±0.05* | 0.49±0.31* | 0.46±0.07 | 0.45±0.06* | 0.46±0.06 | 0.02 |
| | M | 0.44±0.06 | 0.44±0.06 | 0.44±0.06 | 0.46±0.09 | 0.45±0.07 | 0.44±0.07 | 0.071 |
| DBP | F | 111.69±13.11 | 109.39±11.23 | 112.29±12.58* | 112.41±14.30 * | 109.93±11.40 | 111.79±10.81 | 0.014 |
| | M | 120.54±13.90 | 119.34±13.09 | 112.35±13.18* | 120.59±12.9 | 116.23±13.77 * | 118.66±14.36 | <0.02 |
| SBP | F | 67.04±9.90* | 66.40±7.50 | 65.66±9.90 | 66.84±8.87 | 64.62±8.61* | 65.25±9.02 | 0.012 |
| | M | 66.92±9.58* | 65.17±7.88 | 65.15±10.23 | 63.89±9.50* | 63.12±10.19* | 64.73±9.27 | 0.008 |



**Figure 1: ROC curves that associate 4 anthropometric pointers of obesity as discriminators of hypertension. BMI: in centimeter-waist circumference; WHR: C index: conicity index.**

## DISCUSSION:

The occurrence of hypertension is currently 23.6% in men and 13.6% in women. With regard to prehypertension, the rate is slightly lower, at 21.4% for men and 9.7% for women. The current outcomes showed a higher banality rate connected to other similar surveys, showing that, in a limited ability to focus on five years, occurrence of prehypertension and hypertension enlarged despite the fact that the area of examination was extraordinary and yet it was an agent of Pakistan [6]. Added result of our current survey further showed that the banality of overweight and corpulence was also high in men (overweight 12.8%, corpulence 14.8%) and women (overweight 13.8%, corpulence 11.8%), which can be firmly identified with the high prevalence of hypertension. Such a perception strongly recommends early consideration of how to treat this medical problem that can lead to hypertension and weight-related wellness complexity in later life [7]. One of the motivations for this review was to decide on the power of prescience and to establish cut-off points for anthropometric pointers for the prediction of hypertension [8]. As a result of this discovery, a standard well-being test using anthropometric estimation at the school level can be routinely performed to screen and identify young people with signs of hypertension for further investigation in a clinical setting. Of all the anthropometric markers, BMI and waist circumference had the highest affectivity and self-esteem, confirming the evidence from past examinations that both markers were identified in the same way, with hypertension being independent of age and sexual orientation [9]. Nevertheless, in one investigation, such findings were challenged when WC estimated focal weight was considered a more reliable indicator of hypertension. This is established by the huge population concentration in China wherever conclusions reliant on on 505,200 grown-ups described that focal weight was key gauge. In pre-adult examinations, confirmation of anthropometric markers of hypertension in youth and adolescents is uncertain. A few examinations showed that BMI and waist circumference were critical indicators of hypertension with ORs of 3.70 and 2.95, individually. While one review found that gtr and WC are preferred indicators of hypertension over BMI, another review would prohibit gtr and maintain BMI and WC [10].

## CONCLUSION:

In total, grounded on results, this is prescribed to apply the BMI and WC markers to identify hypertension in young people in Sarawak. A cut-off points of 24 kg/m2 for young men and 21.8 kg/m2 for young women was prescribed for BMI, and a cut-off point of 68.2 cm for young men and 69.3 cm for young women was prescribed for WC to identify high circulatory pressure. Both markers might be applied at school,

where a scale, stadiometer and tape measure can be accessed without stretching too much. Recognized adolescents can also be referred for clinical examination and thus benefit from treatment and intervention.

**REFERENCES:**


1. Taiwo, Emmanuel O., and Olusoga A. Sofola. "Baseline anthropometric measurements and obesity of people working in Isale-Oko, Sagamu, Ogun State, Southwest, Nigeria." *The Nigerian Journal of General Practice* 18, no. 1 (2020): 5.
2. Wangui, Anthony Muchiri, Maximilla N. Wanzala, Micky O. Olutende, and Edwin K. Wamukoya. "Assessment Of Pre-Season Anthropometric Tests Of Rugby Players In A Championship Club, Kenyan Rugby Union." *European Journal of Physical Education and Sport Science* (2019).
3. Kim, Tae Jun, Hyuk Lee, Sun-Young Baek, Kyunga Kim, Yang Won Min, Byung-Hoon Min, Jun Haeng Lee, Hee Jung Son, Poong-Lyul Rhee, and Jae J. Kim. "Metabolically Healthy Obesity and the Risk of Erosive Esophagitis: A Cohort Study." *Clinical and translational gastroenterology* 10, no. 9 (2019).
4. Kim, Tae Jun, Hyuk Lee, Sun-Young Baek, Kyunga Kim, Yang Won Min, Byung-Hoon Min, Jun Haeng Lee, Hee Jung Son, Poong-Lyul Rhee, and Jae J. Kim. "Metabolically Healthy Obesity and the Risk of Erosive Esophagitis: A Cohort Study." *Clinical and translational gastroenterology* 10, no. 9 (2019).
5. Ijoma, Uchenna Nkemdilim, Peter Chime, Chinwe Onyekonwu, Birinus Adikaibe Ezeala-Adikaibe, Casmir Orjioke, Obumneme Benneth Anyim, Obinna Donatus Onodugo et al. "Factors Associated with Overweight and Obesity in an Urban Area of South East Nigeria." *Food and Nutrition Sciences* 10, no. 07 (2019): 735.
6. Maine, Andrew, Michael J. Brown, Chantal F. Ski, David R. Thompson, Lynne Marsh, and Lisa O'Leary. "Recruitment settings, delivery contexts, intervention techniques and outcomes of health promotion programmes for young adults with intellectual and developmental disabilities: A systematic review." *Research in Developmental Disabilities* 99 (2020): 103592.
7. Ongaga, Jared Kebaso. "Assessing Treatment Adherence In Hypertension Control Among Hypertensive Patients Post Health Education At Kenyatta National Hospital." PhD diss., University of Nairobi, 2019.
8. Al Muktadir, Mohammad Hamid, Md Ashraful Islam, Mohammad Nurul Amin, Supriya Ghosh, Shafayet Ahmed Siddiqui, Dipti Debnath, Md Monirul Islam, Tufael Ahmed, and Farhana Sultana. "Nutrition transition–Pattern IV: Leads Bangladeshi youth to the increasing prevalence of overweight and obesity." *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* 13, no. 3 (2019): 1943-1947.
9. Hassan, Nayera E., Abeer A. El Ashmawi, Sahar A. El-Masry, Waheba A. Zarouk, Marwa F. Mira, Gamila SM El-Saeed, and Omar H. Dwidar. "Metabolic syndrome in a sample of Egyptian adolescent girls and its association with apolipoprotein E." *Journal of paediatrics and child health* 55, no. 11 (2019): 1344-1350.
10. Hazarika, Himangshu Jyoti, Aziz Ahmed, and Kaushal K. Chandrul. "Miracle of Herbals and Natural Products in Treatment and Regulation of Obesity." (2019).