



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF  
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3555312>Available online at: <http://www.iajps.com>

Research Article

**RAISED HOMOCYSTEINE LEVEL IN CASES WITH  
ISCHEMIC STROKE**<sup>1</sup>Anam Batool, <sup>2</sup>Rahat Naveed, <sup>3</sup>Asma Naseem

Article Received: September 2019 Accepted: October 2019 Published: November 2019

**Abstract:**

*Stroke remains one of the most devastating of all neurological diseases, often causing death or gross physical impairment or disability. Homocysteine is one of the modifiable risk factor regarding this.*

***Objectives:** To determine the frequency of raised homocysteine level in patients of ischemic stroke.*

***Material and methods:** This cross sectional study was done at Services Hospital, Lahore and THQ Kotmomim from the period of July 2018 to December 2018 in which 146 cases with age range of 40 to 70 years of both gender suffering from ischemic stroke were included and were assessed for their homocysteine level*

***Results:** In this study there were 146 cases out of which 76 (52.05%) were males and 70 (47.95%) females. The mean age was  $56.25 \pm 7.67$  years and mean BMI was  $29.52 \pm 3.42$ . There were 40 (27.40%) cases with DM, 48 (32.88%) hypertensive, 46 (31.51%) had history of smoking while 18 (12.33%) had family history of ischemic stroke. Raised homocysteine level was seen in 58 (33.70%) cases. Raised homocysteine level was more seen in females where it was seen in 32 (45.71%) out of 70 females as compared to 26 (34.21%) out of 76 males with  $p$  values of 0.32. This raised level was also seen maximum in age groups of 60 to 70 years affecting 28 (45.14%) out of 62 cases ( $p=0.43$ ). There was no significant association with any of the risk factors with raised homocysteine level.*

***Conclusion:** Ischemic stroke is a fatal condition. Raised homocysteine level is seen in every 3<sup>rd</sup> cases of ischemic stroke. Hypertension and family history of ischemic stroke has almost significant association with it.*

**Key words:** Stroke, homocysteine

**Corresponding author:**

Anam Batool,

QR code



Please cite this article in press Anam Batool et al., **Raised Homocysteine Level In Cases With Ischemic Stroke**, *Indo Am. J. P. Sci.*, 2019; 06(11).

**INTRODUCTION:**

Stroke remains one of the most devastating of all neurological diseases, often causing death or gross physical impairment or disability. Worldwide, Asia, Russia and Eastern Europe experience the highest rates of mortality and disability and loss of quality of life because of stroke. [1-2]

Atherosclerosis, diabetes mellitus (DM), hypertension (HTN), ischemic heart disease (IHD), peripheral arterial diseases are the known risk factors for ischemic stroke. But they are unavoidable and non-modifiable risk factors. It is seen that a good number of cases, these well studied risk factors are even absent too.[3] This opens the horizon to look for new or undiscovered and if possible, the modifiable risk factors associated with stroke. Homocysteine (Hcy) excess is one of those risk factors, which are modifiable and recently work is going on globally to look for its importance and their role in ischemic stroke.

The raised level of homocysteine occurs due to deviation in the metabolism of Methionine, which is an essential amino acid; leading to hyper coagulable state which cause clotting in blood vessels and lead to ischemic infarcts in brain. It is affected by many factors like age, vitamin B6, B12, Folic acid and intake of certain drugs like steroids and cyclosporine.

Han L et al revealed that the raised levels of homocysteine were not only directly associated with increased risk of stroke but also with the severity of the ischemic stroke and folic acid was found useful for its prevention.<sup>4</sup> According to a study by Ashjazed N et al carried on 171 cases of ischemic stroke as compared to 82 matched controls, the homocysteine level was found significantly raised in cases with ischemic stroke as compared to controls. The mean fasting Hcy levels was significantly higher in the cases (16.2  $\mu\text{mol/L}$ , 95% CI: 14.8 to 17.5) than in the controls (13.5  $\mu\text{mol/L}$ , 95% CI: 12.4 to 14.6) (P=0.013) [5]

According to a study by Sadiq M et al it was seen that raised levels of homocysteine was seen in 58.3% of cases in their study at Karachi.<sup>6</sup> Another study from Bangladesh, done by Rehman et al on 36 patients showed that raised homocysteine levels were seen in 75% of the cases.[7]

**Objective:**

The objective of this study was to determine the frequency of raised homocysteine level in patients of ischemic stroke in people of Rahim Yar Khan.

**MATERIALS AND METHODS:**

This cross sectional study was done at Services Hospital, Lahore and THQ Kotmomin from the period of July 2018 to December 2018 in which 146 cases of ischemic stroke were included with following inclusion and exclusion criteria.

**Inclusion criteria.** Gender: both male and female

- Age: 40 – 70 years
- All the patient with ischemic stroke (as per operational definition) presenting within 1 week of the symptoms.

**Exclusion criteria**

- Patients with hemorrhagic stroke (hyperdense area on CT brain plain in at least 3 slices)
- Patients with any history of prior stroke in life time.
- Patients taking drugs like multivitamin/steroids/cyclosporine within last two weeks

**Hemorrhagic Stroke:**

The patients were labeled as having hemorrhagic stroke, when he or she had a history of severe headache and sudden loss of movement in one or more than one parts of the body. The diagnosis was confirmed on CT scan brain (hyper dense area in the brain in at least 3 slices).

**Hyperlipidemia:**

Hyperlipidemia was labeled 'yes' when there was any of the following lipid values observed:

1. High serum LDL-C levels: >100mg/dl
2. Hypertriglyceridemia = serum triglycerides level: >150mg/dl
3. Hypercholesterolemia= Cholesterol level >200 mg/dL
4. Low HDL: <40 mg/dL

**Statistical Analysis:**

In this cross sectional study, all the collected data were entered into SPSS version 21 and analyzed. The qualitative data like sex, DM, HTN, Smoking, family history of ischemic stroke, raised level of homocysteine (yes/no) were assessed as frequency and percentages. Quantitative data like age, weight, height, BMI, homocysteine level ( $\mu\text{mol/L}$ ) and duration of symptoms (days) were assessed as means and standard deviations. Post stratification chi square test was applied and p value  $\leq 0.05$  was considered as significant.

**RESULTS:**

In this study there were 146 cases out of which 76 (52.05%) were males and 70 (47.95%) females. The

mean age was  $56.25 \pm 7.67$  years and mean BMI was  $29.52 \pm 3.42$ . There were 40 (27.40%) cases with DM, 48 (32.88%) hypertensive, 46 (31.51%) had history of smoking while 18 (12.33%) had family history of ischemic stroke. Raised homocysteine level was seen in 58 (33.70%) cases. Raised homocysteine level was more seen in females where it was seen in 32 (45.71%) out of 70 females as compared to 26

(34.21%) out of 76 males with p values of 0.32 as shown in table 1. This raised level was also seen maximum in age groups of 60 to 70 years affecting 28 (45.14%) out of 62 cases followed by 40 to 49 year in 16 (44.44%) out of 36 cases with p values of 0.43 as in table 2. There was no significant association with any of the risk factors with raised homocysteine level as in table 3.

**TABLE 01: RAISED HOMOCYSTEINE LEVEL WITH RESPECT TO GENDER n= 146**

GENDER	RAISED HOMOCYSTEINE LEVEL		Total
	Yes	No	
MALE	26 (34.21%)	50 (65.79%)	76 (100%)
FEMALE	32 (45.71%)	38 (54.29%)	70 (100%)
<b>Total</b>	<b>58 (69.8%)</b>	<b>88 (30.2%)</b>	<b>146 (100%)</b>

Chi square= 1.007, p= 0.32

**TABLE 02: RAISED HOMOCYSTEINE LEVEL WITH RESPECT TO AGE GROUPS n= 146**

Age Group	RAISED HOMOCYSTEINE LEVEL		Total
	Yes	No	
40 to 49	16 (44.44%)	20 (55.56%)	36 (100%)
50 to 59	14 (29.17%)	34 (70.83%)	48 (100%)
60 to 70	28 (45.16%)	34 (54.84%)	62 (100%)
<b>Total</b>	<b>58 (69.8%)</b>	<b>88 (30.2%)</b>	<b>146 (100%)</b>

Chi square= 1.667, p= 0.43

**TABLE 03: RAISED HOMOCYSTEINE LEVEL WITH RESPECT TO STUDY VARIABLES n= 146**

		Raised homocysteine		Significance
		Yes	No	
	Yes	12	28	Chi sq.=1.08
<b>DM</b>	No	46	60	P= 0.29
	Yes	18	30	Chi sq. = 0.07
<b>HTN</b>	No	40	58	p= 0.78
	Yes	14	32	Chi sq.=1.21
<b>Smoking</b>	No	44	56	P= 0.27
	Yes	8	10	
<b>Family h/o Ischemic stroke</b>	No	50	78	Chi sq.= 0.09 P= 0.75
	Less than 30	28	46	Chi sq.= 0.11
<b>BMI</b>				P=0.73
	More than 30	30	42	

**DISCUSSION:**

Stroke remains one of the most devastating of all neurological diseases, often causing death or gross physical impairment or disability. Worldwide, Asia, Russia and Eastern Europe experience the highest rates of mortality and disability and loss of quality of life because of stroke.

Raised homocysteine level was seen in 58 (33.70%) cases. Similar was seen by other studies as well. [8-10] The mechanism behind this ischemic stroke is that it increases the tendency of the blood to clot. The above mentioned studies have also build up the association that raised homocysteine level increases the likelihood of developing stroke not only in the smaller vessels but also with the larger ones.[8-10] Even the studies have shown this raised level in as high as 75% of the cases. The reason of such higher number in contrast to our study is unknown. The mortality was not assessed in this study but a study from China has seen for this association. They have

noted that elevated tHcy levels during the acute phase of an ischemic stroke significantly increased the mortality chances. In this study majority of the cases were in 2nd quartiles of the Hcy level and that study has revealed that the cases with 3rd and the 4th quartiles had a 1.83-fold or higher increased risk of death after the 48 months of stroke compared with patients in the first quartile. After multivariate analysis, this association remained statistically significant.[12]

Raised homocysteine level was seen in females where it was seen in 32 (45.71%) out of 70 females as compared to 26 (34.21%) out of 76 males; though this difference was not statistically significant with p values of 0.32. A study by Hung et al showed that resistant hypertension was associated with a higher risk of IS, especially in women and elderly patients.[13] The Framingham Heart Study also suggested overall higher lifetime stroke susceptibility in women. Estrogen and estrogen-progestogen

therapy can reduce homocysteine levels in menopausal women. Another study suggested that higher levels of estrogen-homocysteine conjugates could lead to a lowering of homocysteine-induced cardiovascular disease (CVD) risk. Here we observed that elderly women tended to have higher tHcy levels, perhaps because of the lower estrogen levels in older women.[14-16]

This raised level was also seen maximum in age groups of 60 to 70 years affecting 28 (45.14%) out of 62 cases followed by 40 to 49 year in 16 (44.44%) out of 36 cases. This difference was also not found significant with p values of 0.43. This was also observed by many studies.[17] It can be explained by many ways. The higher age was with this increased tendency was explained in females due to hormonal changes. The other factors can be multiple nutritional factor deficiencies like vitamin and antioxidant which is decreased in older age groups to either poor intake or development of comorbidities like GI problems, malignancies etc. that can also either enhance their consumption or decrease their absorption. Contrary to all this, on the other hand the other group which was found higher in this study was relatively young group with age 40 to 49 years. The reason for higher age groups in relatively young age group can be due to genetic factors that were also observed by other studies. However they also did not found significant association. [18-19]

In this study the ischemic stroke was seen in 18 out of 30 cases with a near significant p value of 0.07. This was also seen by other studies that found in higher number.[13-14] Moreover, Yan et al reported that women with mild hypertension had a higher risk of stroke than men and showed that a 10 mm Hg increase in SBP was associated with a 38% increased risk of stroke in women. The other mechanisms involved include the history of pre-existing atherosclerosis, which is also an outcome of DM and HTN. Conflicting results have been found regarding association between homocysteine levels and diabetes mellitus.[20]

The cases with family history of ischemic stroke were found in 8 out of 18 cases with p value of 0.09. This higher number can be explained by the present of genetic factors. Multiple genetic mutations have been proposed to support this theory with higher Hcy levels than those without pre-existing atherosclerosis. It seems that there is an association between economic prosperity and the risk of stroke. Higher prevalence of Hcy in many developing countries could indicate the role of inadequate intake of vitamins and antioxidants in the multi-factorial

causes of stroke. The effect of genetic factors on hyperhomocysteinemia is also important. In fact, these factors may confound the results of epidemiological studies and may render the results statistically unstable.[21-23]

There were few limitations to this study. As this study did not check for the type of vessel involved to development of stroke and it also did not check for the association of raised level with the severity of the stroke.

### CONCLUSION:

Ischemic stroke is a fatal condition. Raised homocysteine level is seen in every 3<sup>rd</sup> cases of ischemic stroke. Hypertension and family history of ischemic stroke has almost significant association with it.

### REFERENCES:

1. Kim AS, Johnston SC. Global variation in the relative burden of stroke and ischemic heart disease. *Circulation*. 2011;124:314–23.
2. Mukherjee D, Patil CG. Epidemiology and the global burden of stroke. *World Neurosurg*. 2011;76(6):85-90.
3. Han L, Wu Q, Wang C, Hao Y, Zhao J, Zhang L, et al. Homocysteine, ischemic stroke, and coronary heart disease in hypertensive patients: a population-based, prospective cohort study. *Stroke*. 2015;46(7):1777-86.
4. Fahimfar N, Khalili D, Mohebi R, Azizi F, Hadaegh F. Risk factors for ischemic stroke; results from 9 years of follow-up in a population based cohort of Iran. *BMC Neurology*. 2012;12:117-19.
5. Ashjazadeh N, Fathi M, Shariat A. Evaluation of homocysteine level as a risk factor among patients with ischemic stroke and its subtypes. *Iran J Med Sci*. 2013;38(3):233-39.
6. Sadiq M, Alam MT, Kanpurwala MA, Khan MS. Frequency of hyper-homocysteinemia in ischemic stroke patients of Karachi. *J Pak Med Assoc*. 2014;64(9):1063-66.
7. Rahman A, Dasgupta R, Quraishi FA, Saha UK, Ali Z, Hossain S, et al. Association between plasma homocysteine level and ischemic stroke. *J Bangladesh Coll Phys Surg*. 2013;31:128-33.
8. Iso H, Moriyama Y, Sato S, Kitamura A, Tanigawa T, Yamagishi K, et al. Serum total homocysteine concentrations and risk of stroke and its subtypes in Japanese. *Circulation*. 2004;109:2766–2772.
9. Parnetti L, Caso V, Santucci A, Corea F, Lanari A, Floridi A, et al. Mild hyperhomocysteinemia is a risk-factor in all

- etiologiical subtypes of stroke. *Neurol Sci.* 2004;25:13–17. doi: 10.1007/s10072-004-0219-5.
10. Tay SY, Ampil ER, Chen CP, Auchus AP. The relationship between homocysteine, cognition and stroke subtypes in acute stroke. *J Neurol Sci.* 2006;250:58–61. doi: 10.1016/j.jns.2006.06.028.
  11. Rahman A, Dasgupta R, Quraishi FA, Saha UK, Ali Z, Hossain S, et al. Association between plasma homocysteine level and ischemic stroke. *J Bangladesh Coll Phys Surg.* 2013;31:128-33.
  12. Shi, Z, Guan Y, Huo YR. Elevated Total Homocysteine Levels in Acute Ischemic Stroke Are Associated With Long-Term Mortality. *Stroke.* 2015; 46: 2419-2425
  13. Hung CY, Wang KY, Wu TJ, Hsieh YC, Huang JL, Loh el-W, et al. Resistant hypertension, patient characteristics, and risk of stroke. *PLoS One.* 2014;9:e104362.
  14. Gorgui J, Gorshkov M, Khan N, Daskalopoulou SS. Hypertension as a risk factor for ischemic stroke in women. *Can J Cardiol.* 2014;30:774–782.
  15. Sundström J, Sullivan L, D’Agostino RB, Jacques PF, Selhub J, Rosenberg IH, et al. Plasma homocysteine, hypertension incidence, and blood pressure tracking: the Framingham Heart Study. *Hypertension.* 2003;42:1100–1105.
  16. Bonassi MR, Chada BE, Eduardo FC, Marcelo L E, De Lima G R. Effects of estrogen and estrogen-progestogen therapy on homocysteine levels and their correlation with carotid vascular resistance. *Gynecol Endocrinol.* 2007;23:619–624.
  17. Hao L, Ma J, Zhu J, Stampfer MJ, Tian Y, Willett WC, et al. High prevalence of hyperhomocysteinemia in Chinese adults is associated with low folate, vitamin B12 and vitamin B6 status. *J Nutr* 2007; 137: 407-13
  18. Malinow MR, Bostom AG, Krauss RM. Homocyst(e)ine, diet, and cardiovascular diseases: a statement for healthcare professionals from the Nutrition Committee, American Heart Association. *Circulation* 1999; 99: 178-82.
  19. Madonna P, de Stefano V, Coppola A. Hyperhomocysteinemia and other inherited prothrombotic conditions in young adults with a history of ischaemic stroke. *Stroke* 2002; 33: 51-6.
  20. Adunsky A, Weitzman A, Fleissig Y. The relation of plasma total homocysteine levels to prevent cardiovascular disease in older patients with ischaemic stroke. *Ageing (Milano)* 2000; 12: 48-52.
  21. Christopher R, Nagaraja D, Shankar SK. Homocysteine and cerebral stroke in developing countries. *Curr Med Chem.* 2007;14:2393–401.
  22. Sánchez-Moreno C, Jiménez-Escrig A, Martín A. Stroke: roles of B vitamins, homocysteine and antioxidants. *Nutr Res Rev.* 2009;22:49–67.
  23. Khan U, Crossley C, Kalra L, Rudd A, Wolfe CD, Collinson P, et al. Homocysteine and its relationship to stroke subtypes in a UK black population: the south London ethnicity and stroke study. *Stroke.* 2008;39:2943–9.