



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**Available online at: <http://www.iajps.com>

Research Article

**DESCRIPTIVE STUDY ON DIABETIC FOOT IN DIABETIC
PATIENTS AT BBH RAWALPINDI**¹Dr.Sunnaina Tahir, ²Dr. Maryam Tariq, ³Dr. Kainat¹Bahria University Medical and Dental College²Rawalpindi Medical University; Rawalpindi³Foundation University, Islamabad**Abstract:**

Objective: Diabetic foot is common and frequent problem in medical ward, in this study we will determine the knowledge and foot care, common cause of diabetic foot in patients with DM and treatment outcome. **Study Design:** Descriptive / cross sectional study **Place and Duration of Study:** This study was conducted at the Department of Medicine, BBH Rawalpindi, from June 2017 to December 2017. **Materials and Methods:** 108 patients were selected for this study. Both male and female participated. Informed consent was taken from all the patients. Study was done using questionnaire translated into English. **Results:** 78 were male and 30 were female mean age was 52.61 ± 4.21. Non healing wound was present on right foot in 90 patients and left foot in 28 patients. Gangrene was present in 30 patients RBS, FBS and HAlc was elevated in all patients. Curettage was done in 53 patients and amputation was done in 13 patients. All patients were on Insulin therapy and broad spectrum antibiotics were given to all patients. **Conclusion:** Diabetic foot is common problem in our country. Main reason is uncontrolled blood sugar and awareness of the patients. Most of the patients come with irregular diet on and off treatment. Duration of diabetes was prolonged in all patients. Education of the patient about disease diet and proper treatment of diabetes. Lifestyle of the patient and morbidity and mortality can be reduced.

Key Words: Diabetes mellitus, Diabetic foot, Ischemia, Neuropathy, Gangrene**Corresponding author:****Dr.Sunnaina Tahir,**

Bahria University Medical and Dental College

QR code



Please cite this article in press Sunnaina Tahir et al., *Descriptive Study on Diabetic Foot in Diabetic Patients at BBH Rawalpindi.*, Indo Am. J. P. Sci, 2018; 05(10).

INTRODUCTION:

Diabetes mellitus is global health problem; over past two decades its incidence is increased¹ Epidemiologically 30 million cases of diabetes were present in 1985. 177 million in 2000. 285 million in 2010 and more than 360 million people will be affected by DM in 2030.² DM occurs in developed countries but recent new cases of type 2 DM are found in developing countries [3]. One of the complication of Diabetic foot ulcers, result due to combination of minor trauma infection foot deformity, peripheral neuropathy [4]. One lower limb is lost every 30 seconds due to diabetic foot Globally majority of hospital admissions are due to Diabetic Foot [5]. Diabetic foot is associated with hospitalization, morbidity and long-time of stay in hospital than other complication of DM.⁶ Newly diagnosed type 2 diabetes mellitus have peripheral vascular disease or peripheral neuropathy about 10%. Global prevalence of diabetic foot is increasing worldwide⁷ in diabetic patients stiffness of ligaments is due to non-enzymatic glycation. Loss of protective sensation and loss of coordination of foot and leg muscles are due to neuropathy, both elevate mechanical stress during walking. Majority of diabetic patients are not aware of foot care regular foot examination. Regarding complication of diabetes mellitus amputation of lower limb is preventable [8]. In patients with diabetes mellitus lower limb amputations are due to foot ulcer caused by peripheral vascular disease, poor cleaning of foot, bare foot walking unsuitable foot wear and delay in medical treatment [8]. These risk are modified by education of the patient [8]. Physical examination of the patient having diabetic wound is divided into examination of wound and general condition of limbs, assessment of vascular insufficiency (Ischemia). Sensory examination of foot or assessment of neuropathy and Staging of diabetic foot wound, including depth of soft tissue and bony involvement [9] investigations necessary are blood CP ,RBS, FBS HbA1c, creatinine, x-ray ,CT, MRI ,Doppler ultrasound, CT, MRI.

Treatment of diabetic wound depends upon managing systemic and local factors, give up smoking, glycaemic control, control of blood pressure, hyperlipidaemia, ischemic heart disease if present obesity, renal insufficiency [11] offload the wound by therapeutic footwear [12]. Use of saline similar dressings daily [12]. Indeed debridement and appropriate antibiotic to treat osteomyelitis and infection Application of recombinant growth factors or grafts¹³. If arterial insufficiency is not present, hyperbaric oxygen is beneficial.

MATERIALS AND METHODS:

This study was conducted in the department of Medicine BBH, Rawalpindi. 108 diabetic patients were included in this study. Both male and female patients participated. Informed consent was taken from all the patients. Data collected using questionnaire. Detailed history was taken. Duration of diabetes, type of diabetes, dietary habits, smoking history, treatment history other complication of diabetes, physical Examination, Examination of foot sensory neuropathy was assessed by 10 gm, monofilament, force applied on one or more anatomic sites on the planter surface of foot on each foot area [14].

Vibration sense was observed by using 128Hz tuning fork, ankle jerk and patellar jerk. Palpation of dorsal is pedis, posterior tibial arteries, and Doppler ultrasound of the limb. Inspection of the skin hair nails presence of any callus, corn or ulcers. Any foot deformity static analysis was done using SPSS 15 version.

RESULTS:**Table No.1: Age**

		Freq- uency	Percent	Valid Percent	Cumu- Lative Percent
Valid	35.00	1	.9	.9	.9
	39.00	1	.9	.9	1.9
	44.00	1	.9	.9	2.8
	46.00	1	.9	.9	3.7
	47.00	2	1.9	1.9	5.6
	48.00	3	2.8	2.8	8.3
	49.00	11	10.2	10.2	18.5
	50.00	11	10.2	10.2	28.7
	51.00	17	15.7	15.7	44.4
	52.00	9	8.3	8.3	52.8
	53.00	8	7.4	7.4	60.2
	54.00	9	8.3	8.3	68.5
	55.00	11	10.2	10.2	78.7
	56.00	5	4.6	4.6	83.3
	57.00	3	2.8	2.8	86.1
	58.00	5	4.6	4.6	90.7
	59.00	4	3.7	3.7	94.4
	60.00	3	2.8	2.8	97.2
	61.00	2	1.9	1.9	99.1
	62.00	1	.9	.9	100.0
	Total	108	100.0	100.0	

108 patients were selected for this study 78 were female and 30 were males. 56 were farmers, 29 housewife, 15 businessmen and 6 were in Govt. service. 81 patients were uneducated, 8 patients primary, 13 patients middle and 6 patients matriculate. 83 patients were smokers.

Table No.2: Neuropathy

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	.00	10	9.3	9.3	9.3
	1.00	98	90.7	90.7	100.0
	Total	108	100.0	100.0	

Table No.3: Ischemic

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	.00	21	19.4	19.4	19.4
	2.00	87	80.6	80.6	100.0
	Total	108	100.0	100.0	

Table No.4: HbA1C

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	6.90	2	1.9	1.9	1.9
	7.00	9	8.3	8.3	10.2
	7.50	14	13.0	13.0	23.1
	7.80	2	1.9	1.9	25.0
	8.00	21	19.4	19.4	44.4
	8.30	1	.9	.9	45.4
	8.40	1	.9	.9	46.3
	8.50	16	14.8	14.8	61.1
	9.00	32	29.6	29.6	90.7
	9.50	4	3.7	3.7	94.4
	10.00	6	5.6	5.6	100.0
	Total	108	100.0	100.0	

Table No. 5: Type Diabetes

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1.00	14	13.0	13.0	13.0
	2.00	94	87.0	87.0	100.0
	Total	108	100.0	100.0	

Table No.6: Descriptive Statistics

	N	Min.	Max.	Mean	Std. Deviation
Age	108	35.00	62.00	52.6111	4.21797
Sex	108	1.00	2.00	1.2778	0.44999
Occupation	108	1.00	4.00	1.8056	0.93187
Neuropathy	108	0.00	1.00	0.9074	0.29121
Ischemic	108	0.00	2.00	1.6111	0.79524
FBS	108	145.00	231.00	179.6667	17.91569
RBS	108	240.00	530.00	380.2870	63.87357
HbA1C	108	6.90	10.00	8.3713	0.80513
Haemoglobin	108	7.50	11.00	9.5556	0.91543
L.Count	108	9500	11230	10483.21	472.67972
Diabetes Type	108	1.00	2.00	1.8704	0.33746
Duration Diabetes	108	6.00	33.00	14.3148	5.99478
Education Level	108	1.00	4.00	1.4815	0.91183
Valid N(Listwise)	108				

Table No.7: Anova

	Sum of squares	df	Mean square	F	Sig
Sex					
Between Groups	1.241	1	1.241	6.441	0.013
Within Groups	20.426	106	0.19		
Total	21.667	107			
Occupation					
Between Groups	13.283	1	13.283	17.681	0.000
Within Groups	79.634	106	0.751		
Total	92.917	107			
Neuropathy					
Between Groups	0.138	1	0.138	1.636	0.204
Within Groups	8.936	106	0.084		
Total	0.974	107			
Ischemic					
Between Groups	22.493	1	22.493	52.781	0.000
Within Groups	45.173	106	0.426		
Total	67.667	107			
FBS					
Between Groups	2381.043	1	2381.043	7.896	0.006
Within Groups	31962.957	106			
Total	34344.00	107			
RBS					
Between Groups	48786.255	1	48786.255	13.337	0.000
Within Groups	387755.8	106			
Total	436542.1	107			
HbA1C					
Between Groups	5.382	1	5.382	8.917	0.004
Within Groups	63.979	106	0.604		
Total	69.361	107			
Haemoglobin					
Between Groups	0.123	1	0.123	0.145	0.704
Within Groups	89.544	106	0.845		
Total	89.667	107			
Age					
Between Groups	480.974	1	480.974	35.836	0.000
Within Groups	1422.693	106	13422		
Total	1903.667	107			
L.Count					
Between Groups	51.216	1	51.216	0.000	0.988
Within Groups	23906543	106	225533.423		
Total	23906594	107			
Duration. Diabetes					
Between Groups	2305.035	1	2305.035	158.631	0.000
Within Groups	1540.261	106	14.531		
Total	3845.296	107			
Education Level					
Between Groups	14.428	1	14.428	20.519	0.000
Within Groups	74.535	106	0.703		
Total	88.963	107			

Non healing wound was present in right foot in 90 patients and in left foot in 28 patients. Depth of wound was wegener's class1 in 21 patients, 42 in class2, 32 in class3, and 13 in class 4. Gangrene was present in 30 patients, curettage was done in 53 patients and amputation was done in 13 patients. 21 patients were pure neuropathic, 10 patients were pure ischemic and 77 patients were mixed. On examination of foot pure ischemic were 2pts 82 were mixed neuropathic using the test by monofilament tuning fork and Doppler ultrasound. RBS range – 240 to 53, FBS- 145 to 231, HbA1c- 6.9 to 10, Cholesterol level- increase more than

200 in 92 pts, Retinopathy -30pts, Hypertension-80patients, Blood cp-Hb%- 7.5 to 11, L leukocyte count increase from 9500 to 11230, In statical analysis male sex denoted by 1 and female by 2, education level uneducated by 1, primary by 2, middle by 3 and matric by 4. Occupation farmer by 1, housewife by 2, businessmen by 3 and Govt. service by 4.

DISCUSSION:

Globally number of patients with Diabetes mellitus will increase 366 million in 2030¹⁵ out of 108 patients majority of the patients was male, with irregular dietary pattern. Patients were not aware of the severe complications about diabetes mellitus, it was observed that long duration of diabetes increases the incidence of diabetic foot as pathologic process occurs in about 10 years. This problem may occur due to delayed diagnosis

It was observed in previous study in Srilanka¹⁶. Majority of the patients developed diabetic foot in 5th to 6th decades of life. Age about 55-60 years have been found in other studies.¹⁷ Risk of diabetic foot estimated to be 15% in patients with diabetes mellitus. In recent studies it could be 25%⁷ Peripheral neuropathy observed to be common cause of diabetic foot.¹⁸ Education of the patients is necessary for loss of sensation in feet even in minor trauma. Education about blood glucose control in patients with prolonged history of diabetes. These patients must have other complications like diabetic risk of injury¹⁹ Presence of Micro vascular and macro vascular complication of diabetes mellitus, regular screening is compulsory Diabetic Nephropathy and retinopathy increased diabetic foot could be due to micro angiopathic changes in studies diabetic foot healing. Process delayed due to renal impairment²⁰ It was observed hypertension with diabetic foot several studies.²¹ Smoking with Diabetic foot higher percentage of smokers was found in patients with

amputation of limb.²¹ Presence of char coat joint increases the risk of gangrene and amputation²² Diabetic foot complication are increased by peripheral vascular disease Chronic complication of diabetes coronary artery disease CVD retinopathy associated with diabetic foot gangrene and amputation²³ Mosset al and larvery explained association between long duration of diabetes and foot complications²³ according to researchers diabetic foot and amputation increases with diabetes history more than 10 years²³ obesity was associated with increased risk of diabetic foot. Poor glucose control and HbA1c was contributory factor in diabetic foot and their contribution of peripheral neuropathy and micro vascular complication²⁴ Nurses as members of the diabetic care team play role in public education, patient care, health care health management and quality of life improve. Nurses help patients to have movement for those patients who have lost their foot. Nurses teach patients to use assists devices²⁵

CONCLUSION:

In this study main reason of diabetic foot was poor glycaemic control for prolonged period. Development of peripheral neuropathy peripheral vascular disease associated with retinopathy, diabetic nephropathy hypertension, irregular dietary habits. And poor knowledge about diabetes treatment and its complications Educations about diabetes and diabetic foot self-care is necessary management method. Education of foot care, motivation, support, financial assistance is necessary. Involvements of family member, psychological assistance, management of comorbidities retinopathy vasculopathy are helpful. It is necessary to start education about diabetic foot as the symptoms of neuropathy occurs give up smoking education about diet mobility, patients life style patients quality of life can be improved risk of amputation are decreased.

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES:

1. Ramachandran A, Snehalatha C, Shetty AS, Nandita A. Trends in prevalence of diabetes in Asian countries. *World J Diabetes* 2012;3:110-117.
2. Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: Global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Res Clin Pract* 2011; 94:311-321.
3. Chuang LM, Tsai ST, Huang BY, Tai TY. The status of Diabetes control in Asia—across-sectional survey of 24 317 patients with diabetes mellitus in 1998. *Diabet Med* 2002; 19(12):978-985.

4. Reiber GE, Vileikyte L, Boyko EJ, Del Aguila M, Smith DG, Lavery LA, et al. Causal pathways for incident lower-extremity ulcers in patients with diabetes from two settings. *Diabet Care* 1999; 22: 157-162.
5. Levin ME. Pathophysiology of diabetic foot lesions. *Clinical Diabetes Medicine*. In: Davidson JK, Louis ST, editors. Mosby; 1991.p. 504-520.
6. Boyko EJ, Ahroni JH, Smith DG, Davignon D. increased mortality associated with diabetic foot ulcer. *Diabetic Med* 1996; 13:967-972.
7. Andrew J, Gunne R, Jan A. the global burden of diabetes foot disease. *Lancet* 2005;366:1719-24
8. Wikblad K, Smide B, Bergstrom A, Kessii J, Mugusi F. Outcome of clinical foot examination relation to self-perceived health and glycaemic control in a group of urban Tanzanian diabetic patients. *Diabetes Res Clin Pract* 1997; 37:185-92.
9. Abbas ZG, Archibald LK. Challenges for management of the diabetic foot in Africa. *Diabetes Care* 2001; 24(1): 84-8.
10. Pinzur MS, Slovenkai MP, Trepman E, Sheilds NN. Guidelines for diabetic foot care: recommendations endorsed by the diabetes committee of the American orthopedic foot and ankle society. *Foot Ankle Int* 2005; 26(1):113-9.
11. Margolis DJ, Kantor J, Santanna J, Strom BL, Berlin JA. Risk factors for delayed healing of neuropathic diabetic foot ulcers: a pooled analysis. *Arch Dermatol* 2000; 136(12):1531-5.
12. Hilton JR, Williams DT, Beuker B, B Miller DR, Harding KG. Wound dressings in diabetic foot. *Disease Clin Infect Dis* 2004; 39:S100-S103.
13. Benett SP, Griffiths GD, Schor AM, Leese GP, Schor SL. Growth factors in the treatment of diabetic foot ulcers. *Br J Surg* 2003; 90:133-146.
14. Singh N, Armstrong DG, Lipsky BA Preventing foot ulcers in patients with diabetes. *JAMA* 2005; 293:217-228.
15. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes estimates for the year 2000 and projections for 2030. *Diabetes care* 2004; 27:1047-1053.
16. Weerasuriya N, Siribaddana S, Wijeweera I, Dissanayeka A, Wijeskera J, Frnando DJ. The prevalence of peripheral neuropathy in newly diagnosed patients with non- insulin-dependent diabetes mellitus. *Ceylon Med J* 1998; 43(1):19-21.
17. Nathera A, Siok Beeb C, Yiong Hauke C, Chewa JLL, Lina CB, Neoa Sh CB, et al. Epidemiology of diabetic foot problems and predictive factors for limbs loss. *J Diabet Compl* 2008; 22:77-82.
18. Reiber GE, Vileikyte L, Boyko EJ, Aguila M, Smith DG, Lavery LA, et al. Casual pathways for incident lower- extremity ulcers in patients with Diabetes from two settings. *Diabet Care* 1999; 22:157-62.
19. Pataky Z, Vischer U. Diabetic foot disease in the elderly. *Diabetes Metab* 2007; 33 Suppl 1:S56-65.
20. Bruun C, Siersma V, Guassora AD, Holstein P, de Fine Olivarius N. Amputations and foot ulcers in patients newly diagnosed with type 2 Diabetes mellitus and observed for 19 years. The role of age, gender and co-morbidity. *Diabetic Med*.2013; 30 964-972. doi: 10.1111/dme.12196
21. Deery HG, Sangeorzan JA, Saving the diabetic foot with social reference to the patient with chronic renal failure. *Infect Dis Clin North Am* 2001; 15: 953-981.
22. Sohn MW, Stuck RM, Pinzur M, Lee TA, Budiman Mak E. Lower extremity amputation risk after charcotartgropathy and diabetic foot ulcer. *Diabet Care* 2010; 33:98-100.
23. Lavery LA, Armstrong DG, Vela SA, Quebedeaux TL, Fleischli JG. Practical Criteria for screening patients at high risk for diabetes foot ulceration. *Arch Int Med* 1998; 158:157-162.
24. Dinh T, Tecilazich T, Kafanas A, Doupis J, Gnardelis C, Leal E, et al. Mechanism involved in the development and Healing of diabetic foot ulceration. *Diabetes* 2012;61:2937-2947
25. Armstrong DG, Lavery LA, Wu S, Boulton AJ. Evaluation of removable and irremovable cast walkers in the healing of diabetic foot wounds. *Diabet Care* 2005; 28(3):551-554.