



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF  
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3483721>Available online at: <http://www.iajps.com>

Research Article

**ANALYSIS OF POST TRAUMATIC OPTIC NERVE  
DECOMPRESSION IN PATIENTS WITH NO LIGHT  
PERCEPTION**Abdul Shakoor<sup>1</sup>, Mubeen Ullah Meo<sup>2</sup>, Abdul Majid Mujahid<sup>3</sup><sup>1</sup>DHQ Hospital Lodhran, <sup>2</sup>Jinnah Hospital Lahore, <sup>3</sup>Basic Health Unit daiwal, Khushab.**Article Received:** August 2019**Accepted:** September 2019**Published:** October 2019**Abstract:**

**Objective:** To determine if optic nerve decompression has any efficacy in patients with traumatic optic nerve (TON) damage specifically those patients whose optic nerves have been severely damaged and as a result, they suffer no light perception (NLP).

**Material and Methods:** This cross sectional study was conducted in DHQ hospital Lodhran during December 2018 to June 2019.

**Results:** 35 patients still had no light perception at end of the study. By the time others were discharged some had gained a level or two of visual acuity but no cases of perfect vision restored were reported.

**Conclusion:** The methods were relatively useful in reducing discomfort in patients such as significant reduction in swelling but they were not in any way completely responsible for the total recovery of the patients.

**Corresponding author:**

Abdul Shakoor,

DHQ Hospital Lodhran

QR code



Please cite this article in press Abdul Shakoor et al., *Analysis of Post Traumatic Optic Nerve Decompression in Patients with No Light Perception.*, Indo Am. J. P. Sci, 2019; 06(10).

**INTRODUCTION:**

First described by Hippocrates Traumatic Optic Neuropathy (TON) is defined as the loss of vision with no evidence of injury to the optic nerve [1]. Accordingly, TON is rare but very devastating. Optic neuropathy can be on the anterior or posterior of the optic nerve and each classically divided into direct or indirect injuries [2]. Direct TON can result from penetrating injury or broken bone fragment that may lead to optic nerve avulsion, stretch injury, shearing, contusion, lacerations, and disruption. Compressive optic neuropathy can result from orbital haemorrhages [3].

Indirect TON is the most common type of optic nerve injury and accounts for 0.5-5% of closed head injuries [4]. It is believed to result from transmission of shock from an orbital impact to the intracranial portion of optic nerve [4]. Due to lack of understanding of the pathophysiology involved and uncertainty of clinical results, treatment of TON remains generally controversial. Methods such as optic decompression and use of steroids have been brought forward to try alleviating the cases of TON. Our study focuses mainly on optic nerve decompression.

Optical decompression has been used for the treatment of optic canal fractures [5]. Un-roofing the optical canal could relieve pressure in the optic intracranial portion of the optic nerve halting further damage. However, some arguments brought forward voiced concern that further injuries could be meted on the optic nerve resulting in further loss of vision [6]. In this paper a retrospective study of patients with no light perception, admitted in our hospital was done during their admission time, to determine the efficacy of optic nerve decompression.

**METHODS:**

This cross sectional study was conducted in BHU Daiwal, Khushab during December 2018 to June 2019. A group of 54 patients, 53 males and a female, who has suffered optic nerve trauma were put under observation. 23 of them had injured their left eye, 30 their right and 1 patient had injured both eyes. 27 had

sustained injuries from car accidents, 23 from falls, 4 from motorcycle accidents. The timing from injury to treatment was more than 3-21 days. 12 cases took 3-7 days, 8 took 15-21 days and some 5 cases took more than 21 days. Examination of the patients was as follows:

**VISUAL**

Ideally should be determined using a hand held near card, newspapers, magazines or Snell's chart. One should bear in mind that delayed visual loss develops due to secondary optic nerve injury hence visual acuity should be conducted again after 24 hours [7]. A comprehensive history of the patient should be obtained to ensure that the loss of sight is as a result of the injury and not an underlying optic problem. The only drawback of this test is that it can only be done on patients who were conscious and had no swellings over their eyes. The patient's comfort also had to be ensured so that discomfort from other injuries did not bias the feedback they gave the doctor.

**TREATMENT**

Patients were treated with drugs, craniotomy optical canal decompression and endoscopic nerve decompression.

**Statistics:** SPSS17.0 software  $\chi^2$  test,  $P < 0.05$  statistically significant.

**RESULTS:**

In the group of 54 patients who were followed up for a period between 3 to 12 months. At the end of the follow up, 7 patients had recovered visual acuity to different degrees. The other cases showed ineffectiveness and one patient was lost to the follow up. The earliest time that a period recovered visual acuity was one day and the latest recovery was fifteen days after the operation. Improvement of visual acuity was 1 level in 1 patient, two levels in 1 patient, 3 levels in 1 patient and 4 levels in 2 patients. Recovery occurred between 1 and 2 weeks after recovery but ceased by the end of 2 months and remained as was. 5 could count number of fingers shown to them, 4 could perceive finger movement 50 cm away and 7 could see the eye chart.

case	age (yr)	Cause of injury	DOC	Time of lost vision (day)	Result of CT scan	Injured to operation time (day)	Preoperative vision	Time of vision recovery (day)	Vision at discharge	Time and postoperative vision during follow up
1	23	CA	yes	7	FBF,SH	6	NLP	15	NLP	0.04 (12)
2	28	TF	yes	20	FOC,SH,OF	30	NLP	8	Light perception	Finger movement (10)
3	31	TF	yes	30	CLB, FOC,SH	51	NLP	—	NLP	NLP (10)
4	30	TF	yes	3	FOC,OF,SH	10	NLP	—	NLP	NLP (7)
5	29	TF	yes	10	ASBF,SH	18	NLP	7	Finger movement	Finger movement (6)
6	20	TF	yes	1	CLB, SH	14	NLP	—	NLP	NLP (7)
7	33	TF	yes	3	FBF,FOC,SH	3	NLP	13	NLP	light (6)
8	17	CA	yes	9	CLB,SH	24	NLP	—	NLP	Lost of follow up
9	30	TF	Yes	1	OF	8	NLP	—	NLP	NLP (4)
10	33	CA	Yes	1	FOC	4	NLP	—	NLP	NLP (3)
11	19	TF	Yes	14	CLB,SH,OF	21	NLP	—	NLP	NLP (3)
12	34	CA	Yes	2	SH, FBF	7	NLP	3	Finger movement	0.6 (6)
13	26	TF	Yes	3	FOC,SH	13	NLP	—	NLP	NLP (4)
14	46	CA	Yes	2	FOC, TBF	6	NLP	—	NLP	NLP(6)
15	38	TF	Yes	14	SH,OF	23	NLP	—	NLP	NLP(7)
16	23	CA	Yes	4	FOC, TBF	6	NLP	—	NLP	NLP(9)
17	24	TF	Yes	1	FOC,SH	7	NLP	11	light	light (12)
18	29	CA	Yes	2	SH	6	NLP	—	NLP	NLP(8)
19	31	TF	Yes	1	FOC,SH	8	NLP	—	NLP	NLP(4)
20	26	CA	Yes	1	FBF	3	NLP	—	NLP	NLP(8)
21	29	CA	Yes	1	FOC,OF	7	NLP	10	NLP	light(9)
22	30	CA	Yes	5	SH	8	NLP	—	NLP	NLP(3)
23	36	CA	Yes	3	FBF, OF	5	NLP	8	NLP	light(6)
24	24	TF	Yes	2	FOC,SH	15	NLP	—	NLP	NLP(6)
25	28	CA	Yes	4	SH, FBF	22	NLP	—	NLP	NLP(5)
26	28	CA	Yes	3	SH	5	NLP	7	light	light(7)
27	31	TF	Yes	5	OF, TBF	8	NLP	—	NLP	NLP(6)
28	34	TF	Yes	2	FOC,SH	20	NLP	—	NLP	NLP(7)
29	33	CA	Yes	4	FOC,SH	18	NLP	—	NLP	NLP(3)
30	20	CA	Yes	2	SH	10	NLP	—	NLP	NLP(4)
31	13	CA	Yes	1	FOC,OF	2	NLP	—	NLP	NLP(3)
32	43	MA	Yes	2	FOC,OF	3	NLP	2	Finger Movement	Finger Movement (7)
33	20	CA	Yes	2	FOC,OF	3	NLP	2	Visual Acuity(0.04)	0.04(3)
34	28	CA	Yes	1	FOC,SH	2	NLP	3	Index	Index(5)
35	47	TF	Yes	7	OF,TBF	8	NLP	3	LP	LP(8)
36	51	CA	Yes	3	OF,SH	4	NLP	—	NLP	NLP(6)

37	20	CA	Yes	3	FOC,OF,TB F	5	LP	3	VISUAL ACUITY 0.2	0.2(4)
38	18	CA	Yes	4	OF	5	NLP	2	INDEX FINGER	INDEX (10) FINGER
39	22	CA	Yes	2	OF	3	NLP	5	MOVEMEN T	MOVMENT (3)
40	17	MA	Yes	2	FOC,OF	3	NLP	—	NLP	NLP(9)
41	32	TF	Yes	5	OF,TBF	6	NLP	6	INDEX	INDEX(5)
42	21	MA	Yes	3	FOC,OF	4	LP	—	NLP	NLP(8)
43	39	TF	Yes	2	FOC,SH	3	NLP	1	LP	LP(6)
44	42	CA	Yes	1	FOC,OF	2	LP	4	VISUAL ACUITY 0.08	0.08(9)
45	16	CA	Yes	5	OF	6	LP	3	VISUAL ACUITY 0.6	0.6(7)
46	29	TF	Yes	3	FOC,OF	4	NLP	7	INDEX	INDEX(3)
47	35	MA	Yes	8	FOC,OF,TB F	10	NLP	—	NLP	NLP(10)
48	41	TF	Yes	1	FOC,OF	2	NLP	3	VISUAL ACUITY 0.04	0.04(4)
49	19	CA	Yes	3	FOC,OF,SH	4	NLP	—	NLP	NLP(6)
50	28	TF	Yes	1	FOC,OF	2	NLP	3	INDEX	INDEX (11)
51	31	CA	Yes	1	FOC,OF,TB F	3	NLP	5	NLP	NLP(6)
52	32	TF	Yes	4	FOC	5	NLP	—	NLP	NLP(4)
53	18	CA	Yes	5	FOC	2	NLP	—	NLP	NLP(3)
54	23	TF	Yes	8	FOC	4	NLP	—	NLP	NLP(5)

CLB=contusion and laceration of brain,; FOC=fracture of optic canal,;  
SH=Sinuses hematoma, ; OF=orbital fractures, ; NLP=no light perception, DOC-Disturbance of Consciousness  
CA- car accident. TF- to fall FBF- frontal bone fracture TBF-Temporal bone fracture ASBF- Anterior skull base  
fracture

### DISCUSSION:

Generally any blindness sustained immediately after an injury exhibits a poor prognosis but gradual loss of vision exhibits a better prognosis. Immediate treatment of injuries is believed to be of paramount importance if desired results are expected [18]. However, there has been no observed correlation between immediate treatment and recovery in patients [19]. In the event that surgical decompression occurred immediately after the accident then it meant that the optic nerve was severely damaged or that majority or all the nerves were fractured, and surgical decompression would be of insignificant effect [20, 21]. In a report by Ohlsson and Svensson early decompression of the injured optic nerve reduces axonal degeneration and showed significant improvement in adult rats [23]. However, there was little evidence to support that these cases improved as a direct result of the surgery [8]. A study by the International Optic Nerve Trauma the largest multicentre study of TON to date analysed 133

people with indirect TON who were treated within 7 days of trauma. No clear evidence was found to support benefits of optic nerve decompression surgery [22].

Endoscopic decompression at other times is an alternative method that alleviates pressure in the canal. It is considered to have advantages such low morbidity, retention of a sense of smell, a shorter recovery time as compared to surgical procedure, no surgical scar and does not interfere with children's growth [18, 8]. However, there is no evidence to support that there is significant improvement in patients after the procedure.

### CONCLUSION:

The decompression of the optic canal by surgery or endoscopy served to alleviate some of the symptoms of the optic nerve injury. Swelling of the nerve reduced and in cases eliminated. A level or two of improved visual acuity was reported. However, there is no proof

that they in any way help in the restoration of the vision in patients.

#### REFERENCES:

1. Chandwick, J & Mann WN The Medical works of Hippocrates. Oxford: England, Backwell 1950
2. Levin L Traumatic Optic Neuropathy; Roy and Fraufelder's Current Ocular Therapy 6th edition Eds: Roy FH. Fraunfelders FS. Philadelphia 2007; 5(2):73-78
3. Nishi T, Ueda T et al. Traumatic optic neuropathy caused by blunt injury to the inferior orbital rim. J Neuroophthalmom, 2006; 26:44-46
4. Van S GP, Biousse V, Lynn MJ et al Neuroophthalmic manifestation of the head trauma. J Neuroophthamol, 2000; 21:112-117
5. Sofferman B. The recovery potential of the optic nerve, Laryngoscope, 1995; 105:1-3
6. Guyer RG, Miller RN MD, Long MD, PHD, et al. Visual function following optical Decompression via Craniotum, Balitimore 1985; 7:112-116
7. Levin LA, Beck RW, Joseph MP The treatment of traumatic optic neuropathy: the international Optic Nerve Trauma Study. Ophthalmology, 1999; 106:1268-1277
8. Steinsapir KD, Goldberg RA Traumatic optic neuropathy; a critical update. Comp Ophthalmol Update, 2005; 6:11-612
9. Steinsapir KD, Goldberg RA Traumatic optic neuropathy Surv Ophthalmol, 1994; 38:487-518
10. Lessel L Traumatic optic neuropathy and visual system injury In Shingleton BJ, Hersh PS, Kenyon KR, Editors Eye trauma. St Louis: Mosby Year Book; 2010 June; 23:371-379
11. Tsai HH, Jengb SF, Linb TS, et al. Predictive value of computed tomography in visual outcome in indirect traumatic optic neuropathy complicated with periorbital facil bone structure. Clin. Neurol. Neurosurg. 2005 april;107(3):200-206
12. Lee AG. Imaging for neuroophthalmic manifestations of the head trauma. J Neuroophthalmol, 2004;138:852-862
13. Kountakis SE, Maillard AAJ, Urso R, et al. Endoscopic approach to traumatic visual loss. Otolaryngol. Head Neck Surg. 1997 June, 116(6): 652-655
14. Rajiniganth MG, Gupta AK, Gupta A, et al. Traumatic optic neuropathy: visual outcome following combined therapy protocol. Arch Otolaryngol Head Neck Surg. 2003 Nov, 129(11):1203-1206
15. Wohlrab TM, Maas S, de Carpentier JP. Surgical decompression in traumatic optic neuropathy. Acta Ophthalmol Scand 2002 Jun, 80(3):287-293
16. Gupta AK, Gupta AK, Gupta A, et al. Traumatic optic neuropathy in pediatric population: early intervention or delayed intervention? Int J Pediatr Otorhinolaryngol. 2007 april, 71(4):559-562
17. Lipkin AF, Woodson GE, Miller RH .Visual loss due to Orbital Fracture. Arch Otolaryngol Head Surg, 1983 Jun, 91(3):276-279
18. Spoor TC, Hartel WC, Lensik DB, et al. Treatment of Traumatic Optic Nerve Neuropathy with Corticosteroid Am J Ophthamol, 1990; 110:665-9.