



CODEN [USA]: IAJ PBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3474159>Available online at: <http://www.iajps.com>

Research Article

**STUDY TO DETERMINE BACTERIAL VAGINOSIS
COMPLICATIONS IN PREGNANCY**Dr Irum Siddique¹, Dr Asma Noreen², Dr Rubina Zain²¹ Nishter Medical College, Multan, ² Quaid e Azam Medical College, Bahawalpur.**Article Received:** August 2019**Accepted:** September 2019**Published:** October 2019**Abstract:**

Objective: In women, the utmost usual causes of vaginal discharge is bacterial vaginosis. The risk of preterm birth, premature rupture of membranes (PROM) and abortion has been suggested to be advanced in pregnancies pretentious by bacterial vaginosis. This study was planned to govern the bacterial vaginosis incidence and its relation with preterm labor, miscarriage, premature rupture of membranes and intrauterine fetal death (IUFD).

Study Design: A cohort study.

Place and Duration: In the Gynaecology and Obstetrics department in Tehsil Headquarter Hospital Jalalpur Peerwala, Multan for two year duration from May 2017 to May 2019.

Methods: Before 20 weeks of gestation; 136 women identified with bacterial vaginosis were included in exposure group and in non-exposed group there were 397 pregnant women deprived of bacterial vaginosis. The relationship between premature birth, miscarriage, intrauterine fetal death and bacterial vaginosis and premature rupture of membranes in pregnancy were evaluated. In SPSS version 18.0 data was analyzed. For Fisher's exact test and statistical comparisons; Chi-square test were used and Yates correction were performed when required. $P < 0.05$ was taken as statically significant.

Results: The frequency of intrauterine fetal death, abortion, premature rupture of membranes and preterm delivery were 0.9%, 1.3%, 1.3%, 3.6% correspondingly. There was no arithmetical association between abortion and bacterial vaginosis, but the association between bacterial vaginosis and preterm delivery, intrauterine fetal death and PROM was substantial ($p < 0/001$).

Conclusion: The results of this analysis show that preterm delivery, intrauterine fetal death and premature rupture rates are greater in females with bacterial vaginosis than women who do not.

Key Words: Preterm labor, abortion, bacterial vaginosis.

Corresponding author:**Dr. Irum Siddique,**

Nishter Medical College, Multan.

QR code



Please cite this article in press Irum Siddique et al., *Study to Determine Bacterial Vaginosis Complications in Pregnancy*, Indo Am. J. P. Sci, 2019; 06(10).

INTRODUCTION:

In women, the utmost usual causes of vaginal discharge are bacterial vaginosis. Incidence rates have been reported between 10% and 31%. Bacterial vaginosis can result in pregnancy complications such as preterm labor, miscarriage, chorioamnionitis and PROM [1-3]. With bacterial vaginosis, pregnant women -before the 16th week of gestation have been shown to have an early birth risk of 1.5 to 7.0 times further as compared those who don't have [4-5]. The concept is that the problem should be managed before the four months is grounded on the contrivance of bacterial vaginosis in preterm labor. In pregnancy; the bacterial vaginosis prevalence and its association with preterm birth contrasts between various populations and rely on the sociodemographic factors, clinical setting, gestational age, diagnostic criteria and additional influences [6-7]. There is a lack of information on the bacterial vaginosis effects on pregnancy [8]. The outcome of this analysis will help implement detection, health decision makers design and in pregnancy helps in treatment plans for bacterial vaginosis. This shows the significance of the study.

MATERIALS AND METHODS:

This cohort study was held in the Gynaecology and Obstetrics department in Tehsil Headquarter Hospital Jalalpur Peerwala, Multan for two year duration from May 2017 to May 2019. Total 777 pregnant women listed in the hospital of Southern Punjab before the 20th week of pregnancy were included. Before 20 weeks of gestation; 136 women identified with bacterial vaginosis were included in exposure group and in non-exposed group there were 397 pregnant women deprived of bacterial vaginosis. Criteria that could not be included gravid women with a history of preterm birth, abortion, pregnancy complication history, multifetal pregnancy, and also females under eighteen or above forty years of age. We are also investigating pregnancy consequences. For Fisher's exact test and statistical comparisons; Chi-square test was used and Yates correction were performed when required. $P < 0.05$ was taken as statically significant.

RESULTS:

The bacterial vaginosis incidence in our study among gravid women was 17.5%. There was no substantial variation between the occupation, birth order and age group between the unexposed and exposed groups (Table 1).

Table-I: Characteristics of the pregnant women according to bacterial vaginosis (B.V) status

	<i>B.V positive</i>	<i>B.V Negative</i>	<i>Significance</i>
Pregnancy			NS
First	93 (35.5)	169 (64.5)	
Second	36 (25.5)	105 (74.4)	
>3rd	34 (34.3)	65 (65.6)	
Age Group			NS
<30	124 (32.7)	255 (67.3)	
>30	39 (32)	83 (68)	
Education			NS
Primary	45 (30)	105 (70)	
Intermediate	98 (34.7)	184 (65.2)	
College	18 (27.7)	47 (72.3)	

B.V: Bacterial Vaginosis

The frequency of intrauterine fetal death, abortion, premature rupture of membranes and preterm delivery were 0.9%, 1.3%, 1.3%, 3.6% correspondingly. There was no arithmetical association between abortion and

bacterial vaginosis, but the association between bacterial vaginosis and preterm delivery, intrauterine fetal death and PROM was substantial ($p < 0/001$) Table 2.

Table-II: Outcome of pregnancy based on bacterial vaginosis status

	B.V Positive	B.V Negative	Relative risk(95% CI)	Significance
Abortion				NS
Yes	4 (2.9)	132 (97.1)	3.89 (0.88-17.17)	
No	3 (0.8)	394 (99.2)		
IUFD				<0.0005
Yes	4 (2.9)	132 (97.1)	11/68 (1.32-103.57)	
No	1 (0.3)	396 (99.7)		
Preterm labor				<0.0001
Yes	13 (9.6)	123 (90.4)	6.32(2.24-16.31)	
No	6 (1.5)	391 (98.5)		
PROM				<0.0001
Yes	6 (4.4)	130 (95.6)	25.71(3.29-20.07)	
No	1 (0.3)	396 (99.7)		

B.V: Bacterial Vaginosis

IUFD: Intra uterine fetal death

PROM: Premature rupture of membrane

11.68 was the comparative IVFD risks ($P < 0.0005$), 25.71 ($P < 0.0001$) and 6.32 ($P < 0.0001$) correspondingly, in preterm labor and positive pregnancies of B.V.

DISCUSSION:

The bacterial vaginosis prevalence ranges between 6.9% and 27% in analysis conducted in public hospitals and academic medical centre in the USA¹⁰. And in other analysis, however, our population was not well considered and studied. In our analysis, the frequency of miscarriage, bacterial vaginosis, preterm delivery, IUFD and PROM were 1.3%, 17.5%, 3.6%, 0.9% and 1.3% correspondingly. There was no arithmetical association between abortion and bacterial vaginosis, but the association between bacterial vaginosis and preterm delivery, intrauterine fetal death and PROM was substantial ($p < 0/001$). It has been supposed that the preterm birth risk is greater in pregnancies convoluted by bacterial vaginosis [11-12]. The Kalinkaj et al study shows that the preterm delivery rate was approximately 15.8% compared to 9.2% in individuals without bacterial vaginosis and concluded that bacterial vaginosis must be diagnosed early and treatment may reduce early delivery [13]. In a Paris study, preterm delivery in females with bacterial vaginosis was approximately 26% associated with 10.7% in other females. In Hiller's analysis, with bacterial vaginosis low birth weight infants and preterm birth were very usual in females. In Benedetto et al study, preterm labor and PROM are significantly associated with bacterial vaginosis, but not with IUD. Ralph et al also measured the risk of falling to 18.5% in women with bacterial vaginosis and 18.5% in females deprived of bacterial vaginosis [14]. In other analysis it is suggested that females with B.V have greater preterm labor risk, and that antibacterial therapy may reduce the preterm labor risk. According to this analysis, in some women the bacterial vaginosis

treatment may cause miscarriage, IUFD, preterm labor and decrease in PROM. The study by Ugwumda et al. exhibited that detection and asymptomatic bacterial vaginosis treatment in the 1st month of the 2nd trimester may reduce the preterm delivery rate in gravid females, and many studies have shown same results, but Goffeng et al. There was no association between bacterial preterm labor and vaginosis, and in Toronto another study showed that bacterial vaginosis treatment did not reduce the preterm labor risk and had no effect on pregnancy outcomes¹⁵. Oakeshott et al. determined that bacterial vaginosis did not have better prognosis of spontaneous abortions before the 16th week of pregnancy. Liversedge et al found an important relationship in the 1st trimester.

CONCLUSION:

According to IUFD results, preterm delivery and PROM rates were higher in pregnant females with bacterial vaginosis compared to pregnant women without bacterial vaginosis. The gynaecologist and midwife should evaluate pregnant women before the 20th week of gestation to detect BV and treat them to reduce the risk of IUFD and PROM.

REFERENCES:

1. Lannon, Sophia MR, Kristina M. Adams Waldorf, Tina Fiedler, Raj P. Kapur, Kathy Agnew, Lakshmi Rajagopal, Michael G. Gravett, and David N. Fredricks. "Parallel detection of lactobacillus and bacterial vaginosis-associated bacterial DNA in the chorioamnion and vagina of pregnant women at term." *The Journal of Maternal-Fetal & Neonatal Medicine* 32, no. 16 (2019): 2702-2710.
2. De Vulder, Annelies, Guy Mulinganya, Jerina Boelens, Geert Claeys, Erick Hendwa, Yvette Kujirakwinja, Freddy Kampara et al. "Prevalence, risk factors and associated adverse pregnancy

- outcome of bacterial vaginosis in pregnant women in Bukavu, Democratic Republic of Congo." In *European Congress of Clinical Microbiology & Infectious Diseases*. 2019.
3. Powell, Anna Maya, Judy R. Shary, Christopher Loudon, Vishwanathan Ramakrishnan, Allison Ross Eckard, and Carol L. Wagner. "Association of Bacterial Vaginosis with Vitamin D in Pregnancy: Secondary Analysis from the Kellogg Pregnancy Study." *American Journal of Perinatology Reports* 9, no. 03 (2019): e226-e234.
 4. Shimaoka, Masao, Yoshie Yo, Kunihiko Doh, Yasushi Kotani, Ayako Suzuki, Isao Tsuji, Masaki Mandai, and Noriomi Matsumura. "Association between preterm delivery and bacterial vaginosis with or without treatment." *Scientific reports* 9, no. 1 (2019): 509.
 5. Haahr, Mona Aarenstrup Karlsen, and Lene Hee Christensen. "Bacterial vaginosis in pregnancy and risk of spontaneous preterm delivery." *Update* (2019).
 6. Dunlop, Anne L., Sheila L. Jordan, Erin P. Ferranti, Cherie C. Hill, Shiven Patel, Li Hao, Elizabeth J. Corwin, and Vin Tangpricha. "Total and Free 25-Hydroxy-Vitamin D and Bacterial Vaginosis in Pregnant African American Women." *Infectious diseases in obstetrics and gynecology* 2019 (2019).
 7. Brookheart, Rita T., Warren G. Lewis, Jeffrey F. Peipert, Amanda L. Lewis, and Jenifer E. Allsworth. "Association between obesity and bacterial vaginosis as assessed by Nugent score." *American journal of obstetrics and gynecology* 220, no. 5 (2019): 476-e1.
 8. Rebouças, Karinne F., José Eleutério Jr, Raquel C. Peixoto, Ana Paula F. Costa, Ricardo N. Cobucci, and Ana K. Gonçalves. "Treatment of bacterial vaginosis before 28 weeks of pregnancy to reduce the incidence of preterm labor." *International Journal of Gynecology & Obstetrics* (2019).
 9. Kamga, Yiewou Marguerithe, John Palle Ngunde, and Jane-Francis KT Akoachere. "Prevalence of bacterial vaginosis and associated risk factors in pregnant women receiving antenatal care at the Kumba Health District (KHD), Cameroon." *BMC pregnancy and childbirth* 19, no. 1 (2019): 166.
 10. Vodstrcil, Lenka A., Ms Erica Plummer, Christopher K. Fairley, Gilda Tachedjian, Matthew G. Law, Jane S. Hocking, Ms Karen Worthington, Ms Mieken Grant, Nita Okoko, and Catriona S. Bradshaw. "Combined oral contraceptive pill-exposure alone does not reduce the risk of bacterial vaginosis recurrence in a pilot randomised controlled trial." *Scientific reports* 9, no. 1 (2019): 3555.
 11. Amalokwu, S., P. I. Okonta, and E. Ebonu. "Prevalence of bacterial vaginosis among antenatal attendees with abnormal vaginal discharge in a secondary health facility in Delta State, Nigeria." *Tropical Journal of Obstetrics and Gynaecology* 36, no. 1 (2019): 85-87.
 12. Vieira-Baptista, Pedro, and Jacob Bornstein. "Candidiasis, Bacterial Vaginosis, Trichomoniasis and Other Vaginal Conditions Affecting the Vulva." In *Vulvar Disease*, pp. 167-205. Springer, Cham, 2019.
 13. Udoh, Ubong Aniefiok, Boniface U. Ago, Anthony A. Iwuafor, Ernest A. Ochang, Okokon I. Ita, and Anthony U. Agbakwuru. "Sociodemographic determinants of Bacterial Vaginosis risk among pregnant women in southern Cross River State, Nigeria." (2019).
 14. Rac, Hana, Alyssa P. Gould, Lea S. Eiland, Brooke Griffin, Milena McLaughlin, Kayla R. Stover, Christopher M. Bland, and P. Brandon Bookstaver. "Common Bacterial and Viral Infections: Review of Management in the Pregnant Patient." *Annals of Pharmacotherapy* 53, no. 6 (2019): 639-651.
 15. Balkus, Jennifer E., Kayla A. Carter, and R. Scott McClelland. "Lessons from Suppressive Therapy and Periodic Presumptive Treatment for Bacterial Vaginosis." *Current infectious disease reports* 21, no. 10 (2019): 34.