



CODEN [USA]: IAJPBB

ISSN : 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4062190>Available online at: <http://www.iajps.com>

Research Article

**RECENT ADVANCES AND TREATMENTS OF CHILDHOOD
NEPHROTIC SYNDROME****Dr. Sarwar Rafique¹, Dr. Uswa Majeed¹, Dr. Sumaira Rafique¹**
¹Nishtar Hospital, Multan**Article Received:** August 2020**Accepted:** September 2020**Published:** October 2020**Abstract:**

Introduction: Nephrotic syndrome is characterized by gross proteinuria, hypo albuminemia, hyperlipidemia, and peripheral edema. The etiology of nephrotic syndrome in adults is complex and ranges from primary glomerulonephritis to secondary forms. **Objectives of the study:** The main objectives of the study is to find the recent advances and treatments of childhood nephrotic syndrome in Pakistan. **Methodology of the study:** This analytical study was conducted at Nishtar Hospital, Multan during 2019. The data was collected from 100 patients of both genders which was suffering from nephrotic syndrome. The basic purpose of this data is to find and investigate the recent advances and treatments of nephrotic syndrome in Pakistani hospitals. **Results:** The study group comprised of 100 patients 72 (93.5%) were initial steroid resistant and 28 were late non-responders. Gender distribution showed 49 (63.6%) males and 28 (36.4%) females with a ratio of 1.75. Age range of patients was 1-15 years with a mean of 8.11 ± 3.58 years. Sixty nine (89.6%) patients underwent renal biopsy. Patients with focal segmental glomerulo sclerosis (FSGS) were least likely to respond to treatment followed by mesangio proliferative glomerulonephritis and minimal change disease. **Conclusion:** Nephrotic syndrome can increase your child's risk of infection and blood clots. It always affects both kidneys and usually appears in the early years of your child's life. Most children with this disorder outgrow it by young adulthood. We conclude that NS in children is a difficult disease with significant morbidity.

Corresponding author:**Dr. Sarwar Rafique ***,
Nishtar Hospital, Multan

QR code



Please cite this article in press Sarwar Rafique et al, **Recent Advances And Treatments Of Childhood Nephrotic Syndrome.**, Indo Am. J. P. Sci, 2020; 07(10).

INTRODUCTION:

Nephrotic syndrome is characterized by gross proteinuria, hypo albuminemia, hyperlipidemia, and peripheral edema. The etiology of nephrotic syndrome in adults is complex and ranges from primary glomerulonephritis to secondary forms. Primary forms of nephrotic syndrome in adults are comprised of three histological disease entities: idiopathic membranous nephropathy (iMN), minimal change disease (MCD), and focal segmental glomerulo sclerosis (FSGS)¹. The basis for therapy of primary nephrotic syndrome is mostly of supportive nature. Supportive strategies include antihypertensive and anti-proteinuric therapy and dietary recommendations. Patients with nephrotic syndrome are also at increased risk to develop thromboembolism. In patients with membranous nephropathy, the adjusted hazard ratio for thromboembolism was 10.8 compared to patients with IgA nephropathy³. In contrast, for patients with FSGS the hazard ratio was 5.9. Hence, anticoagulant therapy is recommended in patients with a primary nephrotic syndrome, especially in iMN and serum albumin < 2.5 mg/dl. In 2014, Lee et al. proposed a practical approach to prophylactic anticoagulation therapy in patients with iMN. The presented model takes into account the serum albumin concentration, the individual patient's bleeding risk, and the risk tolerance as reflected by the selected benefit-to risk ratio³.

Idiopathic, or primary, nephrotic syndrome is often used to describe the group of patients for whom no specific cause has been identified, and the histology is relatively non-specific⁴. These patients will usually receive immunosuppression without knowledge of the mechanism and be categorized according to response. So the challenge is to understand and categorize the underlying injury at a molecular level and therefore adapt treatments according to the likely mechanism⁵.

Objectives of the study

The main objectives of the study is to find the recent advances and treatments of childhood nephrotic syndrome in Pakistan.

METHODOLOGY OF THE STUDY:

This analytical study was conducted at Nishtar Hospital, Multan during 2019. This study was done according to the rules and regulations of hospital ethical authority. The data was collected from 100 patients of both genders which was suffering from nephrotic syndrome. The basic purpose of this data is to find and investigate the recent advances and treatments of nephrotic syndrome in Pakistani hospitals. The main objectives in management of nephrotic syndrome are to induce quick remission to ensure freedom from edema and consequences of persistent nephrotic state such as hyperlipidemia, protein malnutrition, thromboembolic episodes, severe infections, etc. Demographic data of the patients, treatment received, and outcome of treatment and complications were recorded.

Statistical analysis

A chi-square test was used to examine the difference in the distribution of the fracture modes (SPSS 19.0 for Windows, SPSS Inc., USA).

RESULTS:

The study group comprised of 100 patients 72 (93.5%) were initial steroid resistant and 28 were late non-responders. Gender distribution showed 49 (63.6%) males and 28 (36.4%) females with a ratio of 1.75. Age range of patients was 1-15 years with a mean of 8.11 ± 3.58 years. Sixty nine (89.6%) patients underwent renal biopsy. Patients with focal segmental glomerulosclerosis (FSGS) were least likely to respond to treatment followed by mesangio proliferative glomerulonephritis and minimal change disease.

Table 01: Demography of 100 selected patients

Category	Number (%age)
Biopsied	69(89.6%)
Un biopsied	08(10.4%)
Initial SR	72(93.5%)
Late SR	05(6.5%)
Males	49(63.6%)
Females	28(36.4%)
Age(Years)	
<4	22(28.6%)
4-10	31(40.2%)
>10	24(31.2%)

The patients in group 1 received CsA plus PDN, MMF plus PDN, combined CsA and MMF plus PDN, and intravenous MP pulses \pm oral PDN and CPM as S1,S2, S3, and S4 treatment respectively. Following S1, 31/61 patients (50.8%) achieved complete remission, 5/61 (8.2%) were partial responders, and 25/61 (41%) were non-responders.

Table 02: Treatment of nephrotic syndrome

Sequential Treatment Step	Drugs	Number of Patients	Complete Remission	Partial Remission	No Remission
Step 1	CsA + PDN	61 (79.2%)	31 (50.8%)	05 (8.2%)	25 (41%)
Step 2	MMF + PDN	30 (49.2%)	03 (4.9%)	07 (11.5%)	20 (32.8%)
Step 3	CsA + MMF +PDN	27 (44.3%)	06 (9.8%)	09 (14.8%)	12 (19.7%)
Step 4	IVMP + PDN ± CPM	21 (34.4%)	04 (6.6%)	02 (3.3%)	15 (24.6%)
Total: Steps 1-4		61 (100%)	44 (72%)	02 (3.3%)	15 (24.6%)

DISCUSSION:

There have been no consistent clinical cues to either whether a patient with nephrotic syndrome has the risk of becoming steroid-resistant in the future or whether they will suffer recurrence post-transplant⁶. There are weak clinical associations with recurrence (for example, age at onset of disease, race, and serum albumin at diagnosis or time to first dialysis/transplant. Interestingly, the last two features may point to the possibility that CFD has a more aggressive presentation and natural course, compared with the monogenic or 'other' groups⁶. To address the question of whether there are clinical features that pertain to CFD, we hypothesized that patients with the archetypal CFD, those with post-transplant recurrence, would have distinct early clinical features regarding their initial response to immunosuppression⁷. If a patient has initial steroid sensitivity (otherwise described as secondary steroid resistance), they are likely to have an immune-mediated circulating factor causing their underlying disease and therefore high risk of recurrence⁸.

Treatment of SRNS in children continues to pose a therapeutic challenge to the pediatric nephrologists. The lack of large-scale randomized controlled trials leads to a paucity of strong evidence to inform treatment decisions. The treatment strategies are heterogeneous with variable efficacy and side effects' profile. Optimal strategies with least toxicity remain to be determined⁹. Without effective treatment, progression to the end-stage kidney disease is very likely¹⁰.

CONCLUSION:

Nephrotic syndrome can increase your child's risk of infection and blood clots. It always affects both kidneys and usually appears in the early years of your child's life. Most children with this disorder outgrow it by young adulthood.

REFERENCES:

1. A. Howman, T. L. Chapman, M. M. Langdon et al., "Immunosuppression for progressive membranous nephropathy: a UK randomised controlled trial," *The Lancet*, vol. 381, no. 9868, pp. 744–751, 2013
2. Abeyagunawardena AS, Sebire NJ, Risdon RA, et al. Predictors of long-term outcome of children with idiopathic FSGS. *Pediatr Nephrol* 2007; 22(2):215-221
3. Mekahli D, Liutkus A, Ranchin B et al. Long-term outcome of idiopathic steroid resistant nephrotic syndrome: a multicenter study. *Pediatr Nephrol* 2009;24:1525-15327.
4. Naudet P. Treatment of childhood nephrotic syndrome with a combination of cyclosporine and prednisone. *French Society of Pediatric Nephrology. Kidney Int* 1997;125:981-86.
5. Lombel RM, Hodson E, Gipson D. Treatment of steroid -resistant nephrotic syndrome in children- new guidelines from KDIGO. *Pediatr Nephrol* 2012; DOI 10.1007/s00467-012-2304-8.
6. Hafeez F, Ahmad TM, Anwar S. Efficacy of steroids, cyclosporin, and cyclophosphamide in steroid resistant idiopathic nephrotic syndrome. *JCPSP* 2005;15(6):329-332
7. Hodson EM, Habashy D, Craig JC. Intervention for idiopathic steroid resistant nephrotic syndrome in children. *Cochrane Database Syst Rev* 2006;(2): CD003594.
8. Kari JA, Halawani M, Mokhtar G, Jalalah SM, Anshasi W. Pattern of steroid resistant nephrotic syndrome in children living in the kingdom of Saudi Arabia. *Saudi J Kidney Dis Transpl* 2009;20(5):854-857.
9. Gulati S, Kher V, Sharma RK, Gupta A. Steroid response pattern in Indian children with nephrotic syndrome. *Acta Pediatr Scand* 1994;83:530-533.
10. Gulati S, Sengupta D, Sharma RK, Sharma A, Gupta RK, Singh U, et al. Steroid resistant nephrotic syndrome: Role of histopathology. *Ind Pediatr* 2006;43(10):55-60.