



CODEN [USA]: IAJPBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4134693>Available online at: <http://www.iajps.com>

Research Article

A COMPARATIVE STUDY OF SYSTOLIC AND DIASTOLIC BLOOD PRESSURE OF HEALTHY PEOPLE WITH DIVERSE BMI RANGES

¹Dr Zarnain Tariq, ²Dr Mirza Aun Muhammad Baig, ³Dr Zulaikha Yaqoob
^{1, 2, 3}Shalamar Medical and Dental College Lahore

Article Received: August 2020

Accepted: September 2020

Published: October 2020

Abstract:

Background and Objectives: Obesity, especially central obesity, is linked to hypertension. The purpose of this study was to compare the systolic and diastolic blood pressure (B.P.) in healthy, overweight and obese controls.

Study design: An analytical cross-sectional study.

Place and Duration: In the Medicine Unit of Shalamar Medical and Dental College Lahore for one-year duration from June 2019 to June 2020.

Material and methods: 81 (81) healthy people, both men and women (aged 18-60) were selected from the general population. Blood pressure was measured with a mercury sphygmomanometer. Body mass index (BMI) was calculated as weight (kilogram) divided by the square of your height (square meters). We divided the participants into a control group, overweight and obese according to their BMI. In the next step, we divided the subjects into categories according to gender and BMI. Group I consisted of fifty-four (54) men, including 18 controls, 18 overweight and 18 obese. Group II comprised twenty-seven (27) women, including 09 controls, 09 overweight and 09 obese.

Results: The results of this study indicate significantly higher systolic and diastolic B.P. in overweight and obese subjects compared to the control group, except for men in the control group and overweight men, who had a statistically insignificant difference in systolic B.P.

Conclusion: We can conclude that the increase in body weight seems to be related to the increase in B.P. systolic and diastolic.

Key words: body mass index (BMI), central obesity, diastolic blood pressure, hypertension, obesity, obesity, overweight, systolic blood pressure.

Corresponding author:**Dr Zarnain Tariq,**

Shalamar Medical and Dental College Lahore

QR code



Please cite this article in press Zarnain Tariq et al, A Comparative Study Of Systolic And Diastolic Blood Pressure Of Healthy People With Diverse BMI Ranges., Indo Am. J. P. Sci, 2020; 07(10).

INTRODUCTION:

Being overweight is the sixth most important risk factor contributing to the overall burden of disease worldwide. More than 1 billion adults and 10% of children are currently classified as overweight or obese¹⁻². In the US population, 30% of adults are obese and another 35% are overweight as calculated by body mass index (BMI) as weight (in kilograms) / height (in meters)³⁻⁴. The increasing incidence of obesity is more and more often accepted as one of the most important risk factors for the development of hypertension⁵⁻⁶. Obesity, and central obesity in particular, is associated with hypertension. It has been reported that at least two-thirds of the occurrence of hypertension can be directly attributed to obesity. Moreover, the combination of obesity and high blood pressure increases the risk of cardiovascular disease. Weight loss is often accompanied by a decrease in B.P., supporting the hypothesis that an increase in BMI contributes to the higher incidence of hypertension⁷⁻⁸. Therefore, lifestyle modifications, including weight loss and increased physical activity, should be considered an early stage in the treatment of obesity-related hypertension. In addition, public health strategies to reduce the obesity epidemic are also expected to significantly reduce the burden of hypertension⁹. Moreover, it is suggested that the effect of weight gain on B.P must be confirmed with future longitudinal studies.

MATERIALS AND METHODS:

The study was conducted at the Medicine Unit of Shalamar Medical and Dental College Lahore for one-year duration from June 2019 to June 2020. Eighty-one (81) healthy individuals, both male and female (aged 18-60 years), were selected from the general population. Known hypertension has been ruled out. In addition, we excluded patients with kidney disease, diabetes and other systemic diseases. We also excluded pregnant women. Informed consent was obtained from all subjects and the relevant history was recorded in the proformas. Height was measured in centimeters (cm) on a standard height scale and weight in kilograms on a Camry scale. Body mass index was calculated as weight (kilogram) divided by the square of height (square meters). Subjects with BMI 20-24.9 kg / m² were considered

the control group, while subjects with BMI 25-29.9 kg / m² and > 30 kg / m² were classified as overweight and obese, respectively. Blood pressure was measured with a mercury sphygmomanometer after the participant rested for at least 05 minutes in a sitting position. Although it is recommended that a diagnosis of hypertension be made only after an elevation was found on three readings on different occasions over a period of several months, the purpose of this study was to compare the BP of people with different BMI ranges and not have a diagnosis of hypertension, so we relied on one reading. Initially, the entire study population was divided according to the BMI index into a control group, overweight and obese. In the next step, we divided the subjects into categories according to gender and BMI. Group I included fifty-four (54) men. This group included 18 controls, 18 overweight and 18 obese. Group II included twenty-seven (27) women. This group included 09 controls, 09 overweight and 09 obese (Table 1). Statistical analysis: Data was entered and analyzed using SPSS 18.0. Means, standard deviation (SD), and standard error of the mean (SEM) were calculated for the systolic and diastolic BP of overweight and obese controls, controls, overweight and obese men, controls, overweight women and obese women. Student's t-test was used to compare the group means. A p value of less than 0.05 (5%) was considered significant.


RESULTS:

Means, SD and SEM for systolic and diastolic blood pressure for the whole study population (Group I + Group II) are given in Table 2. Means, SD and SEM for systolic and diastolic BP for people in Group I are given in Table 3. Mean, SD and SEM for systolic and diastolic blood pressure for people from group II is given in Table 4. In the whole study population (groups I and II) we compared the systolic and diastolic blood pressure of overweight and obese controls. There was a significant difference between the mean BP systolic control group and the overweight group (117.59 ± 9.74 versus 124.07 ± 9.71 , $p < 0.05$). Similarly, there was a significant difference between the mean BP diastolic control group and overweight (76.67 ± 5.71 vs. 84.81 ± 8.93 , $p < 0.05$).

Table 1: Distribution of Control, Over Weight and Obese subjects among Group I (Males) and Group II (Females)

	Control	Over weight	Obese	Total
Group I (Male)	18	18	18	54
Group II (Female)	09	09	09	27
	27	27	27	81

There was a significant difference between the mean BP systolic control group and obesity (117.59 ± 9.74 vs. 132.59 ± 11.63 , $p < 0.05$). Similarly, there was a significant difference between the mean BP diastolic control group and obesity (76.67 ± 5.71 vs 90.00 ± 7.46 , $p < 0.05$) (Fig. 1).

**Fig. 1: Comparison of Mean Systolic and Diastolic B.P of control, overweight and obese in total study population**

To account for gender differences, we also made group comparisons. In group I, no significant difference was found between the mean BP systolic control group and overweight (121.11 ± 9.00 vs. 125.00 ± 9.85 , $p > 0.05$).

Total study population (Group I + Group II)	Mean	SD	SEM
Control Systolic	117.59	9.74	1.87
Control Diastolic	76.67	5.71	1.10
Overweight Systolic	124.07	9.71	1.86
Overweight Diastolic	84.81	8.93	1.71
Obese Systolic	132.59	11.63	2.23
Obese Diastolic	90.00	7.46	1.43

However, there was a significant difference between the mean BP diastolic control group and overweight (78.06 ± 4.58 vs. 85.00 ± 9.85 , $p < 0.05$). There was a significant difference between the mean BP systolic control group and obesity (121.11 ± 9.00 vs. 134.44 ± 10.41 , $p < 0.05$).

Group I	Mean	SD	SEM
Control Systolic	121.11	9.00	2.12
Control Diastolic	78.06	4.58	1.08
Overweight Systolic	125.00	9.85	2.32
Overweight Diastolic	85.00	9.85	2.32
Obese Systolic	134.44	10.41	2.45
Obese Diastolic	91.11	6.76	1.59

Similarly, a significant difference was found between the mean BP diastolic control group and obesity (78.06 ± 4.58 vs 91.11 ± 6.76 , $p < 0.05$) (Fig. 2).




Fig 2: Comparison of Mean systolic and Diastolic B.P. of control, overweight and obese in Group I

In group II, a significant difference was found between the mean BP systolic control group and the overweight (110.56 ± 7.26 vs. 122.22 ± 9.71 , $p < 0.05$). Similarly, there was a significant difference between the mean BP diastolic control group and overweight (73.89 ± 6.97 vs. 84.44 ± 7.26 , $p < 0.05$).

Group II	Mean	SD	SEM
Control Systolic	110.56	7.26	2.42
Control Diastolic	73.89	6.97	2.32
Overweight Systolic	122.22	9.71	3.23
Overweight Diastolic	84.44	7.26	2.42
Obese Systolic	128.89	13.64	4.54
Obese Diastolic	87.78	8.70	2.90

There was a significant difference between the mean BP systolic control group and obesity (110.56 ± 7.26 vs 128.89 ± 13.64 , $p < 0.05$). Similarly, a significant difference was found between the mean BP diastolic control group and obesity (73.89 ± 6.97 vs 87.78 ± 8.70 $p < 0.05$)

DISCUSSION:

Obesity is associated with an increase in systemic B.P, but the exact mechanisms linking obesity with hypertension are not fully understood¹⁰. Obesity-related hypertension can be attributed to the production of several vasoactive factors from adipose tissue, increased activity of the sympathetic nervous system due to the action of leptin on the central nervous system, and low concentrations of atrial natriuretic factor (ANF) leading to sodium retention and an increase in volume¹¹⁻¹². Results of this study reveal significantly higher systolic and diastolic blood pressure in overweight and obese subjects compared to the control group, with the exception of the control group and overweight men, who had a statistically insignificant difference in systolic blood pressure. Consequently, we can conclude that the

increase in body weight seems to be associated with an increase in systolic and diastolic B.P¹³⁻¹⁴. Similar results are reported by Leenen *et al.*, Mohsen A. *et al.*

and Bramlage *et al.* In addition to the increasing prevalence of hypertension in obese people, the pharmacological treatment of obesity-related hypertension is also a challenge. On the one hand, many workers suggested the pathophysiological effect of weight gain on the increase in B.P, while on the other hand, it was found that the opposite is also true. It has been observed that people of equal weight who had a higher initial B.P. gain weight in the future¹⁵. The coexistence of hypertension and obesity can be attributed to a primary increase in sympathetic tone.

CONCLUSION:

We conclude that the increase in body weight seems to be related to the increase in systolic and diastolic B.P.

REFERENCES:

1. Santos AS, Rodrigues AP, Rosa LP, Sarrafzadegan N, Silveira EA. Cardiometabolic

- risk factors and Framingham Risk Score in severely obese patients: Baseline data from DieTBra trial. *Nutrition, Metabolism and Cardiovascular Diseases*. 2020 Mar 9;30(3):474-82.
2. Abbasnezhad A, Falahi E, Gonzalez MJ, Kavehi P, Fouladvand F, Choghakhori R. Effect of different dietary approaches compared with a regular diet on systolic and diastolic blood pressure in patients with type 2 diabetes: A systematic review and meta-analysis. *Diabetes Research and Clinical Practice*. 2020 Apr 4:108108.
 3. Flint AC, Conell C, Ren X, Banki NM, Chan SL, Rao VA, Melles RB, Bhatt DL. Effect of systolic and diastolic blood pressure on cardiovascular outcomes. *New England Journal of Medicine*. 2019 Jul 18;381(3):243-51.
 4. Chan Sun M, Ruhomally ZB, Boojhawon R, Neergheen-Bhujun VS. Consumption of Moringa oleifera Lam leaves lowers postprandial blood pressure. *Journal of the American College of Nutrition*. 2020 Jan 2;39(1):54-62.
 5. Keir DA, Badrov MB, Tomlinson G, Notarius CF, Kimmerly DS, Millar PJ, Shoemaker JK, Floras JS. Influence of Sex and Age on Muscle Sympathetic Nerve Activity of Healthy Normotensive Adults. *Hypertension*. 2020 Sep;76(3):997-1005.
 6. Bae EH, Lim SY, Han KD, Oh TR, Choi HS, Kim CS, Ma SK, Kim SW. Association between systolic and diastolic blood pressure variability and the risk of end-stage renal disease. *Hypertension*. 2019 Oct;74(4):880-7.
 7. Sakaniwa R, Tromp J, Shirai K, Yamagishi K, Tamakoshi A, Iso H. The association of conventionally medicated systolic and diastolic blood pressure level and mortality from cardiovascular disease: is the lower the better in high stroke population?. *Clinical Research in Cardiology*. 2020 Jan 16:1-5.
 8. Byambasukh O, Snieder H, Corpeleijn E. Relation Between Leisure Time, Commuting, and Occupational Physical Activity With Blood Pressure in 125 402 Adults: The Lifelines Cohort. *Journal of the American Heart Association*. 2020 Feb 18;9(4):e014313.
 9. Ntineri A, Kollias A, Zeniodi ME, Vazeou A, Soldatou A, Stergiou GS. Insight into the 24-hour ambulatory central blood pressure in adolescents and young adults. *The Journal of Clinical Hypertension*. 2020.
 10. Tran AH, Flynn JT, Becker RC, Daniels SR, Falkner BE, Ferguson M, Hanevold CD, Hooper SR, Ingelfinger JR, Lande MB, Martin LJ. Subclinical Systolic and Diastolic Dysfunction Is Evident in Youth With Elevated Blood Pressure. *Hypertension*. 2020 Jun;75(6):1551-6.
 11. Reshetnikov E, Ponomarenko I, Golovchenko O, Sorokina I, Batlutskaya I, Yakunchenko T, Dvornyk V, Polonikov A, Churnosov M. The VNTR polymorphism of the endothelial nitric oxide synthase gene and blood pressure in women at the end of pregnancy. *Taiwanese Journal of Obstetrics and Gynecology*. 2019 May 1;58(3):390-5.
 12. Ma Y, Zhang X, Che Q, Wu K, Zhao W, Hu H, Yang Q, Huang A, Chen D. Reference ranges and trajectories for blood pressure in pregnancy: findings from a follow-up study based on China Maternal and Newborn's Health Monitoring System. *Hypertension in Pregnancy*. 2020 Apr 2;39(2):117-25.
 13. Bazerbachi F, Haffar S, Wang Z, Cabezas J, Arias-Loste MT, Crespo J, Darwish-Murad S, Ikram MA, Olynyk JK, Gan E, Petta S. Range of normal liver stiffness and factors associated with increased stiffness measurements in apparently healthy individuals. *Clinical Gastroenterology and Hepatology*. 2019 Jan 1;17(1):54-64.
 14. Pedralli ML, Marschner RA, Kollet DP, Neto SG, Eibel B, Tanaka H, Lehnen AM. Different exercise training modalities produce similar endothelial function improvements in individuals with prehypertension or hypertension: a randomized clinical trial Exercise, endothelium and blood pressure. *Scientific reports*. 2020 May 6;10(1):1-9.
 15. Jobman J, Aguirre T, Koehler A, Struwe L, Wells W. Role of community health fairs in providing health services, improving health of rural residents. *Arc Psy Mental Health*. 2020:042-5.