Maria *et al* 



# CODEN [USA]: IAJPBB

ISSN: 2349-7750

# INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

http://doi.org/10.5281/zenodo.3408379

Available online at: <u>http://www.iajps.com</u>

**Research Article** 

# ANALYSIS OF GASTROINTESTINAL AND LIVER DISEASES IN PAEDIATRICS AMONG LOCAL POPULATION OF PAKISTAN

Dr Maria, Dr Sobia Qadeer, Dr Syeda Hina Haider

## Abstract

**Introduction:** Gastric cancer is the second most common cancer worldwide and almost two-thirds of all cases occur in developing countries. **Objectives of the study:** The basic aim of the study is to analyze the complete blood count (CBC) in gastrointestinal gastrointestinal and liver diseases in paediatrics among local population of Pakistan. **Methodology of the study:** The study was conducted at Health department Punjab during 2019. The data was collected from both genders and the sample size is 70. Detailed history was taken from all patients with special reference to duration of disease, mode of infection, previous history of disease, HBV or HCV or any other kind of infection. A thorough clinical examination was carried out and stigmata of disease and histopathology was noted. **Results:** According to analysis CBC of patients shows that the level of WBCs and RBCs become decreases in gastrointestinal cancer. It shows that GSH level become increases in patients as compared to control. But the GPx become decreases. All other antioxidants become increases in patients group as compared to control. **Conclusion:** It is concluded that the level of antioxidants will be used as a important biomarker in the diagnosis and treatment of gastrointestinal cancer. And the level of blood cells become decreases in diseased condition.

Corresponding author: Dr.Maria,



Please cite this article in press Maria et al., Analysis Of Gastrointestinal And Liver Diseases In Paediatrics Among Local Population Of Pakistan, Indo Am. J. P. Sci, 2019; 06(09).

### IAJPS 2019, 06 (09), 11767-11770

## Maria et al

#### **INTRODUCTION:**

Cancer is a leading cause of death worldwide: it accounted for 7.9 million deaths (around 13% of all deaths) in 2008. The rising life expectancy means that the risk of developing cancer is also increasing. Deaths from cancer worldwide are projected to continue rising, with an estimated 12 million deaths in 2030<sup>1</sup>. As most cancers appear in adults at an advanced age, the burden of cancer is much more important than other diseases in populations with a long life expectancy. The eight leading cancer killers worldwide are also the eight most common in terms of incidence. Together, they account for about 60% of all cancer cases and deaths. They are cancers of the lung, stomach, breast, colon-rectum, mouth, liver, cervix and oesophagus<sup>2</sup>.

Gastric cancer is the second most common cancer worldwide and almost two-thirds of all cases occur in developing countries. It is the fourth most common cancer in men, while in women it is the fifth most common cancer (based on statistic in 2008)<sup>3</sup>. Although the incidence of gastric cancer is declining, it still remains a major health problem and a common cause of cancer mortality worldwide. Gastric cancer carcinogenesis refers to accumulation of genetic alteration of multiple genes such as oncogenes, tumour suppressor and mismatch repair genes<sup>4</sup>.

The dynamic balance between cell proliferation and apoptosis is very important to maintain the homeostasis in human body and gastric carcinogenesis is related to this imbalance. Development of gastric cancer is believed to be a slow process with primary etiological determinants for gastric cancer being exposure to chemical carcinogens and/or infection with *Helicobacter pylori*. It has been reported that gastric cancer also expresses multidrug-resistance associated protein (MRP) and shows lower sensitivity to anti-cancer drugs. Gastric cancer is more common in older populations, usually occurring in the seventh and eighth decades of life<sup>5</sup>.

#### **Objectives of the study**

The basic aim of the study is to analyze the complete blood count (CBC) in gastrointestinal gastrointestinal and liver diseases in paediatrics among local population of Pakistan.

#### **METHODOLOGY OF THE STUDY:**

The study was conducted at Health department Punjab during 2019. The data was collected from both genders and the sample size is 70. Detailed history was taken from all patients with special reference to duration of disease, mode of infection, previous history of disease, HBV or HCV or any other kind of infection. A thorough clinical examination was carried out and stigmata of disease and histopathology was noted. It includes Hemoglobin (Hb), total leucocytes count (TLC), and differential leucocytes count (DLC), platelet count, level of antioxidants and other related factors. We collected data on incidence of gastric cancer. incidence of oesophageal cancer, deaths from gastric cancer, deaths from any cause, and adverse effects arising due to therapy.

#### Statistical analysis

The data were sampled and entered into the SPSS worksheet for analysis. The alpha criterion was set at 0.05 (95% confidence interval [CI]).

#### **RESULTS:**

Table 01 shows the CBC of control and diseased group. According to analysis CBC of patients shows that the level of WBCs and RBCs become decreases in gastrointestinal cancer.

Blood component	Abbreviation used	Reference range	Diseased values				
White blood cells	WBC	4500-11,000/mm <sup>3</sup>	3.8-11.0 x 10 <sup>9</sup> /L				
Red blood cells*	RBC	Male:4.3-5.9million/mm³3.5-5.5million/mm³3.5-5.5	3.9-4.0 million/mm <sup>3</sup>				
Hemoglobin*	HGB	Male: 13.5-17.5 g/dL Female: 12.0-16.0 g/dL	9-11 g/dL				
Hematocrit*	HT	Male: 41%-53% Female: 36%-46%	0.41-0.53				
Mean corpuscular volume	MCV	80-100 μm <sup>3</sup>	70-80				
Mean corpuscular hemoglobin	МСН	25.4-34.6 pg/cell	39-54 pg/cell				
Mean corpuscular hemoglobin concentration	MCHC	31%-36% Hb/cell	4.81-5.58				
Platelets	Platelets	150,000-400,000/mm <sup>3</sup>	150-400 x 10 <sup>9</sup> /L				

 Table 01: CBC in gastrointestinal cancer patients

Table 02 shows the antioxidants level of patients. It shows that GSH level become increases in patients as compared to control. But the GPx become decreases. All other antioxidants become increases in patients group as compared to control.

	group	Ν	Mean	Std. Deviation	Std. Error Mean
GSH	control	10	.93150	.283559	.089669
	patients	17	4.15765	.534536	.129644
GPx	control	10	.78900	.344970	.109089
	patients	17	.17750	.038108	.009242
Catalase	control	10	4.33600	.748750	.236776
	patients	17	3.69382	1.188499	.288253
SOD	control	10	.38020	.165513	.052340
	patients	17	.92471	.814660	.197584
MDA	control	10	2.57200	.814886	.257690
	patients	17	1.80729	1.379333	.334537

#### **DISCUSSION:**

The development of gastric cancer is a multifactorial process and many conditions influence the likelihood of occurrence, of them, family history of gastric cancer<sup>6</sup>, *Helicobacter pylori* infection (a common bacteria that can also cause stomach ulcers), history of an adenomatous gastric polyp larger than 2 centimetres, history of chronic atrophic gastritis, history of pernicious anemia, obesity, alcohol, smoking, red meat and low socioeconomic status are all believed to be important<sup>7</sup>.

In 1994, the International Agency for Research on Cancer and The World Health Organization classified Helicobacter pylori as a type I carcinogen, the exact mechanism leading to gastric carcinoma is not clearly understood. The effects of H. gastric cancer pylori infection on appear multifactorial, involving host and environmental factors as well as differing bacterial strains. H. pylori is most closely associated with intestinal gastric cancers, which follow a stepwise pathway but toward malignancy, similar to that in the colon<sup>8</sup>. In the Correa model of gastric carcinogenesis, gastric inflammation leads to mucosal atrophy, metaplasia, dysplasia, and, ultimately, carcinoma. Studies have shown that H. pylori infection is an independent risk factor for distal gastric cancer, with a 3- to 6-fold increased risk relative to those without the infection<sup>9</sup>. In patients with *H. pylori*, the presence of specific gene polymorphisms increases the risk of developing gastric carcinoma<sup>10</sup>.

#### **CONCLUSION:**

It is concluded that the level of antioxidants will be used as a important biomarker in the diagnosis and treatment of gastrointestinal cancer. And the level of blood cells become decreases in diseased condition.

#### **REFERENCES:**

- 1. Wang R. The epidemiological study of subtype risk factors of gastric cancer. Zhonghua LiuXingBingXue ZaZhi. 1993;14:295–299.
- Sun X, Mu R, Zhou Y, Dai X, Qiao Y, Zhang S, Huangfu X, Sun J, Li L, Lu F. 1990-1992 mortality of stomach cancer in China. Zhonghua ZhongLiu ZaZhi. 2002;24:4– 8.
- Correa P. Human gastric carcinogenesis: a multistep and multifactorial process--First American Cancer Society Award Lecture on Cancer Epidemiology and Prevention. Cancer Res. 1992;52:6735–6740.
- Rugge M, Correa P, Dixon MF, Hattori T, Leandro G, Lewin K, et al. Gastric dysplasia: The Padova international classification. Am J Surg Pathol. 2000;24:167–76.
- Smith MG, Hold GL, Tahara E, El-Omar EM. Cellular and molecular aspects of gastric cancer. World J Gastroenterol. 2006;12:2979– 90.
- Yasui W, Sentani K, Motoshita J, Nakayama H. Molecular pathobiology of gastric cancer. Scand J Surg. 2006;95:225–31.
- Graziano F, Humar B, Guilford P. The role of the E-cadherin gene (CDH1) in diffuse gastric cancer susceptibility: From the laboratory to clinical practice. Ann Oncol. 2003;14:1705– 13.
- Censini S, Lange C, Xiang Z, Crabtree JE, Ghiara P, Borodovsky M, et al. Cag, a pathogenicity island of Helicobacter pylori, encodes type I-specific and disease-associated virulence factors. Proc Natl Acad Sci U S A. 1996;93:14648–53.
- 9. Covacci A, Censini S, Bugnoli M, Petracca R, Burroni D, Macchia G, et al. Molecular characterization of the 128-kDa immunodominant antigen of *Helicobacter*

# IAJPS 2019, 06 (09), 11767-11770

Maria *et al* 

*pylori* associated with cytotoxicity and duodenal ulcer. Proc Natl Acad Sci U S A. 1993;90:5791–5.

 Cunningham D, Starling N, Rao S, Iveson T, Nicolson M, Coxon F, et al. Capecitabine and oxaliplatin for advanced esophagogastric cancer. N Engl J Med. 2008;358:36–46.