



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.3408381>Available online at: <http://www.iajps.com>**Research Article****ANALYSIS OF RELATIONSHIP OF OBESITY, DIABETES
MELLITUS AND PERIODONTITIS IN PAKISTAN**

Dr Maria, Dr Sobia Qadeer, Dr Syeda Hina Haider

Abstract:

Introduction: Diabetes is a major cause of mortality globally, and it has been estimated that 400 million people worldwide will suffer from it by 2030. Despite the fact that hereditary qualities seems to assume an essential part in the advancement of diabetes, examine recommends that dietary decisions driven by natural and financial components are of critical significance. **Objectives of the study:** The main objective is to find the relations among obesity, diabetes mellitus and periodontitis in common people of Pakistan. **Methodology of the study:** The study was conducted at Health department Punjab during 2019. There was 100 patients which was visit the health center during this time period. **Result:** Results shows that values of glucose become high diabetic patients as compared to normal values. IFG factor shows that diabetic people suffer more from oral health problems as compared to others. **Conclusion:** Obesity for many is a sensitive subject and may not be easy to bring up. Hence the need for more education. The demand for dental hygienists who specialize in treating patients with obesity may increase as the condition itself is increasing and the link with inflammatory diseases is relevant.

Corresponding author:

Dr. Maria,

QR code



Please cite this article in press Maria et al., *Analysis Of Relationship Of Obesity, Diabetes Mellitus And Periodontitis In Pakistan*, *Indo Am. J. P. Sci.*, 2019; 06(09).

INTRODUCTION:

Diabetes is a major cause of mortality globally, and it has been estimated that 400 million people worldwide will suffer from it by 2030. Despite the fact that hereditary qualities seems to assume an essential part in the advancement of diabetes, examine recommends that dietary decisions driven by natural and financial components are of critical significance. Amazing eating regimens assume an essential part in diabetes avoidance.¹ Suitable dietary adherence can enhance insulin affectability and glycemic control, and consequently add to way of life change and general personal satisfaction. Nonetheless, past research recommends that dietary adherence is seemingly among the most troublesome foundations of diabetes administration.² Higher HEI scores demonstrate nearer adherence to current dietary rules for singular food and supplement gatherings. For the sufficiency segments, for example, vegetables and natural product, a higher score demonstrates higher utilization. Dietary proposals depend on the useful effects of devouring products of the soil and expressly stress their constructive outcomes of decreasing corpulence and certain sorts of growths. The last three segments of the HEI incorporate refined grains, sodium, and discharge (calories from strong fats, liquor, and included sugars) and a higher score demonstrates bring down utilization.^{3,4}

Obesity creates risk for chronic health problems, is associated with increased mortality and exists in complexes of multiple, clustered behavioral risk factors. Similarly periodontal disease is one of the world's most common chronic diseases. Increasing evidence establishes periodontal disease as a significant risk factor in the etiology of diseases with inflammatory components. Severe periodontal disease is the well-established sixth complication of diabetes and the relationship may be two-way.⁵ Obesity is a systemic disease predisposing to comorbidities and complications that affect overall health; cross-sectional research suggests obesity is associated with periodontal diseases by underlying biologic mechanisms yet to be established.⁶

Obesity rates and its associated health problems have risen to exponential proportions in the United States (U.S.) and globally.¹ Obesity has been targeted as the sixth most significant risk factor worldwide that is known to contribute to both oral and other associated systemic diseases. According to the Centers for Disease Control and Prevention (CDC), the prevalence of obesity among adults in the U.S. is 40% and 18.5% in youth with prevalence higher among middle aged adults (42.8%) than younger adults (35.7%). As of 2016, nearly two billion people worldwide were either overweight or obese. Research indicates accumulated excessive fat from being overweight or obese contributes to serious health impairments.⁷

Objectives of the study

The main objective is to find the relations among obesity, diabetes mellitus and periodontitis in common people of Pakistan.

METHODOLOGY OF THE STUDY:

The study was conducted at Health department Punjab during 2019. There was 100 patients which was visit the health centre during this time period. We assess the level of glucose in those patients who was obsess and suffering from oral health issues. We measure the glucose level both before fasting and after fasting. For this purpose 5cc blood of patients were drawn for CBC analysis. The economic and health status describe the level of awareness regarding disease. The collected data were analyzed using SPSS software (version 17). The results are presented as a mean with 95% confidence interval limits or standard deviations. The significant value for $P < .05$ was accepted as statistically significant.

RESULTS:

In our data most of the respondents are female. The reason is that because female suffering more from obesity and diabetes. As compared to this male respondents has more normal values of glucose. Table 01 represents the level of blood glucose in patients. It shows that values of glucose become high diabetic patients as compared to normal values. IFG factor shows that diabetic people suffer more from oral health problems as compared to others.

Table 01: Level of glucose in obese and diabetic patients

	Normal	Diabetes	IFG	IGT
Fasting Plasma Glucose	<10	≥126	100-125	
Normal glucose		≥200 plus symptoms of diabetes		
Plasma glucose	<14	≥200		140-199

Table 02 shows the demographic history of patients. It clearly indicated that people who suffer from all above mentioned problems also suffer from high blood pressure and some other problems. Because people have less awareness of health issues.

Table 02: Demographic characteristics and history of patients

Variables	Co-efficient	SE
Blood pressure	0.048	0.35
Healthy eating index (HEI)	-0.059	0.05
Smoker	0.060	0.80
Food security	0.106	0.12
Drinker	-0.343	0.08
Belong to city area	0.057	0.01
Belong to rural area	0.59	0.70
BMI	0.5460.24	

Indicate significance at the 99, 95, and 90% level.

DISCUSSION:

This study focuses on the investigating the linkage between diabetes⁶, diet-health behavior, and health outcomes that are frequently discussed in the context of diabetes management, public health, and diet quality and BMI⁷. It is realized that carbohydrates are the supplements that most influence blood glucose levels.⁸ Be that as it may, up to now there is no agreement prove about the perfect measure of carbohydrate intake for individuals with diabetes. Truth be told, in the present investigation, the carbohydrate utilization did not vary between the unhealthy and healthy gathering.⁹

Body weight adjusted for stature (Body Mass Index) has been commonly used in large-scale population surveys as a surrogate for body fat content. Body mass index (BMI), also known as Quetelet's Index, is the most commonly used tool, the ratio defined as body weight (kg) divided by height squared (m²). BMI has been shown to have strong correlation with body fatness, and weak correlation with height.¹⁰ Some studies have suggested a relationship between marital status and obesity, although the relationship is not well established. Higher BMI has been associated with married subjects than subjects living alone. Other studies suggest that no such link exists.¹¹

Studies on diet and obesity have reported inconsistent results, compromised by factors such as weaknesses in study design, methodological errors in estimating energy and nutrient intakes, and confounding factors. Obesity has been suggested in association with a large number of various dietary factors but conclusive evidence is still lacking to prove than one diet would promote obesity more than another.¹²

CONCLUSION:

Obesity for many is a sensitive subject and may not be easy to bring up. Hence the need for more education. The demand for dental hygienists who specialize in treating patients with obesity may increase as the condition itself is increasing and the link with inflammatory diseases is relevant.

Understanding obesity may help the dental hygienist provide quality comprehensive care to those in need.

REFERENCES:

1. Dahiya P, Kamal R, Gupta R. Obesity, periodontal and general health: relationship and management. *Indian J Endocrinol Metab.* 2012 Jan;16(1):88-93.
2. Kangas S, Timonen P, Knuuttila M, Jula A, Ylostalo P, Syrjala AH. Waist circumference and waist-to-height ratio are associated with periodontal pocketing-results of the Health 2000 Survey. *BMC Oral Health.* 2017 Jan 17(48):1-7.
3. Brien S, Katzmarzyk P, Craig C, Gauvin L. Physical activity, cardiorespiratory fitness and body mass index as predictors of substantial weight gain and obesity: the Canadian physical activity longitudinal study. *Can J Public Health.* 2007 Mar/Apr;98(2):121-4.
4. Haas AN, Gaio EJ, Oppermann RV, Rosing CK, Albandar JM, Susin C. Pattern and rate of progression of periodontal attachment loss in an urban population of South Brazil: a 5-years population-based prospective study. *J Clin Periodontol.* 2012 Jan;39(1):1-9.
5. Eke PI, Dye BA, Wei L, Thornton-Evans GO, Genco RJ. Prevalence of periodontitis in adults in the United States: 2009 and 2010. *J Dent Res.* 2012 Oct;91(10):914-20.
6. Dietrich T, Sharma P, Walter C, Weston P, Beck J. The epidemiological evidence behind the association between periodontitis and incident atherosclerotic cardiovascular disease. *J Clin Periodontol.* 2013 Apr;40 Suppl 14:S70-84.
7. Muluke M, Gold T, Kiefhaber K, Al-Sahli A, Celenti R, Jiang H, Cremers S, Van Dyke T, Schulze-Spate U. Diet-Induced Obesity and Its Differential Impact on Periodontal Bone Loss. *J Dent Res.* 2016 Feb;95(2):223-9.
8. Khader YS, Bawadi HA, Haroun TF, Alomari M, Tayyem RF. The association between periodontal disease and obesity among adults in Jordan. *J Clin Periodontol.* 2009 Jan;36(1):18-24.
9. Zimmermann GS, Bastos MF, Dias Goncalves TE, Chambrone L, Duarte PM. Local and

- circulating levels of adipocytokines in obese and normal weight individuals with chronic periodontitis. *J Periodontol.* 2013 May;84(5):624-33.
10. Wheeler ML, Dunbar SA, Jaacks LM, Karmally W, Mayer-Davis EJ, Wylie-Rosett J, Yancy WS Jr. Macronutrients, food groups, and eating patterns in the management of diabetes: a systematic review of the literature, 2010. *Diabetes Care.* 2012;35(2):434–445.
 11. A jala O, English P, Pinkney J. Systematic review and meta-analysis of different dietary approaches to the management of type 2 diabetes. *Am J Clin Nutr.* 2013;97(3):505–516.
 12. Newby PK, Tucker KL. Empirically derived eating patterns using factor or cluster analysis: a review. *Nutr Rev.* 2004;62(5):177–203.